



CHEMISTRY AND PHYSICS

GEORGE TURRELL

$$y(x,0) = \sum_{n=1}^{\infty} B_n$$

Mathematics for Chemistry and Physics

This Page Intentionally Left Blank

Mathematics for Chemistry and Physics

GEORGE TURRELL

University of Science and Technology, Lille, France



An Elsevier Science Imprint

San Diego San Francisco New York Boston London Sydney Tokyo

This book is printed on acid-free paper.

Copyright © 2002 by ACADEMIC PRESS

All Rights Reserved.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher.

Academic Press An Elsevier Science Imprint Harcourt Place, 32 Jamestown Road, London NW1 7BY, UK http://www.academicpress.com

Academic Press

An Elsevier Science Imprint
525 B Street, Suite 1900, San Diego, California 92101-4495, USA http://www.academicpress.com

ISBN 0-12-705051-5

Library of Congress Catalog Number: 2001 091916

A catalogue record for this book is available from the British Library

Typeset by Laserwords Pvt. Ltd., Chennai, India Printed and bound in Great Britain by MPG Books Ltd, Bodmin, Cornwall

Contents

Preface			xiii
1	Varia	ables and Functions	1
	1.1	Introduction	1
	1.2	Functions	2
	1.3	Classification and properties of functions	6
	1.4	Exponential and logarithmic functions	7
	1.5	Applications of exponential and logarithmic	
		functions	10
	1.6	Complex numbers	12
	1.7	Circular trigonometric functions	14
	1.8	Hyperbolic functions	16
		Problems	17
2	Limit	ts, Derivatives and Series	19
	2.1	Definition of a limit	19
	2.2	Continuity	21
	2.3	The derivative	22
	2.4	Higher derivatives	24
	2.5	Implicit and parametric relations	25
	2.6	The extrema of a function and its critical points	26
	2.7	The differential	28
	2.8	The mean-value theorem and L'Hospital's rule	30
	2.9	Taylor's series	32
	2.10	Binomial expansion	34
	2.11	Tests of series convergence	35
	2.12	Functions of several variables	37
	2.13	Exact differentials	38
		Problems	39

vi CONTENTS

3	Integ	gration	43
	3.1	The indefinite integral	43
	3.2	Integration formulas	44
	3.3	Methods of integration	45
		3.3.1 Integration by substitution	45
		3.3.2 Integration by parts	46
		3.3.3 Integration of partial fractions	47
	3.4	Definite integrals	49
		3.4.1 Definition	49
		3.4.2 Plane area	50
		3.4.3 Line integrals	51
		3.4.4 Fido and his master	52
		3.4.5 The Gaussian and its moments	54
	3.5	Integrating factors	56
	3.6	Tables of integrals	59
		Problems	60
4	Vect	or Analysis	63
	4.1	Introduction	63
	4.2	Vector addition	64
	4.3	Scalar product	66
	4.4	Vector product	67
	4.5	Triple products	69
	4.6	Reciprocal bases	71
	4.7	Differentiation of vectors	72
	4.8	Scalar and vector fields	73
	4.9	The gradient	74
	4.10	The divergence	75
	4.11	The curl or rotation	75
	4.12	The Laplacian	76
	4.13	Maxwell's equations	77
	4.14	Line integrals	80
	4.15	Curvilinear coordinates	81
		Problems	83
5	Ordi	nary Differential Equations	85
	5.1	First-order differential equations	85
	5.2	Second-order differential equations	87
	5.4	5.2.1 Series solution	87 87
		J.2.1 Selies solution	0/

ONTENTS			vii
0.11.2.11.0			***
	ONTENTS	ONTENTS	ONTENTS

		5.2.2	The classical harmonic oscillator	89
		5.2.3	The damped oscillator	91
	5.3	The d	ifferential operator	93
		5.3.1	Harmonic oscillator	93
		5.3.2	Inhomogeneous equations	94
		5.3.3	Forced vibrations	95
	5.4	Applio	cations in quantum mechanics	96
		5.4.1	The particle in a box	96
		5.4.2	Symmetric box	99
		5.4.3	Rectangular barrier: The tunnel effect	100
		5.4.4	The harmonic oscillator in quantum	
			mechanics	102
	5.5	-	al functions	104
		5.5.1	Hermite polynomials	104
		5.5.2	Associated Legendre polynomials	107
		5.5.3	The associated Laguerre polynomials	111
		5.5.4	The gamma function	112
		5.5.5	Bessel functions	113
		5.5.6	Mathieu functions	114
		5.5.7	The hypergeometric functions	115
		Proble	ems	116
		–		
6	Part	iai Diffe	erential Equations	119
	6.1	The v	ibrating string	119
		6.1.1	The wave equation	119
		6.1.2	Separation of variables	120
		6.1.3	Boundary conditions	121
		6.1.4	Initial conditions	123
	6.2	The th	nree-dimensional harmonic oscillator	125
		6.2.1	Quantum-mechanical applications	125
		6.2.2	Degeneracy	127
	6.3		wo-body problem	129
		6.3.1	Classical mechanics	129
		6.3.2	Quantum mechanics	130
	6.4	Centra	al forces	132
		6.4.1	Spherical coordinates	132
		6.4.2	Spherical harmonics	134
	6.5	The d	iatomic molecule	135
		6.5.1	The rigid rotator	136

viii CONTENTS

		6.5.2 The vibrating rotator	136
		6.5.3 Centrifugal forces	137
	6.6	The hydrogen atom	138
		6.6.1 Energy	139
		6.6.2 Wavefunctions and the probability density	140
	6.7	Binary collisions	142
		6.7.1 Conservation of angular momentum	142
		6.7.2 Conservation of energy	143
		6.7.3 Interaction potential: LJ (6-12)	143
		6.7.4 Angle of deflection	145
		6.7.5 Quantum mechanical description:	
		The phase shift	146
		Problems	147
7	Oper	rators and Matrices	149
	7.1	The algebra of operators	149
	7.2	Hermitian operators and their eigenvalues	151
	7.3	Matrices	153
	7.4	The determinant	157
	7.5	Properties of determinants	158
	7.6	Jacobians	159
	7.7	Vectors and matrices	161
	7.8	Linear equations	163
	7.9	Partitioning of matrices	163
	7.10	Matrix formulation of the eigenvalue problem	164
	7.11	Coupled oscillators	166
	7.12	Geometric operations	170
	7.13	The matrix method in quantum mechanics	172
	7.14	The harmonic oscillator	175
		Problems	177
8	Grou	ıp Theory	181
	8.1	Definition of a group	181
	8.2	Examples	182
	8.3	Permutations	184
	8.4	Conjugate elements and classes	185
	8.5	Molecular symmetry	187
	8.6	The character	195
	8.7	Irreducible representations	196
	8.8	Character tables	198

CONTENTS ix

	8.9	Reduction of a representation:	
		The "magic formula"	200
	8.10	The direct product representation	202
	8.11	Symmetry-adapted functions: Projection operators	204
	8.12	Hybridization of atomic orbitals	207
	8.13	Crystal symmetry	209
		Problems	212
9	Mole	ecular Mechanics	215
	9.1	Kinetic energy	215
	9.2	Molecular rotation	217
	· · -	9.2.1 Euler's angles	218
		9.2.2 Classification of rotators	220
		9.2.3 Angular momenta	221
		9.2.4 The symmetric top in quantum mechanics	222
	9.3	Vibrational energy	224
	,	9.3.1 Kinetic energy	225
		9.3.2 Internal coordinates: The <i>G</i> matrix	226
		9.3.3 Potential energy	227
		9.3.4 Normal coordinates	227
		9.3.5 Secular determinant	228
		9.3.6 An example: The water molecule	229
		9.3.7 Symmetry coordinates	231
		9.3.8 Application to molecular vibrations	233
		9.3.9 Form of normal modes	234
	9.4	Nonrigid molecules	236
		9.4.1 Molecular inversion	236
		9.4.2 Internal rotation	238
		9.4.3 Molecular conformation: The molecular	
		mechanics method	240
		Problems	242
10	Prob	ability and Statistics	245
	10.1	Permutations	245
	10.1	Combinations	246
	10.2	Probability	249
	10.3	Stirling's approximation	251
	10.5	Statistical mechanics	253
	10.5	The Lagrange multipliers	255

x CONTENTS

	10.7	The partition function	256
	10.8	Molecular energies	257
		10.8.1 Translation	258
		10.8.2 Rotation	259
		10.8.3 Vibration	261
	10.9	Quantum statistics	262
		10.9.1 The indistinguishability of identical	
		particles	262
		10.9.2 The exclusion principle	263
		10.9.3 Fermi–Dirac statistics	264
		10.9.4 Bose–Einstein statistics	266
	10.10	Ortho- and para-hydrogen	267
		Problems	270
11	Integ	ral Transforms	271
	11.1	The Fourier transform	271
		11.1.1 Convolution	272
		11.1.2 Fourier transform pairs	273
	11.2	The Laplace transform	279
		11.2.1 Examples of simple Laplace transforms	279
		11.2.2 The transform of derivatives	281
		11.2.3 Solution of differential equations	282
		11.2.4 Laplace transforms: Convolution	
		and inversion	283
		11.2.5 Green's functions	284
		Problems	286
12	Appr	oximation Methods in Quantum Mechanics	287
	12.1	The Born-Oppenheimer approximation	287
	12.2	Perturbation theory: Stationary states	290
		12.2.1 Nondegenerate systems	290
		12.2.2 First-order approximation	291
		12.2.3 Second-order approximation	293
		12.2.4 The anharmonic oscillator	293
		12.2.5 Degenerate systems	296
		12.2.6 The Stark effect of the hydrogen atom	298
	12.3	Time-dependent perturbations	300
		12.3.1 The Schrödinger equation	300
		12.3.2 Interaction of light and matter	301

CONTENTS xi

		12.3.3 Spectroscopic selection rules	305
	12.4	The variation method	308
		12.4.1 The variation theorem	308
		12.4.2 An example: The particle in a box	309
		12.4.3 Linear variation functions	311
		12.4.4 Linear combinations of atomic	
		orbitals (LCAO)	312
		12.4.5 The Hückel approximation	316
		Problems	322
13	Num	erical Analysis	325
	13.1	Errors	325
	13.1	13.1.1 The Gaussian distribution	326
		13.1.2 The Poisson distribution	327
	13.2	The method of least squares	328
	13.3	Polynomial interpolation and smoothing	330
	13.4	The Fourier transform	334
	13.4	13.4.1 The discrete Fourier transform (DFT)	334
		13.4.2 The fast Fourier transform (FFT)	336
		13.4.3 An application: interpolation	330
		and smoothing	339
	13.5	Numerical integration	341
		13.5.1 The trapezoid rule	342
		13.5.2 Simpson's rule	343
		13.5.3 The method of Romberg	343
	13.6	Zeros of functions	345
		13.6.1 Newton's method	345
		13.6.2 The bisection method	346
		13.6.3 The roots: an example	346
		Problems	347
Арр	endic	es	
I	The C	Greek alphabet	349
II		nsions and units	351
III		ic orbitals	355
IV		ll wavefunctions for hydrogenlike species	361
V		Laplacian operator in spherical coordinates	363
VΙ		livergence theorem	367
VII		mination of the molecular symmetry group	369
,		minutes of the moreovalar symmetry group	20)

xii	C	ONTENTS

ΙX	common point groups
X	Further reading
	Chemical physics

Preface

This book has been written in an attempt to provide students with the mathematical basis of chemistry and physics. Many of the subjects chosen are those that I wish that I had known when I was a student. It was just at that time that the no-mans-land between these two domains – chemistry and physics – was established by the "Harvard School", certainly attributable to E. Bright Wilson, Jr., J. H. van Vleck and the others of that epoch. I was most honored to have been a product, at least indirectly, of that group as a graduate student of J. C. Decius. Later, in my post-doc years, I profited from the Harvard–MIT seminars. During this experience I listened to, and tried to understand, the presentations by those most prestigious persons, who played a very important role in my development in chemistry and physics. The essential books at that time were most certainly the many publications by John C. Slater and the "Bible" on mathematical methods, by Margeneau and Murphy. They were my inspirations.

The expression "Chemical Physics" appears to have been coined by Slater. I should like to quote from the preface to his book, "*Introduction to Chemical Physics*" (McGraw-Hill, New York, 1939).

It is probably unfortunate that physics and chemistry ever were separated. Chemistry is the science of atoms and of the way in which they combine. Physics deals with the interatomic forces and with the large-scale properties of matter resulting from those forces. So long as chemistry was largely empirical and nonmathematical, and physics had not learned how to treat small-scale atomic forces, the two sciences seemed widely separated. But with statistical mechanics and the kinetic theory on the one hand and physical chemistry on the other, the two sciences began to come together. Now [1939!] that statistical mechanics has led to quantum theory and wave mechanics, with its explanations of atomic interactions, there is really nothing separating them any more

A wide range of study is common to both subjects. The sooner we realize this the better. For want of a better name, as Physical Chemistry is already preempted, we may call this common field Chemical Physics. It is an overlapping field in which both physicists and chemists should xiv PREFACE

be trained. There seems no valid reason why their training in it should differ . . .

In the opinion of the present author, nobody could say it better.

That chemistry and physics are brought together by mathematics is the "raison d'être" of the present volume. The first three chapters are essentially a review of elementary calculus. After that there are three chapters devoted to differential equations and vector analysis. The remainder of the book is at a somewhat higher level. It is a presentation of group theory and some applications, approximation methods in quantum chemistry, integral transforms and numerical methods.

This is not a fundamental mathematics book, nor is it intended to serve a textbook for a specific course, but rather as a reference for students in chemistry and physics at all university levels. Although it is not computer-based, I have made many references to current applications – in particular to try to convince students that they should know more about what goes on behind the screen when they do one of their computer experiments. As an example, most students in the sciences now use a program for the fast Fourier transform. How many of them have any knowledge of the basic mathematics involved?

The lecture notes that I have written over many years in several countries have provided a basis for this book. More recently, I have distributed an early version to students at the third and fourth years at the University of Lille. It has been well received and found to be very useful. I hope that in its present form the book will be equally of value to students throughout their university studies.

The help of Professor Daniel Couturier, the ASA (Association de Solidarité des Anciens de l'Université des Sciences et Technologies de Lille) and the CRI (Centre de Resources Informatiques) in the preparation of this work is gratefully acknowledged. The many useful discussions of this project with Dr A. Idrissi, Dr F. Sokolić, Dr R. Withnall, Prof. M. Walters, Prof. D. W. Robinson and Prof. L. A. Veguillia-Berdicía are much appreciated.

My wife, Irène, and I have nicknamed this book "Mathieu". Throughout its preparation Irène has always provided encouragement – and patience when Mathieu was a bit trying or "Miss Mac" was in her more stubborn moods.

George Turrell Lille, May 1, 2001

1 Variables and Functions

1.1 INTRODUCTION

The usual whole numbers, integers such as 1, 2, 3, 4..., are usually referred to as Arabic numerals. It seems, however, that the basic decimal counting system was first developed in India, as it was demonstrated in an Indian astronomic calendar which dates from the third century AD. This system, which was composed of nine figures and the zero, was employed by the Arabs in the ninth century. The notation is basically that of the Arabic language and it was the Arabs who introduced the system in Europe at the beginning of the eleventh century.

In Europe the notion of the zero evolved slowly in various forms. Eventually, probably to express debts, it was found necessary to invent negative integers. The requirements of trade and commerce lead to the use of fractions, as ratios of whole numbers. However, it is obviously more convenient to express fractions in the form of decimals. The ensemble of whole numbers and fractions (as ratios of whole numbers) is referred to as rational numbers. The mathematical relation between decimal and rational fractions is of importance, particularly in modern computer applications.

As an example, consider the decimal fraction $x = 0.616161 \cdots$. Multiplication by 100 yields the expression $100 x = 61.6161 \cdots = 61 + x$ and thus, x = 61/99, is a rational fraction. In general, if a decimal expression contains an infinitely repeating set of digits (61 in this example), it is a rational number. However, most decimal fractions do not contain a repeating set of digits and so are not rational numbers. Examples such as $\sqrt{3} = 1.732051 \cdots$ and $\pi = 3.1415926536 \cdots$ are irrational numbers.* Furthermore, the logarithms and trigonometric functions of most arguments are irrational numbers.

^{*}A mnemonic for π based on the number of letters in words of the English language is quoted here from "the Green Book", Ian Mills, *et al.* (eds), "Quantities, Units and Symbols in Physical Chemistry", Blackwell Scientific Publications, London (1993):

^{&#}x27;How I like a drink, alcoholic of course, after the heavy lectures involving quantum mechanics!'

In practice, in numerical calculations with a computer, both rational and irrational numbers are represented by a finite number of digits. In both cases, then, approximations are made and the errors introduced in the result depend on the number of significant figures carried by the computer – the machine precision.* In the case of irrational numbers such errors cannot be avoided.

The ensemble of rational and irrational numbers are called real numbers. Clearly, the sum, difference and product of two real numbers is real. The division of two real numbers is defined in all cases but one – division by zero. Your computer will spit out an error message if you try to divide by zero!

1.2 FUNCTIONS

If two real variables are related such that, if a value of x is given, a value of y is determined, y is said to be a function of x. Thus, values may be assigned to x, the independent variable, leading to corresponding values of y, the dependent variable.

As an example, consider one mole of a gas at constant temperature. The volume V is a function of the applied pressure P. This relation can be expressed mathematically in the form

$$V = f(P), \tag{1}$$

or, V = V(P). Note that to complete the functional relationship, the nature of the gas, as well as the temperature T, must be specified. A physical chemist should also insist that the system be in thermodynamic equilibrium.

In the case of an ideal gas, the functional relationship of Eq. (1) becomes

$$V = \frac{C(T)}{P},\tag{2}$$

where C(T) is a positive constant which is proportional to the absolute temperature T. Clearly, the roles of V and P can be reversed leading to the relation

$$P = \frac{C(T)}{V}. (3)$$

The question as to which is the independent variable and which is the dependent one is determined by the way in which the measurements are made and, mathematically, on the presentation of the experimental data.

^{*}Note that the zero is a special case, as its precision is not defined. Normally, the computer automatically uses the precision specified for other numbers.

Suppose that a series of measurements of the volume of a gas is made, as the applied pressure is varied. As an example, the original results obtained by Boyle* are presented as in Table 1.

In this case V is a function of P, but it is not continuous. It is the discrete function represented by the points shown in Fig. 1a. It is only the mathematical function of Eq. (2) that is continuous. If, from the experimental data, it is of interest to calculate values of V at intermediate points, it is necessary to estimate them with the use of, say, linear interpolation, or better, a curve-fitting procedure. In the latter case the continuous function represented by Eq. (2)

	~	-
V, Volume ^a	P, Pressure (inches Hg)	1/V
12	29.125	0.0833
10	35.3125	0.1
8	44.1875	0.125
6	58.8125	0.1667
5	70.6875	0.2
4	87.875	0.25
3	116.5625	0.3333

Table 1 Volume of a gas as a function of pressure.

^aMeasured distance (inches) in a tube of constant diameter.




Fig. 1 Volume of a gas (expressed as distance in a tube of constant diameter) *versus* pressure (a). Pressure as a function of reciprocal volume (b).

^{*}Robert Boyle, Irish physical chemist (1627–1691)

would normally be employed. These questions, which concern the numerical treatment of data, will be considered in Chapter 13.

In Boyle's work the pressure was subsequently plotted as a function of the reciprocal of the volume, as calculated here in the third column of Table 1. The graph of P vs. 1/V is shown in Fig. 1b. This result provided convincing evidence of the relation given by Eq. (3), the mathematical statement of Boyle's law. Clearly, the slope of the straight line given in Fig. 1b yields a value of C(T) at the temperature of the measurements [Eq. (3)] and hence a value of the gas constant R. However, the significance of the temperature was not understood at the time of Boyle's observations.

In many cases a series of experimental results are not associated with a known mathematical function. In the following example Miss X weighed herself each morning beginning on the first of February. These data are presented graphically as shown in Fig. 2. Here interpolated points are of no significance, nor is extrapolation. By extrapolation Miss X would weigh nearly nothing in a year-and-a-half or so. However, as the data do exhibit a trend over a relatively short time, it is useful to employ a curve-fitting procedure. In this example Miss X might be happy to conclude that on the average she lost 0.83 kg per week during this period, as indicated by the slope of the straight line in Fig. 2.

Now reconsider the function given by Eq. (3). It has the form of a hyperbola, as shown in Fig. 3. Different values of C(T) lead to other members of the family of curves shown. It should be noted that this function is antisymmetric with respect to the inversion operation $V \to -V$ (see Chapter 8). Thus, P is said to be an odd function of V, as P(V) = -P(-V).

It should be evident that the negative branches of *P vs. V* shown in Fig. 3 can be excluded. These branches of the function are correct mathematically,




Fig. 2 Miss X's weight as a function of the date in February. The straight line is obtained by a least-squares fit to the experimental data (see Chapter 13).




Fig. 3 Pressure *versus* volume [Eq. (3)], with $C_3(T) > C_2(T) > C_1(T)$.

but are of no physical significance for this problem. This example illustrates the fact that functions may often be limited to a certain domain of acceptability. Finally, it should be noted that the function P(V) presented in Fig. 3 is not continuous at the origin (V=0). Therefore, from a physical point of view the function is only significant in the region $0 < V < \infty$. Furthermore, physical chemists know that Eqs. (2) and (3) do not apply at high pressures because the gas is no longer ideal.

As C(T) is a positive quantity, Eq. (3) can be written in the form

$$ln P = ln C - ln V. (4)$$

Clearly a plot of $\ln P$ vs. $\ln V$ at a given (constant) temperature yields a straight line with an intercept equal to $\ln C$. This analysis provides a convenient graphical method of determining the constant C.

It is often useful to shift the origin of a given graph. Thus, for the example given above consider that the axes of V and P are displaced by the amounts v and p, respectively. Then, Eq. (3) becomes

$$P - p = \frac{C(T)}{V - v},\tag{5}$$

and the result is as plotted in Fig. 4. The general hyperbolic form of the curves has not been changed, although the resulting function P(V) is no longer odd – nor is it even.




Fig. 4 Plots of Eq. (5) for given values of v and p.

1.3 CLASSIFICATION AND PROPERTIES OF FUNCTIONS

Functions can be classified as either algebraic or transcendental. Algebraic functions are rational integral functions or polynomials, rational fractions or quotients of polynomials, and irrational functions. Some of the simplest in the last category are those formed from rational functions by the extraction of roots. The more elementary transcendental functions are exponentials, logarithms, trigonometric and inverse trigonometric functions. Examples of these functions will be discussed in the following sections.

When the relation y = f(x) is such that there is only one value of y for each acceptable value of x, f(x) is said to be a single-valued function of x. Thus, if the function is defined for, say, $x = x_1$, the vertical line $x = x_1$ intercepts the curve at one and only one point, as shown in Fig. 5. However, in many cases a given value of x determines two or more distinct values of y.




Fig. 5 Plot of $y = x^2$.




Fig. 6 Plot of $x = \pm \sqrt{y}$.

The curve shown in Fig. 5 can be represented by

$$y = x^2, (6)$$

where y has the form of the potential function for a harmonic oscillator (see Chapter 5). This function is an even function of x, as y(x) = y(-x). Clearly, y is a single-valued function of x. Now, if Eq. (6) is rewritten in the equivalent form

$$x^2 = y, \quad y \ge 0 \tag{7}$$

it defines a double-valued function whose branches are given by $x = \sqrt{y}$ and $x = -\sqrt{y}$. These branches are the upper and lower halves of the parabola shown in Fig. 6. It should be evident from this example that to obtain a given value of x, it is essential to specify the particular branch of the (in general) multiple-valued function involved. This problem is particularly important in numerical applications, as carried out on a computer (Don't let the computer choose the wrong branch!).

1.4 EXPONENTIAL AND LOGARITHMIC FUNCTIONS

If
$$y = f(x)$$
 is given by
$$y = a^x$$
 (8)

y = f(x) is an exponential function. The independent variable x is said to be the argument of f. The inverse relation, the logarithm, can then be defined by

$$x = \log_a y \tag{9}$$

and a is called the base of the logarithm. It is clear, then, that $log_a a = 1$ and $log_a 1 = 0$. The logarithm is a function that can take on different values