


TEXTBOOKS IN MATHEMATICS

A Course in Differential Equations with Boundary-Value Problems SECOND EDITION



Stephen A. Wirkus
Randall J. Swift
Ryan S. Szypowski



CRC Press
Taylor & Francis Group

A CHAPMAN & HALL BOOK

**A COURSE IN
DIFFERENTIAL
EQUATIONS WITH
BOUNDARY-VALUE
PROBLEMS**

SECOND EDITION

TEXTBOOKS in MATHEMATICS

Series Editors: Al Boggess and Ken Rosen

PUBLISHED TITLES

ABSTRACT ALGEBRA: A GENTLE INTRODUCTION

Gary L. Mullen and James A. Sellers

ABSTRACT ALGEBRA: AN INTERACTIVE APPROACH, SECOND EDITION

William Paulsen

ABSTRACT ALGEBRA: AN INQUIRY-BASED APPROACH

Jonathan K. Hodge, Steven Schlicker, and Ted Sundstrom

ADVANCED LINEAR ALGEBRA

Hugo Woerdeman

ADVANCED LINEAR ALGEBRA

Nicholas Loehr

ADVANCED LINEAR ALGEBRA, SECOND EDITION

Bruce Cooperstein

APPLIED ABSTRACT ALGEBRA WITH MAPLETM AND MATLAB[®], THIRD EDITION

Richard Klima, Neil Sigmon, and Ernest Stitzinger

APPLIED DIFFERENTIAL EQUATIONS: THE PRIMARY COURSE

Vladimir Dobrushkin

A BRIDGE TO HIGHER MATHEMATICS

Valentin Deaconu and Donald C. Pfaff

COMPUTATIONAL MATHEMATICS: MODELS, METHODS, AND ANALYSIS WITH MATLAB[®] AND MPI,
SECOND EDITION

Robert E. White

A COURSE IN ORDINARY DIFFERENTIAL EQUATIONS, SECOND EDITION

Stephen A. Wirkus and Randall J. Swift

DIFFERENTIAL EQUATIONS: THEORY, TECHNIQUE, AND PRACTICE, SECOND EDITION

Steven G. Krantz

DIFFERENTIAL EQUATIONS: THEORY, TECHNIQUE, AND PRACTICE WITH BOUNDARY VALUE PROBLEMS

Steven G. Krantz

PUBLISHED TITLES CONTINUED

DIFFERENTIAL EQUATIONS WITH APPLICATIONS AND HISTORICAL NOTES, THIRD EDITION

George F. Simmons

DIFFERENTIAL EQUATIONS WITH MATLAB®: EXPLORATION, APPLICATIONS, AND THEORY

Mark A. McKibben and Micah D. Webster

DISCOVERING GROUP THEORY: A TRANSITION TO ADVANCED MATHEMATICS

Tony Barnard and Hugh Neill

ELEMENTARY NUMBER THEORY

James S. Kraft and Lawrence C. Washington

EXPLORING CALCULUS: LABS AND PROJECTS WITH MATHEMATICA®

Crista Arangala and Karen A. Yokley

EXPLORING GEOMETRY, SECOND EDITION

Michael Hvidsten

EXPLORING LINEAR ALGEBRA: LABS AND PROJECTS WITH MATHEMATICA®

Crista Arangala

EXPLORING THE INFINITE: AN INTRODUCTION TO PROOF AND ANALYSIS

Jennifer Brooks

GRAPHS & DIGRAPHS, SIXTH EDITION

Gary Chartrand, Linda Lesniak, and Ping Zhang

INTRODUCTION TO ABSTRACT ALGEBRA, SECOND EDITION

Jonathan D. H. Smith

INTRODUCTION TO ANALYSIS

Corey M. Dunn

INTRODUCTION TO MATHEMATICAL PROOFS: A TRANSITION TO ADVANCED MATHEMATICS, SECOND EDITION

Charles E. Roberts, Jr.

INTRODUCTION TO NUMBER THEORY, SECOND EDITION

Marty Erickson, Anthony Vazzana, and David Garth

LINEAR ALGEBRA, GEOMETRY AND TRANSFORMATION

Bruce Solomon

MATHEMATICAL MODELLING WITH CASE STUDIES: USING MAPLE™ AND MATLAB®, THIRD EDITION

B. Barnes and G. R. Fulford

MATHEMATICS IN GAMES, SPORTS, AND GAMBLING—THE GAMES PEOPLE PLAY, SECOND EDITION

Ronald J. Gould

PUBLISHED TITLES CONTINUED

THE MATHEMATICS OF GAMES: AN INTRODUCTION TO PROBABILITY

David G. Taylor

A MATLAB® COMPANION TO COMPLEX VARIABLES

A. David Wunsch

MEASURE AND INTEGRAL: AN INTRODUCTION TO REAL ANALYSIS, SECOND EDITION

Richard L. Wheeden

MEASURE THEORY AND FINE PROPERTIES OF FUNCTIONS, REVISED EDITION

Lawrence C. Evans and Ronald F. Gariepy

NUMERICAL ANALYSIS FOR ENGINEERS: METHODS AND APPLICATIONS, SECOND EDITION

Bilal Ayyub and Richard H. McCuen

ORDINARY DIFFERENTIAL EQUATIONS: AN INTRODUCTION TO THE FUNDAMENTALS

Kenneth B. Howell

PRINCIPLES OF FOURIER ANALYSIS, SECOND EDITION

Kenneth B. Howell

REAL ANALYSIS AND FOUNDATIONS, FOURTH EDITION

Steven G. Krantz

RISK ANALYSIS IN ENGINEERING AND ECONOMICS, SECOND EDITION

Bilal M. Ayyub

SPORTS MATH: AN INTRODUCTORY COURSE IN THE MATHEMATICS OF SPORTS SCIENCE AND
SPORTS ANALYTICS

Roland B. Minton

TRANSFORMATIONAL PLANE GEOMETRY

Ronald N. Umble and Zhigang Han

TEXTBOOKS in MATHEMATICS

A COURSE IN DIFFERENTIAL EQUATIONS WITH BOUNDARY-VALUE PROBLEMS

SECOND EDITION

Stephen A. Wirkus

Arizona State University
Glendale, USA

Randall J. Swift

California State Polytechnic University
Pomona, USA

Ryan S. Szypowski

California State Polytechnic University
Pomona, USA



CRC Press

Taylor & Francis Group
Boca Raton London New York

CRC Press is an imprint of the
Taylor & Francis Group, an **informa** business
A CHAPMAN & HALL BOOK

MATLAB® is a trademark of The MathWorks, Inc. and is used with permission. The MathWorks does not warrant the accuracy of the text or exercises in this book. This book's use or discussion of MATLAB® software or related products does not constitute endorsement or sponsorship by The MathWorks of a particular pedagogical approach or particular use of the MATLAB® software.

CRC Press
Taylor & Francis Group
6000 Broken Sound Parkway NW, Suite 300
Boca Raton, FL 33487-2742

First issued in paperback 2022

© 2017 by Taylor & Francis Group, LLC
CRC Press is an imprint of Taylor & Francis Group, an Informa business

No claim to original U.S. Government works

ISBN 13: 978-1-03-247696-4 (pbk)
ISBN 13: 978-1-4987-3605-3 (hbk)
ISBN 13: 978-1-315-36961-7 (ebk)

DOI: 10.1201/9781315369617

This book contains information obtained from authentic and highly regarded sources. Reasonable efforts have been made to publish reliable data and information, but the author and publisher cannot assume responsibility for the validity of all materials or the consequences of their use. The authors and publishers have attempted to trace the copyright holders of all material reproduced in this publication and apologize to copyright holders if permission to publish in this form has not been obtained. If any copyright material has not been acknowledged please write and let us know so we may rectify in any future reprint.

Except as permitted under U.S. Copyright Law, no part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, please access www.copyright.com (<http://www.copyright.com/>) or contact the Copyright Clearance Center, Inc. (CCC), 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. CCC is a not-for-profit organization that provides licenses and registration for a variety of users. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

Trademark Notice: Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation without intent to infringe.

Publisher's Note

The publisher has gone to great lengths to ensure the quality of this reprint but points out that some imperfections in the original copies may be apparent.

Library of Congress Cataloging-in-Publication Data

Names: Wirkus, Stephen Allen, 1971- | Swift, Randall J. | Szypowski, Ryan. |
Wirkus, Stephen Allen, 1971- A course in ordinary differential equations.
Title: A course in differential equations with boundary value problems /
Stephen A. Wirkus, Randall J. Swift, and Ryan Szypowski.
Other titles: Course in ordinary differential equations
Description: Second edition. | Boca Raton : CRC Press, 2017. | Previous
title: A course in ordinary differential equations / Stephen A. Wirkus,
Randall J. Swift (Boca Raton : CRC Press, 2015). | Includes
bibliographical references and index.
Identifiers: LCCN 2016034689 | ISBN 9781498736053
Subjects: LCSH: Differential equations. | Differential equations, Linear. |
Boundary value problems.
Classification: LCC QA372 .S9254 2015 | DDC 515/.352--dc23
LC record available at <https://lccn.loc.gov/2016034689>

Visit the Taylor & Francis Web site at
<http://www.taylorandfrancis.com>

and the CRC Press Web site at
<http://www.crcpress.com>

To our families

Erika Tatiana, Alan, Abdi, and Avani,

Kelly, Kaelin, Robyn, Erin, and Ryley,

and

Jillian

for bringing us more joy than math and showing us the true concept

and meaning of

∞ infinity ∞

with their tireless

patience, love, and understanding.



Taylor & Francis

Taylor & Francis Group

<http://taylorandfrancis.com>

Contents

About the Authors	xiii
Preface	xv
1 Traditional First-Order Differential Equations	1
1.1 Introduction to First-Order Equations	1
1.2 Separable Differential Equations	7
1.3 Linear Equations	13
1.4 Some Physical Models Arising as Separable Equations	24
1.5 Exact Equations	33
1.6 Special Integrating Factors and Substitution Methods	43
1.6.1 Bernoulli Equation	45
1.6.2 Homogeneous Equations of the Form $g(y/x)$	47
Chapter 1 Review	54
Computer Labs: MATLAB®, Maple™, <i>Mathematica</i>	55
Chapter 1 Projects	66
Project 1A: Particles in the Atmosphere	66
Project 1B: Insights into Graphing	67
2 Geometrical and Numerical Methods for First-Order Equations	69
2.1 Direction Fields—Geometry of Differential Equations	69
2.2 Existence and Uniqueness for First-Order Equations	73
2.3 First-Order Autonomous Equations—Geometrical Insight	78
2.3.1 Graphing Factored Polynomials	85
2.3.2 Bifurcations of Equilibria	88
2.4 Modeling in Population Biology	91
2.4.1 Nondimensionalization	94
2.5 Numerical Approximation: Euler and Runge–Kutta Methods	99
2.6 Introduction to Autonomous Second-Order Equations	107
Chapter 2 Review	112
Computer Labs: MATLAB, Maple, <i>Mathematica</i>	114
Chapter 2 Projects	127
Project 2A: Spruce Budworm	127
Project 2B: Multistep Methods of Numerical Approximation	127
3 Elements of Higher-Order Linear Equations	129
3.1 Introduction to Higher-Order Equations	129
3.1.1 Operator Notation	133
3.2 Linear Independence and the Wronskian	137
3.3 Reduction of Order—the Case $n = 2$	145
3.4 Numerical Considerations for n th-Order Equations	150
3.5 Essential Topics from Complex Variables	153

3.6	Homogeneous Equations with Constant Coefficients	160
3.7	Mechanical and Electrical Vibrations	169
	Chapter 3 Review	183
	Computer Labs: MATLAB, Maple, <i>Mathematica</i>	185
	Chapter 3 Projects	198
	Project 3A: Runge–Kutta Order 2	198
	Project 3B: Stiff Differential Equations	199
4	Techniques of Nonhomogeneous Higher-Order Linear Equations	201
4.1	Nonhomogeneous Equations	201
4.2	Method of Undetermined Coefficients via Superposition	210
4.3	Method of Undetermined Coefficients via Annihilation	221
4.4	Exponential Response and Complex Replacement	230
4.5	Variation of Parameters	239
4.6	Cauchy–Euler (Equidimensional) Equation	248
4.7	Forced Vibrations	252
	Chapter 4 Review	257
	Computer Labs: MATLAB, Maple, <i>Mathematica</i>	259
	Chapter 4 Projects	267
	Project 4A: Forced Duffing Equation	267
	Project 4B: Forced van der Pol Oscillator	268
5	Fundamentals of Systems of Differential Equations	269
5.1	Useful Terminology	269
5.2	Gaussian Elimination	277
5.3	Vector Spaces and Subspaces	283
	5.3.1 The Nullspace and Column Space	287
5.4	Eigenvalues and Eigenvectors	292
5.5	A General Method, Part I: Solving Systems with Real and Distinct or Complex Eigenvalues	302
5.6	A General Method, Part II: Solving Systems with Repeated Real Eigenvalues	308
5.7	Matrix Exponentials	319
5.8	Solving Linear Nonhomogeneous Systems of Equations	328
	Chapter 5 Review	334
	Computer Labs: MATLAB, Maple, <i>Mathematica</i>	338
	Chapter 5 Projects	350
	Project 5A: Transition Matrix and Stochastic Processes	350
	Project 5B: Signal Processing	352
6	Geometric Approaches and Applications of Systems of Differential Equations	353
6.1	An Introduction to the Phase Plane	353
6.2	Nonlinear Equations and Phase Plane Analysis	361
	6.2.1 Systems of More Than Two Equations	366
6.3	Bifurcations	370
6.4	Epidemiological Models	379
6.5	Models in Ecology	392
	Chapter 6 Review	400
	Computer Labs: MATLAB, Maple, <i>Mathematica</i>	402
	Chapter 6 Projects	415

Project 6A: An MSEIR Model	415
Project 6B: Routh–Hurwitz Criteria	416
7 Laplace Transforms	419
7.1 Introduction	419
7.2 Fundamentals of the Laplace Transform	427
7.3 The Inverse Laplace Transform	438
7.3.1 Laplace Transform Solution of Linear Differential Equations	442
7.4 Translated Functions, Delta Function, and Periodic Functions	448
7.5 The s -Domain and Poles	456
7.6 Solving Linear Systems Using Laplace Transforms	461
7.7 The Convolution	465
Chapter 7 Review	470
Computer Labs: MATLAB, Maple, <i>Mathematica</i>	472
Chapter 7 Projects	482
Project 7A: Carrier-Borne Epidemics	482
Project 7B: Integral Equations	483
8 Series Methods	487
8.1 Power Series Representations of Functions	487
8.2 The Power Series Method	497
8.3 Ordinary and Singular Points	505
8.4 The Method of Frobenius	513
8.5 Bessel Functions	529
Chapter 8 Review	541
Computer Labs: MATLAB, Maple, <i>Mathematica</i>	541
Chapter 8 Projects	549
Project 8A: Asymptotic Series	549
Project 8B: Hypergeometric Functions	550
9 Boundary-Value Problems and Fourier Series	553
9.1 Two-Point Boundary-Value Problems	553
9.2 Orthogonal Functions and Fourier Series	558
9.3 Even, Odd, and Discontinuous Functions	565
9.4 Simple Eigenvalue-Eigenfunction Problems	571
9.5 Sturm–Liouville Theory	575
9.6 Generalized Fourier Series	581
Computer Labs: MATLAB, Maple, <i>Mathematica</i>	586
Chapter 9 Projects	592
Project 9: Chebyshev Polynomials	592
10 Partial Differential Equations	593
10.1 Separable Linear Partial Differential Equations	593
10.2 Heat Equation	600
10.3 Wave Equation	609
10.4 Laplace Equation	620
10.5 Nonhomogeneous Boundary Conditions	630
10.6 Non-Cartesian Coordinate Systems	636
Computer Labs: MATLAB, Maple, <i>Mathematica</i>	645
Chapter 10 Projects	652
Project 10: Finite Difference Approximation	652

A	An Introduction to MATLAB[®], Maple[™], and <i>Mathematica</i>	653
A.1	MATLAB	653
A.1.1	Some Helpful MATLAB Commands	655
A.1.2	Programming with a script and a function in MATLAB	658
A.2	Maple	661
A.2.1	Some Helpful Maple Commands	662
A.2.2	Programming in Maple	664
A.3	<i>Mathematica</i>	666
A.3.1	Some Helpful <i>Mathematica</i> Commands	668
A.3.2	Programming in <i>Mathematica</i>	669
B	Selected Topics from Linear Algebra	673
B.1	A Primer on Matrix Algebra	673
B.2	Matrix Inverses, Cramer's Rule	685
B.2.1	Calculating the Inverse of a Matrix	685
B.2.2	Cramer's Rule	689
B.3	Linear Transformations	692
B.4	Coordinates and Change of Basis	701
B.4.1	Similarity Transformations	703
	Computer Labs: MATLAB, Maple, <i>Mathematica</i>	707
	Answers to Odd Problems	717
	References	755
	Index	759

About the Authors

Stephen A. Wirkus completed his Ph.D. at Cornell University under the direction of Richard Rand. He began guiding undergraduate research projects while in graduate school and came to California State Polytechnic University, Pomona (Cal Poly Pomona) in 2000 after being a Visiting Professor at Cornell for a year. He co-founded the Applied Mathematical Sciences Summer Institute (AMSSI), an undergraduate research program jointly hosted by Loyola Marymount University, that ran from 2005 through 2007. He came to Arizona State University in 2007 as a tenured associate professor and won the 2013 Professor of the Year Award at ASU as well as the 2011 NSF AGEP Mentor of the Year award. He was a Visiting MLK Professor at the Massachusetts Institute of Technology in 2013–2014. He has guided over 80 undergraduate students in research and has served as chair for four M.S. students, and two Ph.D. students. He has over 30 publications and technical reports with over 40 students and has received grants from the NSF and NSA for guiding undergraduate research.

Randall J. Swift completed his Ph.D. at the University of California, Riverside under the direction of M. M. Rao. He began his career at Western Kentucky University and taught there for nearly a decade before moving to Cal Poly Pomona in 2001 as a tenured associate professor. He is active in research and teaching, having authored more than 80 journal articles, three research monographs and three textbooks in addition to serving as chair for 25 M.S. students. Now a professor, he received the 2011–12 Ralph W. Ames Distinguished Research Award from the College of Science at Cal Poly Pomona. The award honors Swift for his outstanding research in both pure and applied mathematics, and his contributions to the mathematics field as a speaker, journal editor, and principal investigator on numerous grants. He was also a visiting professor in 2007–2008 at the Australian National University in Canberra, Australia as well as having taught at the Claremont Colleges.

Ryan S. Szypowski completed his Ph.D. at the University of California, San Diego under the direction of Michael Holst. He was hired at Cal Poly Pomona in 2011 and became a tenured Associate Professor in 2016. His main research is in the area of computational methods for approximation of solutions to partial differential equations, but studies broad problems from applied mathematics and science with his students and colleagues. He had an active NSF research grant from 2012 to 2015 to study adaptive techniques in finite element exterior calculus and has won numerous smaller grants to support his work with students. In 2015, he won the Department of Mathematics and Statistics Excellence in Teaching Award.



Taylor & Francis

Taylor & Francis Group

<http://taylorandfrancis.com>

Preface

This book is based on lectures given by the first author at Cal Poly Pomona, Arizona State University (ASU), and the Massachusetts Institute of Technology (MIT), the second author at Western Kentucky University (WKU) and Cal Poly Pomona, and the third author at Cal Poly Pomona. The first eight chapters and two appendices are identical to those in *A Course in Ordinary Differential Equations, 2nd Edition*. The text can be used for a traditional one-semester sophomore-level course in ordinary differential equations (such as WKU's MATH 331), a one-semester sophomore-level course in differential equations in which partial differential equations replace Laplace transforms, or a two-semester sophomore- or junior-level course in differential equations. There is ample material for a two-quarter sequence (such as Cal Poly Pomona's MAT 216-431 or MAT 431-432), as well as sufficient linear algebra in the text so that it can be used for a one-quarter course that combines differential equations and linear algebra (such as Cal Poly Pomona's Math 224), or a one-semester course in differential equations that brings in linear algebra in a significant way (such as ASU's MAT 275 or MIT's 18.03). Most significantly, computer labs are given in MATLAB[®],¹ Maple[™], and *Mathematica* at the end of each chapter so the book may be used for a course to introduce and equip the student with a knowledge of the given software (such as ASU's MAT 275). Near the end of this Preface, we give some sample course outlines that will help show the independence of various sections and chapters. The focus of the text is on applications and methods of solution, both analytical and numerical, with emphasis on methods used in the typical engineering, physics, or mathematics student's field of study. We have tried to provide sufficient problems of a mathematical nature at the end of each section so that even the pure math major will be sufficiently challenged.

Key Features

This second edition of the book keeps many of the key features from the first edition:

- MATLAB, Maple, and *Mathematica* are incorporated at the end of each chapter, helping students with pages of tedious algebra and many of the differential equations topics; the goal of the software is still to show students how to make informed use of the relevant software in the field; all three software packages have parallel code and exercises;
- There are numerous problems of varying difficulty for both the applied and pure math major, as well as problems for the nonmathematician (engineers, etc.);
- An appendix that gives the reader a “crash course” in the three software packages; no prior knowledge is assumed;
- Chapter reviews at the end of each chapter to help the students review;
- Projects at the end of each chapter that go into detail about certain topics and sometimes introduce new topics that the students are now ready to see;
- Answers to most of the odd problems in the back of the book;

¹MATLAB is a registered trademark of The MathWorks, Inc. For product information, please contact:
The Mathworks, Inc.
3 Apple Hill Drive
Natick, MA, 01760-2098 USA
Tel: 508-647-7000
Fax: 508-647-7001
E-mail: info@mathworks.com
Web: www.mathworks.com

- An appendix on linear algebra to supplement the treatment within the text, should it be appropriate for the reader/course;
- A full solutions manual for the qualified instructor.

It also incorporates new features, many of which have been suggested by professors and students who have taught/learned from the first edition:

- The computer codes are moved to the end of each chapter as **Computer Labs** to facilitate reading of the book by students and professors who either choose not to use the technology or who do not have access to it immediately;
- The latest software versions are used; significant changes have occurred in certain aspects of MATLAB, Maple, and *Mathematica* since the first edition in 2006, and the relevant changes are incorporated;
- Much of the linear algebra discussion has been moved to Chapter 5 (from Chapter 3), which deals with linear systems;
- Sections have been added on complex variables (Chapter 3), the exponential response formula for solving nonhomogeneous equations (Chapter 4), forced vibrations (Chapter 4) as well as a subsection on nondimensionalization (Chapter 2), and a combining of the sections on Euler and Runge-Kutta methods (Chapter 2);
- Many rewritten sections highlight applications and modeling within many fields;
- Exercises flow from easiest to hardest;
- Color graphs to help the reader better understand crucial concepts in ordinary differential equations;
- Updated and extended projects at the end of each chapter to reflect changes within the chapters.

Approaches to Teaching Differential Equations

The second edition of this book has evolved with our understanding of how to teach the material in the best possible way. Some notable examples from the above list:

1. The structure of the course in covering linear systems in their entirety before covering applications to nonlinear systems (phase plane, etc.) was a direct result of numerous conversations with MIT's Professor Haynes Miller (who frequently teaches MIT's 18.03) as was the incorporation of the new sections on essential topics from complex variables, exponential response, and complex replacement (developed by Haynes) for solving nonhomogeneous differential equations, and the s -domain and poles as an important use of Laplace transforms by engineers.
2. Combining the computer codes into Computer Labs at the end of each section rather than having snippets of code embedded throughout the text was a direct result of a switch in ASU's method of teaching this course. Setting aside six class periods for such labs is the way differential equations is now taught at ASU.
3. The presentation of essential linear algebra topics to aid in the understanding of differential equations was helped by discussions with MIT's Professor Gil Strang as well as seeing some of his lectures firsthand.

Most differential equations we have encountered in practice have needed analytical approximations or numerical approximations to gain insight into their behavior. We don't feel that students use technology wisely if they simply ask the computer to solve a given problem. We thus focus on what we consider to be the basics necessary for adequately preparing a student for study in her or his respective fields, including mathematics. We present the syntax from MATLAB, Maple, and *Mathematica* in Computer Labs at the end of each chapter. We feel

that this provides the readers a better understanding of the theory and allows them to gain more insight into real-world problems they are likely to encounter. The vast majority of our students also have *no previous experience with MATLAB, Maple, or Mathematica* and we start from the basics and teach informed use of the relevant mathematical software. The student whom we “typically encounter” has had one year of calculus and is usually a major in a field other than pure mathematics.

Our book is traditional in its approach and coverage of basic topics in ordinary differential equations. However, we cover a number of “modern” topics such as direction fields, phase lines, the Runge–Kutta method, and nondimensionalization in Chapter 2 and epidemiological and ecological models in Chapter 6. As mentioned earlier, we also bring elements from linear algebra, such as eigenvectors, bases, and transformations, in order to best equip the reader of the book with a solid understanding of the material. Besides the Computer Labs there are also Projects at the end of each chapter that give useful insight into past and future topics covered in the book. The topics covered in these projects include a mix of traditional, modeling, numerical, and linear algebra aspects of ordinary differential equations. It is the intent that students who study this book and work *most* of the problems contained in these pages will be very prepared to continue their studies in engineering and mathematics.

Some Sample Course Outlines

While we could not begin to prescribe how this book may best be used for each school, we include some *possible* sections covered for various course outlines. There is sufficient material for a two-quarter or two-semester course sequence involving ordinary differential equations and partial differential equations with or without an emphasis on linear algebra that would utilize most of the book. We stress that if you intend to incorporate MATLAB, Maple, or *Mathematica* into your course, it is crucial to assign Exercises 1–4 (plus a few others) from Appendix A and the Chapter 1 Computer Lab early in the course. Appendix A requires only a knowledge of college algebra and some calculus (Taylor series), while Chapter 1 Computer Lab requires knowledge of calculus as it is applied to differential equations. Thus both can be assigned within the first 2 weeks of the course (and likely together).

Traditional semester ODE course:

Chap. 1	Chap. 2	Chap. 3	Chap. 4	Chap. 5	Chap. 7	Chap. 8
1.1–1.6	2.1–2.2	3.1–3.3 3.5–3.6	4.1, 4.3 4.5–4.6	5.1 5.4–5.8	7.1–7.4	8.1–8.5

Semester ODE course with modeling or application emphasis:

Chap. 1	Chap. 2	Chap. 3	Chap. 4	Chap. 5	Chap. 6	Chap. 7
1.1–1.4	2.1–2.6	3.1–3.2 3.4–3.7	4.1–4.2 4.4–4.7	5.1, 5.4 5.5, 5.7	6.1–6.5	7.1–7.5

Semester DE course with PDEs instead of Laplace Transforms:

Chap. 1	Chap. 2	Chap. 3	Chap. 4	Chap. 5	Chap. 6	Chap. 10
1.1–1.6	2.1–2.2	3.1–3.3 3.5–3.6	4.1, 4.3 4.5–4.6	5.1 5.4–5.8	6.1	10.1–10.6

Semester ODE course with linear algebra emphasis and no separate computer labs:

Ch. 1	Ch. 2	Ch. 3	Ch. 4	Ch. 5	Ch. 6	Ch. 7	App. B
1.1–1.4	2.1–2.2 2.5	3.1–3.2 3.4–3.7	4.1–4.2 4.4, 4.7	5.1–5.5 5.7–5.8	6.1	7.1–7.7	B.1–B.4

Semester DE course with linear algebra emphasis and no separate computer labs:

Ch. 1	Ch. 2	Ch. 3	Ch. 4	Ch. 5	Ch. 6	Ch. 10	App. B
1.1–1.4	2.1–2.2 2.5	3.1–3.2 3.4–3.7	4.1–4.2 4.4, 4.7	5.1–5.5 5.7–5.8	6.1	10.1–10.6	B.1–B.4

Semester ODE course with linear algebra emphasis and 6 computer labs:

Ch. 1	Ch. 2	Ch. 3	Ch. 4	Ch. 5	Ch. 7	Comp. Labs
1.1–1.4	2.1–2.2 2.5	3.1–3.2 3.4–3.7	4.1–4.2 4.4, 4.7	5.1–5.5	7.1–7.6	A & 1, 2, 3, 4, 5 & B, 7

Quarter ODE course with linear algebra emphasis:

Ch. 1	Ch. 2	Ch. 3	Ch. 4	Ch. 5	App. B
1.1–1.4	2.1–2.2 2.5	3.1–3.2 3.4–3.7	4.1–4.2 4.7	5.1–5.5	B.1–B.4

Acknowledgments

Students, with their questions both in class and during office hours, helped shaped this second edition as did those professors who used the first edition and/or provided constructive feedback to us, including Erika Camacho, Andrew Knyazev, Luis Melara, Jenny Switkes, Karen Vaughn, Steven Weintraub, and many others. Various chapters were read by Jonathan Burkow, Alexandra Churikova, Maytee Cruz-Aponte, Clay Goggil, Chris Graham, Christine Sowa, and Kathryn Stefanko, and their feedback has been of great help. Mike Pappas, in particular, was a big help in proofreading near-final drafts of several chapters. Valerie Cheathon provided a valuable check of all the codes as did Joshua Grosso (MATLAB) and Alan Wirkus-Camacho (Maple and Mathematica). Scott Wilde, again, provided invaluable help in revising the solutions manual.

As texts based upon lecture notes seemingly develop, many of the examples, exercises, and projects have been collected over many years for various courses taught by the authors. Some were taken from others' textbooks and papers. We have tried to give proper credit throughout this text; however, it was not always possible to properly acknowledge the original sources. It is our hope that we repay this explicit debt to earlier writers by contributing our (and their) ideas to further student understanding of differential equations.

We particularly wish to thank our project editor, Karen Simon, the production coordinator for the ODE version of the book, Jessica Vakili, as well as Michele Dimont, Amy Blalock, Hayley Ruggieri, and Sherry Thomas. Bob Stern and Bob Ross, our editors at Chapman & Hall/CRC Press, both deserve a big thanks for believing in this project and for helpful guidance, advice, and patience. We sincerely thank all these individuals; without their assistance, this text would not have succeeded.

URL for typos and errata:

<http://www.public.asu.edu/~swirkus/ACourseInDEs>

Finally, we would appreciate any comments that you might have regarding this book.

Stephen A. Wirkus (e-mail: swirkus@asu.edu)

Randall J. Swift (e-mail: rjswift@cpp.edu)

Ryan Szypowski (e-mail: rsszypowski@cpp.edu)

From the first edition:

We owe a very special thanks to Erika Camacho (Arizona State University) for her help in writing the MATLAB and Maple code for this book and for detailed suggestions on numerous sections. John Fay and Gary Etgen reviewed earlier drafts of this text and provided helpful feedback. Scott Wilde provided valuable assistance in writing and preparing the solutions manual for the book. We owe a big thanks to our former students David Monarres, for help in preparing portions of this book, and Walter Sosa and Moore Chung, for their help in preparing solutions. We would also like to acknowledge our Cal Poly Pomona colleagues Michael Green, Jack Hofer, Tracy McDonald, Jim McKinney, Siew-Ching Pye, Dick Robertson, Paul Salomaa, Jenny Switkes, Karen Vaughn, and Mason Porter (Caltech/Oxford) for their willingness to use draft versions of this text in their courses and their important suggestions, which improved the overall readability of the text. The faculty and students of AMSSI and MTBI also deserve a special thanks for comments on early drafts of the computer code. Mary Jane Hill assisted us with certain aspects of the text and helped in typesetting some of the chapters of the initial drafts of the book; her effort is greatly appreciated. The production and support staff at Chapman & Hall/CRC Press have been very helpful. We particularly wish to thank our project coordinator, Theresa Del Forn and project editor, Prudence Board. Our editor Bob Stern deserves a special thanks for believing in this project and for his guidance, advice, and patience. We sincerely thank all these individuals; without their assistance this text would not have succeeded.

A few remarks for students and professors:

This book will succeed if any fears and reservations about learning one of the three computer algebra systems used in this book are put aside. Computers are not here to supplant us, but rather they are here to help illustrate and illuminate concepts and insights that we have. Nothing is foolproof and we stress the importance of *informed use of the relevant mathematical software*. Numerical answers, although quite accurate most of the time, should always be examined carefully because computers are as smart as the programmer allows them to be. There should never be a blind trust in an answer.

It is essential that the technology that you choose—MATLAB, Maple, or Mathematica—be introduced early in the class, just as it is introduced early in the book. While certain mathematical software packages may be better suited for studying differential equations, none have the versatility that the above three programs have to give insight into other areas of mathematics. The two keys to learning the programs are (1) learning the syntax and (2) learning to use the help menus to figure out some of the commands. Setting aside one class, for example, to give a brief tutorial on one of these software packages in the computer lab is a very worthwhile investment. It is by no means necessary and the typical student will be able to learn the material on his/her own by carefully following Appendix A. For reinforcement, it is crucial to include at least one or two technology problems with each homework assignment. The conscientious student will be well prepared to use the same software package in any upper division course in *any* branch of the mathematical sciences and its applications.

It is not necessary to bring computer demonstrations into the classroom. Both authors have taught their courses successfully without classroom demonstrations; handouts sometimes are useful, especially from the appendices. The students, for better or worse, are generally far less afraid of technology than one might expect. If students are sent to the computer lab with an assignment to do and aided with Appendix A, the vast majority will come back with satisfactory answers. Yes, you may bang your head against your desk in frustration at times, but just ask the person next to you for help and also seek the help menus and you will be able to learn MATLAB, Maple, and *Mathematica* quite well.



Taylor & Francis

Taylor & Francis Group

<http://taylorandfrancis.com>

Chapter 1

Traditional First-Order Differential Equations

A Very Brief History

The study of Differential Equations began very soon after the invention of Differential and Integral Calculus, to which it forms a natural sequel. In 1676 Newton solved a differential equation by the use of an infinite series, only 11 years after his discovery of the *fluxional* form of differential calculus in 1665. These results were not published until 1693, the same year in which a differential equation occurred for the first time in the work of Leibniz (whose account of differential calculus was published in 1684).

In the next few years progress was rapid. In 1694–1697 John Bernoulli explained the method of “Separating the Variables,” and he showed how to reduce a homogeneous differential equation of the first order to one in which the variables were separable. He applied these methods to problems on orthogonal trajectories. He and his brother Jacob (after whom the “Bernoulli Equation” is named; see Section 1.6.1) succeeded in reducing a large number of differential equations to forms they could solve. Integrating Factors were probably discovered by Euler (1734) and (independently of him) by Fontaine and Clairaut, though some attribute them to Leibniz. Singular Solutions, noticed by Leibniz (1694) and Brook Taylor (1715), are generally associated with the name of Clairaut (1734). The geometrical interpretation was given by Lagrange in 1774, but the theory in its present form was not given until much later by Cayley (1872) and M.J.M. Hill (1888).

Today, differential equations are used in many different fields. They can often accurately capture the behavior of continuous models or a large number of discrete objects where the current state of the system determines the future behavior of the system. Such models are called **deterministic** (as opposed to **stochastic** or **random**). The study of **nonlinear** differential equations is still a very active area of research. Although this text will consider some nonlinear differential equations, here the focus will be on the linear case. We will begin with some basic terminology.

1.1 Introduction to First-Order Equations

Order, Linear, Nonlinear

We begin our study of differential equations by explaining what a differential equation is. From our experience in calculus, we are familiar with some differential equations. For example, suppose that the acceleration of a falling object is $a(t) = -32$, measured in ft/sec². Using the fact that the derivative of the velocity function $v(t)$ (measured in ft/sec) is the acceleration function $a(t)$, we can solve the equation

$$v'(t) = a(t) \quad \text{or} \quad \frac{dv}{dt} = a(t) = -32.$$

Many different types of differential equations can arise in the study of familiar phenomena in subjects ranging from physics to biology to economics to chemistry. We give examples from various fields throughout the text and engage the reader with many such applications.

It is clearly necessary (and expedient) to study, independently, more restricted classes of these equations. The most obvious classification is based on the nature of the derivative(s) in the equation. A differential equation involving derivatives of a function of one variable (ordinary derivatives) is called an **ordinary differential equation**, whereas one containing partial derivatives of a function of more than one independent variable is called a **partial differential equation**. In this text, we will focus on ordinary differential equations.

The **order** of a differential equation is defined as the order of the highest derivative appearing in the equation.

Example 1 The following are examples of differential equations with indicated orders:

- (a) $dy/dx = ay$ (first-order)
- (b) $x''(t) - 3x'(t) + x(t) = \cos t$ (second-order)
- (c) $(y^{(4)})^{3/5} - 2y'' = \cos x$ (fourth-order)

where the superscript (4) in (c) represents the fourth derivative.

Our focus will be on **linear** differential equations, which are those equations that have an unknown function, say y , and each of its higher derivatives appearing in linear functions. That is, we do *not* see them as y^2 , yy' , $\sin y$, or $(y^{(4)})^{3/5}$.¹ More precisely, a linear differential equation is one in which the dependent variable and its derivatives appear in additive combinations of their first powers. Equations where one or more of y and its derivatives appear in nonlinear functions are called **nonlinear** differential equations. In the above example, only (c) is a nonlinear differential equation.

Example 2 Classify the equations as linear or nonlinear.

- (a) $y'' + 3y' - x^2y = \cos x$
- (b) $y'' - 3y' + y^2 = 0$
- (c) $y^{(3)} + yy' + \sin y = x^2$

Solution

The first of these equations is linear, as it consists of an additive combination of y , y' , and y'' , each of which is raised to the first power. In contrast to this, the second equation is nonlinear because of the y^2 term. The last equation is nonlinear because of both the yy' term and the $\sin y$ term—either of these terms by itself would have made the equation nonlinear. Our study of nonlinear differential equations will focus on techniques for specific equations or on understanding the qualitative behavior of a nonlinear differential equation, since general techniques of solution are rarely applicable.

Much of this book is concerned with the solutions of linear differential equations. Thus we need to explain what we mean by a solution. First we note that any n th-order differential equation can be written in the form

$$F(x, y, y', \dots, y^{(n)}) = 0, \quad (1.1)$$

where n is a positive integer. For example, $y' = x^2 + y^2$ can be written as

$$y' - x^2 - y^2 = 0.$$

¹Most of the equations we consider will involve an unknown function y that depends on x . Two other common variables used are (i) the unknown function y that depends on t and (ii) the unknown function x that depends on t , the latter being used in Example 1b.

Here $F(x, y, y') = y' - x^2 - y^2$. The second-order equation $y'' - 3x^2y' + 5y = \sin x$ can be written as

$$y'' - 3x^2y' + 5y - \sin x = 0$$

and we see that $F(x, y, y', y'') = y'' - 3x^2y' + 5y - \sin x$.

Definition 1.1.1

A *solution* to an n th-order differential equation is a function that is n times differentiable and that satisfies the differential equation. Symbolically, this means that a solution of differential equation (1.1) is a function $y(x)$ whose derivatives $y'(x), y''(x), \dots, y^{(n)}(x)$ exist and that satisfies the equation

$$F(x, y(x), y'(x), \dots, y^{(n)}(x)) = 0$$

for all values of the independent variable x in some interval (a, b) where

$$F(x, y(x), y'(x), \dots, y^{(n)}(x))$$

is defined. (Note that the solution to a differential equation does not contain any derivatives, although the derivatives of this solution exist.) The interval (a, b) may be infinite; that is, $a = -\infty$, or $b = \infty$, or both.

Example 3

The function $y(x) = 2e^{3x}$ is a solution of the differential equation

$$\frac{dy}{dx} = 3y,$$

for $x \in (-\infty, \infty)$ because it satisfies the differential equation by giving an identity:

$$\frac{dy}{dx} = 2 \frac{de^{3x}}{dx} = 6e^{3x} = 3y.$$

Initial-Value vs. Boundary-Value Problems

We will soon see that solving a general differential equation gives rise to a solution that has constants. These constants can be eliminated by specifying the initial state of the system or conditions that the solution must satisfy on its domain of definition or “boundary.” An example of the first situation is specifying the position and velocity of a mass on a spring. An example of the second is a rope hanging from two supports, given the location of these two supports.

Consider a first-order differential equation

$$\frac{dy}{dx} = f(x, y)$$

and suppose that the solution $y(x)$ was subject to the condition that $y(x_0) = y_0$. This is an example of an **initial-value problem**. The condition $y(x_0) = y_0$ is called an **initial condition** and x_0 is called the **initial point**. More generally, we have the following:

Definition 1.1.2

An *initial-value problem* consists of an n th-order differential equation together with n initial conditions of the form

$$y(x_0) = a_0, \quad y'(x_0) = a_1, \dots, \quad y^{(n-1)}(x_0) = a_{n-1}$$

that must be satisfied by the solution of the differential equation and its derivatives at the initial point x_0 .

Example 4

The following are examples of initial-value problems:

(a) $dy/dx = 2y - 3x$, $y(0) = 2$ (here $x = 0$ is the initial point).

(b) $x''(t) + 5x'(t) + \sin(tx(t)) = 0$, $x(1) = 0$, $x'(1) = 7$ (here $t = 1$ is the initial point).

(Note that the differential equation in (a) is linear, whereas the equation in (b) is non-linear.) We define a *solution* to an n th-order initial-value problem as a function that is n times differentiable on an interval (a, b) ; this satisfies the given differential equation on that interval, and satisfies the n given initial conditions with the requirement that $x_0 \in (a, b)$. As before, the interval (a, b) might be infinite.

In contrast to an initial-value problem, a **boundary-value problem** consists of a differential equation and a set of conditions *at different x -values* that the solution $y(x)$ must satisfy. Although any number of conditions (≥ 2) may be specified, usually only two are given. Rather than specifying the initial state of the system, we can think of a boundary-value problem as specifying the state of the system at two different physical locations, say $x_0 = a$, $x_1 = b$, $a \neq b$.

Example 5

The following are examples of boundary-value problems:

(a) $d^2y/dx^2 + 5xy = \cos x$, $y(0) = 0$, $y'(\pi) = 2$

(b) $dy/dx + 5xy = 0$, $y(0) = y(1) = 2$

Although a boundary-value problem may not seem too different from an initial-value problem, methods of solution are quite varied. We will focus on initial-value problems. We ask whether an initial-value problem has a unique solution. Essentially this is two questions:

1. Is there a solution to the problem?
2. If there is a solution, is it the only one?

As we see in the next two examples, the answer may be “no” to each question.

Example 6

An *initial-value problem with no solution*.

The initial-value problem

$$\left(\frac{dy}{dx}\right)^2 + y^2 + 1 = 0$$

with $y(0) = 1$ has no real-valued solutions, since the left-hand side is always positive for real-valued functions.

Example 7

An *initial-value problem with more than one solution*.

The initial-value problem

$$\frac{dy}{dx} = xy^{1/3}$$

with $y(0) = 0$ has at least two solutions in the interval $-\infty < x < \infty$. Note that the functions

$$y = 0 \text{ and } y = \frac{x^3}{3\sqrt{3}}$$

both satisfy the initial condition and the differential equation. ■

Two Important Models

One of the most fundamental models in biology deals with population growth and one of the most fundamental models in physics deals with a mass on a spring. In the next two examples, we examine how differential equations describe the behavior of these two phenomena.

Example 8 The change in the population of bacteria with sufficient nutrients and space to grow is known to be proportional to its current population. The differential equation can be written as

$$\frac{dP}{dt} = kP \quad (1.2)$$

where $P(t)$ is the current population of bacteria and k is a constant determined by its growth rate. We can verify that

$$P(t) = P(0)e^{kt} \quad (1.3)$$

is a solution to this differential equation. Because of the presence of the constant $P(0)$, we say that Equation (1.3) is a family of solutions parameterized by the constant $P(0)$. To verify this is indeed a solution, we take the derivative to get $P(0)ke^{kt}$. Substituting this into the left side of the differential equation and the supposed solution into the right side,

$$\underbrace{\frac{dP}{dt}}_{P(0)ke^{kt}} = k \cdot \underbrace{P}_{P(0)e^{kt}}$$

we see that, with a slight rearrangement of the expressions underneath, we have equality for all t . Thus (1.3) is a solution to differential equation (1.2) for all t , and we see the solution describes the *exponential growth* of the population. ■

Example 9 In a later chapter we will learn that a mass on a spring moving along a slippery² surface can be described by the differential equation

$$mx'' + kx = 0$$

where $x(t)$ is the distance the spring has stretched from its resting length, k is the spring constant, and m is the mass, as shown in Figure 1.1. We can verify that $x = \cos\left(\sqrt{\frac{k}{m}}t\right)$ is a solution. To do so we take the second derivative to get $x'' = -\frac{k}{m}\cos\left(\sqrt{\frac{k}{m}}t\right)$ and substitute it into the equation along with the assumed form of x :

$$m \cdot \underbrace{\left[-\frac{k}{m}\cos\left(\sqrt{\frac{k}{m}}t\right)\right]}_{x''} + k \cdot \underbrace{\cos\left(\sqrt{\frac{k}{m}}t\right)}_x = 0.$$

²Physicists use the word “slippery” to mean “ignore frictional forces.”

Simplification shows that it is indeed a solution and it holds for all t .




FIGURE 1.1: Model of spring system for Example 9. $x = 0$ marks the position if the spring were at its natural (unstretched) length and we will take x to the right as positive.

In the next several sections we will develop methods for finding solutions to first-order differential equations. We will then discuss existence and uniqueness of solutions in Chapter 2.

Problems

In Problems 1–12, classify the differential equations by specifying (i) the order, (ii) whether it is linear or nonlinear, and (iii) whether it is an initial-value or boundary-value problem (where appropriate).

1. $3y'' + y = \sin x$
2. $y'' = \sin x$
3. $y'' + y' - y = 0$
4. $y'' + 3y' = 0, \quad y(0) = 1, y'(1) = 0$
5. $y^{(3)} + (\sin x)y^{(2)} + y = x, \quad y(0) = 1, y'(0) = 0, y''(0) = 2$
6. $y'' = 0, \quad y(1) = 1, y'(1) = 2$
7. $y' + e^x y = y^4, \quad y(0) = 0$
8. $y'' - 3yy' = x$
9. $y'' + \sin y = 0$
10. $y'' - 4y' + 4y = 0, \quad y(0) = 1, y'(0) = 1$
11. $y'' + e^x y' + y^2 = 0, \quad y(0) = 1, y(\pi) = 0$
12. $x^2 y'' + y' + (\ln x)y = 0$

In Problems 13–24, verify that the given function is a solution to the differential equation by substituting it into the differential equation and showing that the equation holds true. Assume the interval is $(-\infty, \infty)$ unless otherwise stated. Do NOT attempt to solve the differential equation.

13. $y(x) = 2x^3, \quad x \frac{dy}{dx} = 3y$
14. $y(x) = x, \quad y'' + y = x$
15. $y = 2, \quad \frac{dy}{dx} = x^3(y - 2)^2$
16. $y(x) = x^3, \quad \frac{dy}{dx} = 3y^{2/3}$
17. $y(x) = e^x - x, \quad \frac{dy}{dx} + y^2 = e^{2x} + (1 - 2x)e^x + x^2 - 1$
18. $y(x) = \sin x + 2 \cos x, \quad y'' + y = 0$
19. $y(x) = x^2 - x^{-1}, \quad x^2 y'' = 2y, \quad x \neq 0$
20. $y(x) = x + C \sin x, \quad y'' + y = x, \quad C = \text{constant}$
21. $y(x) = \frac{-1}{x - 3}, \quad \frac{dy}{dx} = y^2, \quad (-\infty, 3)$
22. $y(x) = \frac{-1}{5x + 4}, \quad \frac{dy}{dx} = 5y^2, \quad (-4/5, \infty)$
23. $y_1(x) = e^x$ and $y_2(x) = e^{2x}, \quad y'' - 3y' + 2y = 0$
24. $y_1(x) = e^x$ and $y_2(x) = xe^x, \quad y'' - 2y' + y = 0$

In Problems **25–28**, determine which of the functions solve the given differential equation.

25. $y'' + 6y' + 9y = 0$: (a) e^x , (b) e^{-3x} , (c) xe^{-3x} , (d) $4e^{3x}$, (e) $2e^{-3x} + xe^{-3x}$

26. $y'' + 9y = 0$: (a) $\sin 3x$, (b) $\sin x$, (c) $\cos 3x$, (d) e^{3x} , (e) x^3

27. $y'' - 7y' + 12y = 0$: (a) e^{2x} , (b) e^{3x} , (c) e^{4x} , (d) e^{5x} , (e) $e^{3x} + 2e^{4x}$

28. $y'' + 4y' + 5y = 0$: (a) e^{-2x} , (b) $e^{-2x} \sin 2x$, (c) $e^{-2x} \cos 2x$, (d) $\cos 2x$

In Problems **29–32**, find values of r for which the given function solves the differential equation on $(-\infty, \infty)$.

29. $y(x) = e^{rx}$, $y'' + 3y' + 2y = 0$

30. $y(x) = e^{rx}$, $y'' + 3y' - 4y = 0$

31. $y(x) = xe^{rx}$, $y'' + 6y' + 9y = 0$

32. $y(x) = xe^{rx}$, $y'' + 4y' + 4y = 0$

1.2 Separable Differential Equations

We will now introduce the simplest first-order differential equations. Although these are the simplest class of differential equations we will encounter, they appear in numerous applications and aspects of subsequent theory. We make the following definition:

Definition 1.2.1

A first-order differential equation that can be written in the form

$$g(y)y' = f(x) \quad \text{or} \quad g(y)dy = f(x)dx,$$

where $y = y(x)$, is called a separable differential equation.

Separable differential equations are solved by collecting all the terms involving the dependent variable y on one side of the equation and all the terms involving the independent variable x on the other side. Once this is completed (it may require some algebra), both sides of the resulting equations are integrated. That is, the equation

$$g(y)y' = f(x)$$

can be written in “differential form”

$$g(y) \frac{dy}{dx} = f(x)$$

so that treating dy/dx as a fraction, we have

$$g(y)dy = f(x)dx.$$

Here the variables are separated, so that integrating both sides gives

$$\int g(y)dy = \int f(x)dx. \tag{1.4}$$

The Method of Separation of Variables, which we just applied to (1.4), is the name given to the method we use to solve Separable Equations—it is one of the simplest and most useful methods for solving differential equations. (Incidentally, it is an important technique for solving certain classes of partial differential equations, too.)

Sometimes we will be able to solve (1.4) for y . When we can do so, we will say we can express the **explicit solution** and will write $y = h(x)$. Other times, we will not be able

to solve (1.4) or it will not be worth our time and efforts to do so. In these situations, we say that we are giving the **implicit solution** with (1.4). When our solution can be written explicitly, it will be easy to plot solutions in the x - y plane, by hand or with the computer; however, when the solution is implicit, plotting solutions by hand is challenging at best. The various computer programs discussed in Appendix A and the end of each chapter will allow us to view plots in the x - y plane without much additional work. We now consider a number of examples.

Example 1 Solve $y' = ky$ where k is a constant.

Solution

Writing y' as $\frac{dy}{dx}$ gives

$$\frac{dy}{dx} = ky.$$

Treating $\frac{dy}{dx}$ as a “fraction” and rearranging terms gives

$$\frac{dy}{y} = k dx.$$

This step will only be valid if $y \neq 0$. We note that $y = 0$ is also a solution to the original differential equation. Integrating gives

$$\int \frac{dy}{y} = \int k dx,$$

which is

$$\ln |y| = kx + C_1, \implies |y| = e^{kx+C_1}.$$

This gives

$$y = \pm e^{kx} e^{C_1}.$$

Now e^{C_1} is a positive constant, so that we may let $C = \pm e^{C_1}$. In the above process, we encountered the constant solution $y = 0$, which also gives us the possibility that $C = 0$. Thus, we have

$$y = Ce^{kx} \tag{1.5}$$

as our solution, where $x \in (-\infty, \infty)$ and C is any real constant. We say that (1.5) defines a **one-parameter family of solutions** of $y' = ky$. It is also important to remember the “trick” used above for getting rid of the absolute values—it will come up quite often in practice! We will consider a few more examples with similar standard “tricks.”

Example 2 Solve $\frac{dx}{dt} = e^{t-x}$, $x(0) = \ln 2$, for $x(t)$.

Solution

Separating the variables gives

$$\frac{dx}{dt} = e^t e^{-x}$$

and thus

$$e^x dx = e^t dt.$$

Integrating both sides of this equation gives

$$e^x = e^t + C.$$

Solving for x , we have

$$x = \ln |e^t + C|.$$

Applying the initial condition $x(0) = \ln 2$ yields

$$\ln 2 = \ln |1 + C|, \quad \text{so that} \quad C = 1.$$

Thus

$$x = \ln(e^t + 1),$$

which is defined for all t . Note that $e^t + 1$ is always positive so that we can drop the absolute value signs. *We should also note that, after integrating, we could have applied the initial condition to determine C and then proceeded to solve for x instead of first solving for x and then applying the initial condition to determine C . Both methods will result in the same final answer.* See Figure 1.2 for a plot of the solution.




FIGURE 1.2: Plot of solution for Example 2.

Example 3 Solve $(x - 4)y^4 - x^3(y^2 - 3) \frac{dy}{dx} = 0$.

Solution

To separate variables, we divide by x^3y^4 , which implicitly assumes that $x \neq 0$ and $y \neq 0$. Doing so gives

$$\frac{x - 4}{x^3} dx = \frac{y^2 - 3}{y^4} dy.$$

This simplifies to $(x^{-2} - 4x^{-3}) dx = (y^{-2} - 3y^{-4}) dy$. Integrating gives

$$\frac{-1}{x} + \frac{2}{x^2} = \frac{-1}{y} + \frac{1}{y^3} + C$$

as the general solution, which is valid when $x \neq 0$ and $y \neq 0$. This is definitely a case where giving the solution in an implicit representation is acceptable! See Figure 1.3 for a plot of the implicit solution. We refer the reader to the end of this chapter for the computer code used to plot these types of solutions with one of the software packages. There is, however, a more important idea that is illustrated by this example. If we assume $x \neq 0$ and $y^2 - 3 \neq 0$, we can rewrite the original differential equation as

$$\frac{dy}{dx} = \frac{(x - 4)y^4}{x^3(y^2 - 3)},$$




FIGURE 1.3: Implicit plot for Example 3. The curves plotted here satisfy the implicit solution. We note here that the C -values superimposed on the curves were good for this problem, but it often takes ingenuity, experience, trial and error, or some combination of these to get a “nice” picture.

and then one can clearly see that $y = 0$ is a solution. (That is, when $y = 0$ is substituted into both sides of the equation, we get an identity for all x .) This problem shows that the separation process can lose solutions.

How can we verify that

$$\frac{-1}{x} + \frac{2}{x^2} = \frac{-1}{y} + \frac{1}{y^3} + C$$

is a solution? We need to substitute it into the differential equation as before. This will require us to find y' and we will do so with implicit differentiation. Taking the derivative of both sides of the equation gives

$$\frac{1}{x^2} - \frac{4}{x^3} = \frac{1}{y^2}y' - \frac{3}{y^4}y'.$$

We solve for y' and then simplify the complex fraction to obtain

$$y' = \frac{y^4(x-4)}{x^3(y^2-3)},$$

which is an equivalent form of our original differential equation. ■

Although the separation process will work on any differential equation in the form of Definition 1.2.1, evaluating the integrals in (1.4) can sometimes be a daunting, if not impossible, task. As discussed in calculus, certain indefinite integrals such as

$$\int e^{x^2} dx$$

cannot be expressed in finite terms using elementary functions. When such an integral is encountered while solving a differential equation, it is often helpful to use definite integration by assuming an initial condition $y(x_0) = y_0$.

Example 4 Solve the initial-value problem

$$\frac{dy}{dx} = e^{x^2} y^2, \quad y(2) = 1$$

and use the solution to give an approximate answer for $y(3)$.

Solution

We would like to divide both sides by y^2 and we note that $y = 0$ is a solution. We set this solution aside and now assume $y \neq 0$, divide by y^2 , and integrate from $x = 2$ to $x = x_1$ to obtain

$$\begin{aligned}\int_2^{x_1} [y(x)]^{-2} \frac{dy}{dx} dx &= -[y(x)]^{-1} \Big|_2^{x_1} \\ &= \frac{-1}{y(x_1)} + \frac{1}{y(2)} \\ &= \int_2^{x_1} e^{x^2} dx.\end{aligned}$$

If we let t be the variable of integration and replace x_1 by x and $y(2)$ by 1, then we can express the solution to the initial-value problem by

$$y(x) = \frac{1}{1 - \int_2^x e^{t^2} dt}.$$

With an explicit solution, we often want to be able to find the corresponding y -value given any x . The right-hand side still cannot be solved exactly but can be approximated if x is given. For example, $y(3) \approx -0.0007007$. We note that we will have a point $x > 2$ that will make the denominator zero (and thus is not in the domain of our solution) and our function will become unbounded.

It is sometimes the case that a substitution or other “trick” will convert the given differential equation into a form that we can solve. A differential equation of the form

$$\frac{dy}{dx} = f(ax + by + k),$$

where a , b , and k are constants, is separable if $b = 0$; however, if $b \neq 0$, the substitution

$$u(x) = ax + by + k$$

makes it a separable equation.

Example 5

Solve

$$\frac{dy}{dx} = (x + y - 4)^2$$

by first making an appropriate substitution.

Solution

We let $u = x + y - 4$ and thus $\frac{dy}{dx} = u^2$. We need to calculate $\frac{du}{dx}$. For this example, taking the derivative with respect to x gives

$$\frac{du}{dx} = 1 + \frac{dy}{dx}.$$

Substitution into the original differential equation gives

$$\frac{du}{dx} - 1 = u^2.$$

This equation is separable. Dividing by $1 + u^2$, we obtain

$$\frac{du}{1+u^2} = dx$$

and integrating gives

$$\arctan(u) = x + c.$$

Thus $u = \tan(x + c)$. Since $u = x + y - 4$, we then have

$$y = -x + 4 + \tan(x + c),$$

which is defined wherever $\tan(x + c)$ is defined.

Problems

In Problems **1–20**, solve each of the following differential equations. Explicitly solve for $y(x)$ or $x(t)$ when possible.

1. $\frac{dy}{dx} = \cos x$
2. $x \frac{dy}{dx} = (1 + y)^2$
3. $x \frac{dx}{dt} + t = 1$
4. $(1 + x) \frac{dy}{dx} = 4y$
5. $\tan x \, dy + 2y \, dx = 0$
6. $\frac{dy}{dx} = 2\sqrt{xy}$
7. $4xy \, dx + (x^2 + 1) \, dy = 0$
8. $\frac{dy}{dx} = \frac{x^2}{1+y^2}$
9. $y' = 10^{x+y}$
10. $xy' = \sqrt{1 - y^2}$
11. $y' = xye^{x^2}$, $y(0) = 1$. Explain why this differential equation guarantees that its solution is symmetric about $x = 0$.
12. $y' = 2x^2(y^2 + 1)$, $y(0) = 1$
13. $(e^x + 1) \cos y \, dy + e^x(\sin y + 1) \, dx = 0$, $y(0) = 3$
14. $(\tan x)y' = y$, $y(\frac{\pi}{2}) = \frac{\pi}{2}$
15. $2x(y^2 + 1) \, dx + (x^4 + 1) \, dy = 0$, $y(1) = 1$
16. $(x^2 - 1)y' + 2xy^2 = 0$, $y(\sqrt{2}) = 1$
17. $(y + 2) \, dx + y(x + 4) \, dy = 0$, $y(-3) = -1$
18. $8 \cos^2 y \, dx + \csc^2 x \, dy = 0$, $y(\pi/12) = \pi/4$
19. $y' = e^{x^2}$, $y(0) = 0$
20. $\frac{dy}{dx} = \frac{y^3 + 2y}{x^2 + 3x}$, $y(1) = 1$
21. Find the solution of the following equation that satisfies the given conditions for $x \rightarrow +\infty$: $x^2 y' - \cos 2y = 1$, $y(+\infty) = \frac{9\pi}{4}$.
22. Find the solution of the following equation that satisfies the given conditions for $x \rightarrow +\infty$: $3y^2 y' + 16x = 2xy^3$, $y(x)$ is bounded for $x \rightarrow +\infty$.

In Problems **23–27**, make an appropriate substitution to solve each of the following differential equations. Explicitly solve for $y(x)$ or $x(t)$ when possible.

23. $xydx + (x+1)dy = 0$

24. $y' - y = 2x - 3$

25. $(x+2y)y' = 1 \quad y(0) = -2$

26. $y' = \cos(y-x)$

27. $y' = \sqrt{4x+2y-1}$

28. Suppose that the population $N(t)$ of a given species (bacteria, elves, Toolie birds, college students, etc.) is not always zero and varies at a rate proportional to its current value. That is,

$$\frac{dN}{dt} = rN,$$

where $r \in \mathbb{R}$ is some measured constant proportionality factor. If the initial population is assumed to be $N(0) = N_0 > 0$, solve this exponential differential equation and discuss the behavior of the solution as $t \rightarrow \infty$ for different values of r .

29. An equivalent way of thinking of the exponential growth problem 28 is to assume the per capita growth rate, $\frac{1}{N} \frac{dN}{dt}$, is constant. That is, we assume $\frac{1}{N} \frac{dN}{dt} = r$. It is more realistic to assume that the per capita growth rate decreases as the population grows. If we assume this decrease is linear and agrees with the exponential growth model for small populations, we can write the equation

$$\frac{1}{N} \frac{dN}{dt} = r \left(1 - \frac{N}{K} \right)$$

where the left-hand side is the per capita growth rate and the right-hand side is a linearly decreasing function in N that has y -intercept r and x -intercept K . Multiplying both sides by N gives

$$\frac{dN}{dt} = r \left(1 - \frac{N}{K} \right) N,$$

which is the well-known *logistic* differential equation. If the initial population is given as $N(0) = N_0 > 0$, solve this differential equation and discuss the behavior of the solution as $t \rightarrow \infty$. From this behavior, why is K called a *carrying capacity*?

1.3 Linear Equations

Linear first-order differential equations are perhaps the most commonly arising class of differential equations in applications. A linear differential equation is defined as follows:

Definition 1.3.1

A first-order ordinary differential equation is linear in the dependent variable y and the independent variable x if it can be written as

$$\frac{dy}{dx} + P(x)y = Q(x). \quad (1.6)$$

More generally, we often see equations of the form

$$a_1(x)y' + a_0(x)y = b(x)$$

but, provided $a_1(x) \neq 0$ for all x , we can always divide by $a_1(x)$ and define $P(x) = a_0/a_1$ and $Q(x) = b/a_1$ to obtain an equation of the form of (1.6).

In our work to follow, specifically in Chapters 3 and 4, we will refer to an equation of this form as a “linear nonhomogeneous equation.” In the case when $Q(x) = 0$, we refer to the equation as “homogeneous,” but we caution the reader to be careful with the word “homogeneous” as it can also have other meanings; see Section 1.6. While it is an unfortunate fact that mathematicians often use the same term for different mathematical notions, our use of it should be clear by context.

In the following pages, we present two techniques for solving linear differential equations. It is likely the case that only one of these methods will be presented in class, depending on the emphasis of your course. The first is variation of parameters while the second is the integrating factor technique.

Variation of Parameters

The first method of solving linear equations that we consider has a nice generalization for higher-order equations. If we consider (1.6) with $Q(x) = 0$,

$$\frac{dy}{dx} + P(x)y = 0,$$

we can solve this linear homogeneous equation by using separation of variables. We obtain y_c , the **complementary solution**.³ We know that we can multiply y_c by any constant and it will still be a solution; however, we instead consider uy_c where u is a function of x and try to find a function u that will make this work. In order for $u(x)y_c$ to be a solution, it needs to satisfy the differential equation. Substituting the assumed solution into (1.6), we obtain

$$(u'(x)y_c + u(x)y_c') + P(x)u(x)y_c = Q(x), \quad (1.7)$$

which we can regroup and then simplify as

$$\begin{aligned} u'(x)y_c + u(x)\underbrace{[y_c' + P(x)y_c]}_{=0} &= Q(x) \\ \implies u'(x)y_c &= Q(x) \end{aligned}$$

since y_c is a solution to the homogeneous equation. We then solve for $u'(x)$ and integrate to obtain

$$u(x) = \int \frac{Q(x)}{y_c} dx. \quad (1.8)$$

As we only care about finding one function $u(x)$ that will work, we don't introduce the typical $+C$ upon integration. Thus we have found a function $u(x)$ that makes uy_c a solution—we call this a particular solution and denote it y_p . Our general solution to (1.6), with $y_p = u(x)y_c$, is then

$$y = Cy_c + y_p, \quad (1.9)$$

where C is a constant that is determined by the initial condition.

Example 1 Solve $\frac{dy}{dx} + 2xy = 3x$ using variation of parameters.

Solution

This equation is linear with $P(x) = 2x$ and $Q(x) = 3x$. We solve the homogeneous equation

³This solution is sometimes called the **homogeneous solution** and is denoted y_h . The terms are used interchangeably.

first:

$$\begin{aligned}\frac{dy}{dx} &= -2xy \Rightarrow \int \frac{dy}{y} = \int -2x dx \\ &\Rightarrow \ln |y| = -x^2 + C_1 \\ &\Rightarrow y = Ce^{-x^2}.\end{aligned}\tag{1.10}$$

We now assume that a particular solution can be written as

$$y_p = u(x)y_c = u(x)e^{-x^2}.$$

The function $u(x)$ that will allow this to be a solution of the original linear equation is

$$\begin{aligned}u(x) &= \int \frac{Q(x)}{y_c} dx \\ &= \int \frac{3x}{e^{-x^2}} dx \\ &= 3 \int xe^{x^2} dx \\ &= \frac{3}{2}e^{x^2}.\end{aligned}\tag{1.11}$$

Recalling that $y_p = u(x)y_c$, our solution is then given by

$$\begin{aligned}y &= Cy_c + y_p \\ &= \underbrace{Ce^{-x^2}}_{y_c} + \underbrace{\frac{3}{2}e^{x^2}}_{u(x)} \underbrace{e^{-x^2}}_{y_c}\end{aligned}\tag{1.12}$$

which simplifies to

$$y = \frac{3}{2} + Ce^{-x^2}.$$

We can easily check that this is a solution of the original differential equation. ■

Example 2 Solve

$$\frac{dy}{dx} + \left(\frac{2x+1}{x} \right) y = e^{-2x}$$

using variation of parameters.

Solution

This is clearly linear and we first solve the homogeneous equation

$$\frac{dy}{dx} + \left(\frac{2x+1}{x} \right) y = 0.$$

Separation of variables gives us

$$y_c = C \frac{e^{-2x}}{x}.$$

We now assume a particular solution of the form $y_p = u(x) \frac{e^{-2x}}{x}$. From the derivation, we know that things will cancel out so that we need to solve for u in (1.8):

$$u(x) = \int \frac{e^{-2x}}{e^{-2x}/x} = \int x dx$$

so that

$$u(x) = \frac{x^2}{2} \implies y_p = u(x)y_c = \frac{x^2}{2} \frac{e^{-2x}}{x} = \frac{1}{2}xe^{-2x}.$$

Our general solution is $y_c + y_p$:

$$y = \frac{C}{x}e^{-2x} + \frac{1}{2}xe^{-2x}.$$

Superposition

A key idea in the study of linear differential equations is that of **superposition**. We have been studying the basic linear equation (1.6)

$$\frac{dy}{dx} + P(x)y = Q(x).$$

We can state a very useful theorem that will serve as an important tool in our further study.

THEOREM 1.3.1 Superposition

Suppose that y_1 is a solution to $y' + P(x)y = Q_1(x)$ and y_2 is a solution to $y' + P(x)y = Q_2(x)$. Then

$$c_1y_1 \quad \text{is a solution to} \quad y' + P(x)y = c_1Q_1(x)$$

for any constant c_1 . For any constants c_1, c_2 , we also have that

$$c_1y_1 + c_2y_2 \quad \text{is a solution to} \quad y' + P(x)y = c_1Q_1(x) + c_2Q_2(x).$$

Example 3 Verify that e^{2x} is a solution to $\frac{dy}{dx} + y = 3e^{2x}$ and $5x - 5$ is a solution to

$\frac{dy}{dx} + y = 5x$. Then find a solution to

$$\frac{dy}{dx} + y = e^{2x} + 4x.$$

Solution

We can easily verify that $y_1 = e^{2x}$ and $y_2 = 5x - 5$ are the solutions of the respective differential equations. Let $Q_1(x) = 3e^{2x}$, $Q_2(x) = 5x$, and $Q(x) = e^{2x} + 4x$ denote the right-hand sides of the three differential equations. We observe that

$$Q(x) = \frac{1}{3}Q_1(x) + \frac{4}{5}Q_2(x).$$

By superposition, it follows that

$$y = \frac{1}{3}y_1 + \frac{4}{5}y_2 = \frac{1}{3}(e^{2x}) + \frac{4}{5}(5x - 5)$$

is a solution of $y' + y = Q(x)$.

Integrating Factor Technique

In studying separable equations, we put all the terms of one variable on the left side of the equation and the terms of the other variable on the right side of the equation. This allowed

us to integrate functions of just one variable. Another trick that we will use is to rewrite the left side so that it looks like the result of the product rule (from calculus). To remind ourselves, for y, μ that are both functions of the same variable, the product rule states that

$$(y\mu)' = y'\mu + \mu'y.$$

We know how to integrate the left-hand side so the goal is to somehow rewrite part of our equation so that it looks like the right-hand side. Looking at $y'\mu + \mu'y$ and recalling our basic linear equation $y' + Py = Q$, we want to multiply the left-hand side by a function μ that satisfies

$$\mu' = \mu P. \quad (1.13)$$

In this equation, $P = P(x)$ is known, whereas $\mu = \mu(x)$ (called the **integrating factor**) is unknown. We can find $\mu(x)$ because Equation (1.13) is separable. Thus

$$\frac{d\mu}{\mu} = P(x) dx.$$

Integrating gives

$$\mu(x) = e^{\int P(x) dx}. \quad (1.14)$$

Since (1.14) is an integrating factor, we have

$$\underbrace{e^{\int P(x) dx}}_{\mu} \underbrace{\frac{dy}{dx}}_{y'} + \underbrace{e^{\int P(x) dx} P(x) y}_{\mu'} = Q(x) e^{\int P(x) dx},$$

which is the same as

$$\frac{d}{dx} \left(\underbrace{e^{\int P(x) dx}}_{\mu} y \right) = Q(x) e^{\int P(x) dx}.$$

So

$$e^{\int P(x) dx} y = \int Q(x) e^{\int P(x) dx} dx + C,$$

which gives

$$y = e^{-\int P(x) dx} \left(\int Q(x) e^{\int P(x) dx} dx + C \right) \quad (1.15)$$

as the solution of the differential equation (1.6). Note that we have explicitly written the constant of integration even though the integral has not yet been evaluated. Depending upon your situation, one can memorize the formula (1.15) for the solution of a first-order linear equation; however, it is just as easy (if not out right preferable) to simply apply the method of solution each time.

Summary: Solving linear equations via an integrating factor

1. Write the linear equation in the form of Equation (1.6).
2. Calculate the integrating factor $e^{\int P(x) dx}$.
3. Evaluate the integral $\int Q(x) e^{\int P(x) dx} dx$ and then multiply this result by $e^{-\int P(x) dx}$.
4. The general solution to (1.6) is

$$y = C e^{-\int P(x) dx} + e^{-\int P(x) dx} \int Q(x) e^{\int P(x) dx} dx.$$

In the event that we are given an initial condition $y(x_0) = y_0$, we would apply it at the time of integration, going from x_0 to a final (general) value x . If we let $\bar{p}(x) = \int P(x)dx$, then the general formula becomes

$$y = Ce^{-\bar{p}(x)} + e^{-\bar{p}(x)} \int Q(x)e^{\bar{p}(x)} dx,$$

and applying the initial condition gives us the solution

$$y = y_0 e^{\bar{p}(x_0) - \bar{p}(x)} + e^{-\bar{p}(x)} \int_{x_0}^x Q(t)e^{\bar{p}(t)} dt, \quad (1.16)$$

where the variable of integration has changed to a dummy variable t .

Example 4 Solve

$$\frac{dy}{dx} + \left(\frac{2x+1}{x} \right) y = e^{-2x}.$$

This is linear with

$$P(x) = \frac{2x+1}{x} \quad \text{and} \quad Q(x) = e^{-2x}$$

so that an integrating factor is

$$\begin{aligned} e^{\int P(x)dx} &= e^{\int \frac{2x+1}{x} dx} \\ &= e^{(2x + \ln|x|)} \\ &= |x|e^{2x}. \end{aligned}$$

We note that integrating factors are not unique. For instance, dropping the absolute value to obtain xe^{2x} gives another integrating factor of the differential equation. Thus, multiplying the original equation by this expression gives

$$xe^{2x} \frac{dy}{dx} + e^{2x}(2x+1)y = x.$$

If we had multiplied by $-xe^{2x}$, we would have obtained the *same* equation. This equation can be simplified to give

$$\frac{d}{dx}(xe^{2x}y) = x.$$

Integrating this equation gives

$$xe^{2x}y = \frac{1}{2}x^2 + C,$$

which becomes

$$y = \frac{1}{2}xe^{-2x} + \frac{C}{x}e^{-2x}.$$

These last few steps could have been avoided by using (1.15). ■

Example 5 Solve $(x^2 + 1)\frac{dy}{dx} + 4xy = x$ with the initial condition $y(0) = 10$.

Solution

Rewriting this equation gives

$$\frac{dy}{dx} + \left(\frac{4x}{x^2 + 1} \right) y = \frac{x}{x^2 + 1},$$

hence

$$P(x) = \frac{4x}{x^2 + 1} \quad \text{and} \quad Q(x) = \frac{x}{x^2 + 1}$$

so that an integrating factor is

$$\begin{aligned} e^{\int P(x)dx} &= e^{\int \frac{4x}{x^2+1} dx} \\ &= e^{\ln(x^2+1)^2} \\ &= (x^2 + 1)^2. \end{aligned}$$

Once we have our integrating factor, we can use the solution as given in (1.15), first noting that

$$\begin{aligned} e^{-\int P(x)dx} &= e^{-\ln(x^2+1)^2} \\ &= (x^2 + 1)^{-2}. \end{aligned} \tag{1.17}$$

Then

$$\begin{aligned} y &= \frac{1}{(x^2 + 1)^2} \left(\int x(x^2 + 1) dx \right) \\ &= \frac{1}{(x^2 + 1)^2} \left(\frac{1}{4}x^4 + \frac{1}{2}x^2 + C \right). \end{aligned}$$

Now the initial condition, $y(0) = 10$, gives $C = 10$ and thus

$$y = \frac{\frac{1}{4}x^4 + \frac{1}{2}x^2 + 10}{(x^2 + 1)^2}$$

is the solution we seek. ■

Now that we know the techniques of solving linear equations, we consider some applications. In Section 1.4, we will consider **Newton's law of cooling**, which describes how the temperature of an object changes due to the constant temperature of the medium surrounding it. This is not always realistic, as in some settings the temperature of the surroundings varies. For example, determining the temperature inside a building over a span of a 24-hour day is complicated because the outside temperature varies. If we assume that the building has no heating or air conditioning, the differential equation that needs to be solved to find the temperature $u(t)$ at time t inside the building is

$$\frac{du}{dt} = k(C(t) - u(t)), \tag{1.18}$$

where $C(t)$ is a function that describes the outside temperature and $k > 0$ is a constant that depends on the insulation of the building. Note that (1.18) is a linear equation. According to this equation, if $C(t) > u(t)$, then

$$\frac{du}{dt} > 0,$$

which implies that $u(t)$ increases, and if $C(t) < u(t)$, then

$$\frac{du}{dt} < 0,$$

so that $u(t)$ decreases.

Example 6 Suppose that on a given day during the month of April in Pomona, California, the outside temperature in degrees Fahrenheit is given by

$$C(t) = 70 - 10 \cos\left(\frac{\pi t}{12}\right)$$

for $0 \leq t \leq 24$. Determine the temperature in a building that has an initial temperature of 60°F if $k = 1/4$. See Figure 1.4.




FIGURE 1.4: Outside temperature over 24 hours for Example 6.

Solution

We see that the average temperature (i.e., the average of $C(t)$) is 70°F because

$$\int_0^{24} \cos\left(\frac{\pi t}{12}\right) dt = 0.$$

The initial-value problem that we must solve is

$$\frac{du}{dt} = k \left(70 - 10 \cos\left(\frac{\pi t}{12}\right) - u \right)$$

with initial condition $u(0) = 60$. The differential equation can be rewritten as

$$\frac{du}{dt} + ku = k \left(70 - 10 \cos\left(\frac{\pi t}{12}\right) \right),$$

which is a linear equation and is thus solvable. This gives (check it!)

$$u(t) = \frac{10}{9 + \pi^2} \left(63 + 7\pi^2 - 9 \cos\left(\frac{\pi t}{12}\right) - 3\pi \sin\left(\frac{\pi t}{12}\right) \right) + C_1 e^{-t/4}.$$

We then apply the initial condition $u(0) = 60$ to determine the arbitrary constant C_1 and obtain the solution

$$u(t) = \frac{10}{9 + \pi^2} \left(63 + 7\pi^2 - 9 \cos\left(\frac{\pi t}{12}\right) - 3\pi \sin\left(\frac{\pi t}{12}\right) \right) - \frac{10\pi^2}{9 + \pi^2} e^{-t/4}.$$

A graph of this solution is shown in Figure 1.5. The graph shows that the temperature reaches its maximum of about 77°F near $t = 15.5$, which is about 3:30 p.m.




FIGURE 1.5: Inside temperature over 24 hours for Example 6.

Sometimes an equation may not immediately appear to be linear.

Example 7 Consider the differential equation

$$y^2 dx + (3xy - 1) dy = 0.$$

This equation is not linear in y . What do we do? Look harder. If we consider y as the independent variable and x as the dependent variable, we can write

$$\frac{dx}{dy} = \frac{1 - 3xy}{y^2},$$

which is

$$\frac{dx}{dy} + \frac{3x}{y} = \frac{1}{y^2},$$

and we see that it is in the form

$$\frac{dx}{dy} + P(y)x = Q(y),$$

which is linear in x , so that this equation can be solved using the theory we have just developed.

Hence, an integrating factor is

$$e^{\int P(y)dy} = e^{\int \frac{3}{y} dy} = e^{\ln |y|^3} = y^3.$$

We also have $\exp\left(-\int P(y)dy\right) = 1/y^3$. Then our solution is

$$\begin{aligned} x &= \frac{1}{y^3} \left(\int \frac{1}{y^2} (y^3) dy \right) + \frac{C}{y^3} \\ &= \frac{1}{y^3} \left(\frac{y^2}{2} \right) + \frac{C}{y^3}. \end{aligned}$$

This becomes

$$x = \frac{1}{2y} + \frac{C}{y^3}$$

which is defined for all $y \neq 0$.

Problems

Solve the linear equations in Problems 1–18 by considering y as a function of x , that is, $y = y(x)$.

- | | |
|--|---|
| 1. $y' + y = e^x$ | 2. $y' + 2y = 4$ |
| 3. $y' + 2y = -3x$ | 4. $y' - 2xy = e^{x^2}$ |
| 5. $y' - 3x^2y = x^2$ | 6. $3xy' + y = 12x$ |
| 7. $\frac{dy}{dx} + \frac{1}{x}y = x$ | 8. $y' + \frac{1}{x}y = e^x$ |
| 9. $\frac{dy}{dx} - \frac{2x}{1+x^2}y = x^2$ | 10. $xy' + (1+x)y = e^{-x} \sin 2x$ |
| 11. $\frac{dy}{dx} + y = \cos x$ | 12. $(2x+1)y' = 4x+2y$ |
| 13. $\frac{dy}{dx} - y = 4e^x, \quad y(0) = 4$ | 14. $y' + 2y = xe^{-2x}, \quad y(1) = 0$ |
| 15. $y' + y \tan x = \sec x, \quad y(\pi) = 1$ | 16. $y' = (1-y) \cos x, \quad y(\pi) = 2$ |
| 17. $\frac{dy}{dx} + \frac{y}{x} = \frac{\cos x}{x}, \quad y\left(\frac{\pi}{2}\right) = \frac{4}{\pi}, \quad x > 0$ | 18. $xy' + 2y = \sin x, \quad y\left(\frac{\pi}{2}\right) = 1, \quad x > 0$ |

Solve the linear equations in Problems 19–21 by considering x as a function of y , that is, $x = x(y)$.

- | | |
|----------------------------------|------------------------|
| 19. $(x + y^2)dy = ydx$ | 20. $(2e^y - x)y' = 1$ |
| 21. $(\sin 2y + x \cot y)y' = 1$ | |

Problems 22–23 address aspects of superposition.

22. Recall that a linear equation is called **homogeneous** if $Q(x) = 0$, i.e., if it can be written as

$$\frac{dy}{dx} + P(x)y = 0.$$

- (a) Show that $y = 0$ is a solution (called the *trivial* solution).
 (b) Show that if $y = y_1(x)$ is a solution and k is a constant, then $y = ky_1(x)$ is also a solution.
 (c) Show that if $y = y_1(x)$ and $y = y_2(x)$ are solutions, then $y = y_1(x) + y_2(x)$ is a solution.
23. (a) If $y = y_1(x)$ satisfies the homogeneous linear equation $\frac{dy}{dx} + P(x)y = 0$ and $y = y_2(x)$ satisfies the nonhomogeneous linear equation $\frac{dy}{dx} + P(x)y = r(x)$, show that

$y = y_1(x) + y_2(x)$ is a solution to the nonhomogeneous linear equation

$$\frac{dy}{dx} + P(x)y = r(x).$$

(b) Show that, if $y = y_1(x)$ is a solution of $\frac{dy}{dx} + P(x)y = r(x)$, and $y = y_2(x)$ is a solution of $\frac{dy}{dx} + P(x)y = q(x)$, then $y = y_1(x) + y_2(x)$ is a solution of

$$\frac{dy}{dx} + P(x)y = q(x) + r(x).$$

(c) Use the results obtained in parts (a) and (b) to solve

$$\frac{dy}{dx} + 2y = e^{-x} + \cos x.$$

- 24.** A pond that initially contains 500,000 gal of unpolluted water has an outlet that releases 10,000 gal of water per day. A stream flows into the pond at 12,000 gal/day containing water with a concentration of 2 g/gal of a pollutant. Find a differential equation that models this process, and determine what the concentration of pollutant will be after 10 days.
- 25.** When wading in a river or stream, you may notice that microorganisms like algae are frequently found on rocks. Similarly, if you have a swimming pool, you may notice that, in the absence of maintaining appropriate levels of chlorine and algaecides, small patches of algae take over the pool surface, sometimes overnight. Underwater surfaces are attractive environments for microorganisms because water removes waste and provides a continuous supply of nutrients. On the other hand, the organisms must spread over the surface without being washed away. If conditions become unfavorable, they must be able to free themselves from the surface and recolonize on a new surface.

The rate at which cells accumulate on a surface is proportional to the rate of growth of the cells and the rate at which the cells attach to the surface. An equation describing this situation is given by

$$\frac{dN(t)}{dt} = r(N(t) + A),$$

where $N(t)$ represents the cell density, r the growth rate, A the attachment rate, and t time.

(a) If the attachment rate, A , is constant, solve

$$\frac{dN(t)}{dt} = r(N(t) + A)$$

with the initial condition $N(0) = 0$.

(b) If $A = 3$ in a particular colony of cells, use the following table to find the growth rate at the end of each hour:

t	1	2	3	4
$N(t)$	3	9	21	45

Using this growth rate, estimate the algae population size at the end of 24 hours and 36 hours.

- 26.** In Section 3.7, you will learn about electric circuits as an application of a second-order differential equation. However, consider the circuit with an inductor and resistor only, whose differential equation is first-order and linear and is given by

$$LI' + RI = V,$$

where I is the to-be-determined current in the circuit, L measures the inductance, R measures the resistance, and V is the constant applied voltage. Find an equation describing the current in the circuit.

27. Suppose $a(t) > 0$, and $f(t) \rightarrow 0$ for $t \rightarrow \infty$. Show that every solution of the equation

$$\frac{dx}{dt} + a(t)x = f(t)$$

approaches 0 for $t \rightarrow \infty$.

28. In the same equation suppose that $a(t) > 0$, and let $x_0(t)$ be the solution for which the initial condition $x(0) = b$ is satisfied. Show that, for every positive $\varepsilon > 0$, there is a $\delta > 0$ such that, if we perturb the function $f(t)$ and the number b by a quantity less than δ , then the solution $x(t), t > 0$, is perturbed by less than ε . The word **perturbed** is understood in the following sense: $f(t)$ is replaced by $f_1(t)$ and b is replaced by b_1 where

$$|f_1(t) - f(t)| < \varepsilon, \quad |b_1 - b| < \delta.$$

This property of the solution $x(t)$ is called **stability for persistent disturbances**.

1.4 Some Physical Models Arising as Separable Equations

Now that we have studied separable equations in detail, we consider some applications. The wide variety of application problems that we will consider all lead to equations in which variables can be separated.

Free Fall, Neglecting Air Resistance

We will begin this application section with an easy problem from elementary physics. This application should be very familiar.

If $x(t)$ represents the position of a particle at time t , then the velocity of the particle is given by

$$v(t) = \frac{dx}{dt}.$$

Similarly, the acceleration of the particle is

$$a(t) = \frac{dv}{dt} = \frac{d^2x}{dt^2}.$$

Thus, if we consider a particle that is in free fall, where the acceleration of the particle is due to gravity alone, we have

$$a(t) = -g.$$

Here g is assumed to be a constant, and we use $-g$ as gravity acts downward. For the moment, we ignore the effects of air resistance. Thus,

$$\frac{dv}{dt} = -g,$$

which is a simple separable equation, so that

$$v(t) = -gt + c.$$

If we assume that the particle has an initial velocity v_0 , so that $v(0) = v_0$, then $v(t) = -gt + v_0$. Now this gives the separable equation

$$\frac{dx}{dt} = -gt + v_0$$

which has solution

$$x(t) = -\frac{g}{2}t^2 + v_0t + C_1.$$

If the particle has initial position x_0 , then

$$x(0) = x_0$$

which gives

$$x(t) = -\frac{g}{2}t^2 + v_0t + x_0 \quad (1.19)$$

as the position $x(t)$ of the particle in free fall, at time t .

Example 1 A man standing on a cliff 60 m high hurls a stone upward at a rate of 20 m/sec. How long does the stone remain in the air and with what speed does it hit the ground below the cliff?

Solution

Here $x_0 = 60$ and $v_0 = 20$. We take $g = 9.8$ m/sec². Thus,

$$x(t) = -\frac{9.8}{2}t^2 + 20t + 60$$

and

$$v(t) = -9.8t + 20.$$

The stone is in the air while $x(t) > 0$, so to find the time t that the stone is in the air, we set $x(t) = 0$ and solve for t . Using the quadratic equation,

$$t = \frac{-20 \pm \sqrt{(20)^2 - 4(-4.9)(60)}}{2(-4.9)} = -2.01, 6.09.$$

The stone is thus in the air for about 6.1 sec. We use this time to find the velocity upon impact:

$$v(6.1) = -9.8(6.1) + 20 = -39.78 \text{ m/sec.}$$

Air Resistance

We will now consider the effects of **air resistance**. The amount of air resistance (sometimes called the **drag force**) depends upon the size and velocity of the object, but there is no general law expressing this dependence. Experimental evidence shows that at very low velocities for small objects it is best to approximate the resistance R as proportional to the velocity, while for larger objects and higher velocities it is better to consider it as proportional to the square of the velocity [38].

By Newton's second law, $F = ma$, so that, if $v(t)$ is the velocity of the object, we have

$$m \frac{dv}{dt} = F_1 + F_2$$

where F_1 is the weight of the object,

$$F_1 = mg,$$

and F_2 is the force of the air resistance on the object as it falls, so

$$F_2 = k_1 v \quad \text{or} \quad F_2 = k_2 v^2$$

where k_1, k_2 are proportionality constants. Note that $k_i < 0$ because air resistance is always opposite the velocity; see Examples 2 and 3 below. We also point out that the units of k_1 and k_2 are different. In SI units, force has units of Newtons = N = kg · m/sec². Thus k_1 must have the units of kg/sec. On the other hand, k_2 can be written as

$$k_2 = -\frac{1}{2}C\rho A$$

where ρ is the air density (SI units of kg/m³), A is the cross-sectional area of the object (SI units of m²), and C is the drag coefficient (unitless) [38].

Example 2 An object weighing 8 pounds falls from rest toward earth from a great height. Assume that air resistance acts on it with a force equal to $2v$. Calculate the velocity $v(t)$ and position $x(t)$ at any time. Find and interpret $\lim_{t \rightarrow \infty} v(t)$.

Solution

Remembering that pounds is a **force** (not a mass), we see that we need to calculate the mass of the object in order to apply Newton's second law. Using $g = 32$ ft/sec² gives $m = w/g = 8/32 = 1/4$. Thus, by Newton's second law,

$$m \frac{dv}{dt} = F_1 + F_2,$$

that is

$$\frac{1}{4} \frac{dv}{dt} = 8 - 2v.$$

This is a separable equation and can be written as

$$\frac{dv}{8 - 2v} = 4dt$$

so that, upon integrating both sides, we have

$$-\frac{1}{2} \ln |8 - 2v| = 4t + c.$$

Using the condition that the object fell from rest, so that $v(0) = 0$, we can determine the constant c and solve for $v(t)$. We have

$$v(t) = 4 - 4e^{-8t}$$

as the velocity of the object at any time. A graph of this velocity is shown in Figure 1.6. Analytically, we see that $v(t)$ approaches 4 as $t \rightarrow \infty$. This value is known as the **limiting** or **terminal velocity** of the object.

Now since $\frac{dx}{dt} = v(t)$, we have

$$\frac{dx}{dt} = 4 - 4e^{-8t}.$$

This is easily integrated to obtain $x(t) = 4t + \frac{1}{2}e^{-8t} + c$. If we take the initial position of the object as zero, so that $x(0) = 0$, then

$$x(t) = 4t + \frac{1}{2}e^{-8t} - \frac{1}{2}.$$




FIGURE 1.6: Approach to terminal velocity of free-falling object of Example 2.

A Cool Problem

In addition to free-fall problems, separable equations arise in some simple thermodynamics applications. One such application is the following example.

Suppose that a pie is removed from a hot oven and placed in a cool room. After a given period of time, the pie has a temperature of 150 °F. We want to determine the time required to cool the pie to a temperature of 80 °F, when we can finally enjoy eating it.

This example is an application of **Newton's law of cooling**, which states

the rate at which the temperature $T(t)$ changes in a cooling body is proportional to the difference between the temperature of the body and the constant temperature T_s of the surrounding medium.

Symbolically we know the rate of change is the derivative, and the statement is expressed as

$$\frac{dT}{dt} = k(T - T_s), \quad (1.20)$$

with the initial temperature of the body $T(0) = T_0$ and k a constant of proportionality. We observe that, if the initial temperature T_0 is larger than the temperature of the surrounding T_s , then $T(t)$ will be a decreasing function of t (as the body is cooling), so $dT/dt < 0$, but $T_0 - T_s > 0$ so that the proportionality constant k must be negative. A similar analysis with $T_0 < T_s$ also gives $k < 0$. This condition on k also follows by noting that the temperature of the body will approach that of the surrounding medium as time gets large.

To solve (1.20), we seek a function $T(t)$ that describes the temperature at time t . For this equation, separating the variables, we have

$$\frac{dT}{T - T_s} = k dt.$$

Integrating both sides of this equation gives

$$\int \frac{dT}{T - T_s} = \int k dt.$$

Evaluating both integrals, we obtain

$$\ln |T - T_s| = kt + C,$$

where C is the constant of integration. Exponentiating both sides and simplifying gives

$$|T - T_s| = e^{kt} e^C \implies T - T_s = \pm e^C e^{kt}.$$

Solving for the temperature, we see that

$$T(t) = C_1 e^{kt} + T_s$$

where $C_1 = \pm e^C$. We can then apply the initial condition $T(0) = T_0$, which implies $T_0 = C_1 + T_s$, so that $C_1 = T_0 - T_s$ and the solution is then

$$T(t) = (T_0 - T_s)e^{kt} + T_s. \quad (1.21)$$

We know that the temperature of the body approaches that of its surroundings and this can be seen mathematically as

$$\lim_{t \rightarrow \infty} T(t) = T_s,$$

which is true because $k < 0$.

Let's now consider a specific pie-cooling example.

Example 3

Suppose that a pie is removed from a 350 °F oven and placed in a room with a temperature of 75 °F. In 15 min the pie has a temperature of 150 °F. We want to determine the time required to cool the pie to a temperature of 80 °F, when we can finally enjoy eating it.

Solution

Comparing with the above derivation, we see that $T_0 = 350$ and $T_s = 75$. Substituting these values in (1.21) gives

$$T(t) = 275e^{kt} + 75.$$

We still need to find k or equivalently e^k , which quantifies how fast the cooling of the pie occurs. We were given the temperature after 15 min, i.e., $T(15) = 150$. Thus

$$275e^{15k} + 75 = 150,$$

and solving for e^k gives

$$e^k = \left(\frac{3}{11} \right)^{1/15},$$

or $k = -0.08662$. Thus

$$T(t) = 275 \left(\frac{3}{11} \right)^{t/15} + 75,$$

and this can be used to find the temperature of the pie at any given time. We can also calculate the time it takes to cool to any given temperature. We want to know when $T(t) = 80$ °F. Thus we solve

$$275 \left(\frac{3}{11} \right)^{t/15} + 75 = 80$$

for t to obtain

$$t = \frac{-15 \ln 55}{\ln 3 - \ln 11} \approx 46.264.$$

Thus, the pie will reach a temperature of 80°F after approximately 46 min.

It is interesting to note that the first term in our equation for the pie temperature satisfies

$$275 \left(\frac{3}{11} \right)^{t/15} > 0$$

for all $t > 0$. Thus

$$T(t) = 275 \left(\frac{3}{11} \right)^{t/15} + 75 > 75.$$

The pie never actually reaches room temperature! This is an artifact of our model; we do note, however, that

$$\lim_{t \rightarrow \infty} 275 \left(\frac{3}{11} \right)^{t/15} + 75 = 75,$$

which can also be seen in Figure 1.7.




FIGURE 1.7: Graph of pie temperature vs. time of Example 3.

We present another example of Newton's law of cooling from forensic science.

Example 4 In the investigation of a homicide, the time of death is important. The normal body temperature of most healthy people is 98.6°F . Suppose that when a body is discovered at noon, its temperature is 82°F . Two hours later it is 72°F . If the temperature of the surroundings is 65°F , what was the approximate time of death?

Solution

This problem is solved as the last example. Here $T(0)$ represents the temperature when the body was discovered and $T(2)$ is the temperature of the body 2 hours later.

Thus, $T_0 = 82$ and $T_s = 65$ so that (1.21) becomes

$$T(t) = 17e^{kt} + 65.$$

Using $T(2) = 72$, we solve $17e^{2k} + 65 = 72$ for e^k to find

$$e^k = \left(\frac{7}{17} \right)^{1/2}$$

so that

$$T(t) = 17 \left(\frac{7}{17} \right)^{t/2} + 65.$$

This equation gives us the temperature of the body at any given time. To find the time of death, we use the fact that the body temperature was at 98.6°F at this time. Thus we solve

$$17 \left(\frac{7}{17} \right)^{t/2} + 65 = 98.6$$

for t and find that

$$t = \frac{2 \ln(1.97647)}{\ln 7 - \ln 17} \approx -1.53569.$$

This means that the time of death occurred approximately 1.53 hours before the body was discovered. Therefore, the time of death was approximately 10:30 a.m. because the body was found at noon.

Mixture Problems

Problems involving mixing typically give rise to separable differential equations. A typical mixture problem is given in the following example.

Example 5 A bucket contains 10 L of water and to it is being added a salt solution that contains 0.3 kg of salt per liter. This salt solution is being poured in at the rate of 2 L/min. The solution is being thoroughly mixed and drained off. The mixture is drained off at the same rate so that the bucket contains 10 L at all times. How much salt is in the bucket after 5 min?

Solution

Let $y(t)$ be the number of kilograms of salt in the bucket at the end of t minutes. We need to derive a differential equation for this problem, and we do so by considering change in this system over a small time interval. We first find the amount of salt added to the bucket between time t and time $t + \Delta t$. Each minute, 2 L of solution is added so that in Δt minutes, $2\Delta t$ liters is added.

In these $2\Delta t$ liters the amount of salt is

$$0.3 \text{ kg/L} \times (2\Delta t) \text{ L} = (0.6\Delta t) \text{ kg}.$$

On the other hand, $2\Delta t$ liters of solution is withdrawn from the bucket in an interval Δt . Now at time t the 10 L in the flask contains $y(t)$ kilograms of salt. Then $2\Delta t$ of these liters contains approximately $(0.2\Delta t)(y(t))$ kilograms of salt if we suppose that the change in the amount of salt $y(t)$ is small in the short period of time Δt .

We have computed the amount of salt added in the interval $(t, t + \Delta t)$, as well as the amount subtracted in the same interval. But the difference between the amounts of salt present at times $t + \Delta t$ and t is $y(t + \Delta t) - y(t)$, so that we have obtained the equation

$$y(t + \Delta t) - y(t) = 0.6\Delta t - (0.2\Delta t)(y(t)).$$

We now divide by Δt and let $\Delta t \rightarrow 0$. The left side approaches the derivative $y'(t)$, and the right side is $0.6 - 0.2y(t)$. The differential equation is thus

$$y'(t) = 0.6 - 0.2y(t), \tag{1.22}$$

which can be thought of as *the rate of change in the number of kilograms of salt in the bucket* $y'(t)$ *being equal to the rate of salt (in kg) flowing into the bucket* $0.6 (= 0.3 \text{ kg/L} \times 2 \text{ L})$ *minus the rate of salt flowing out of the bucket* $0.2y(t)$.

Equation (1.22) is a separable equation and can be written as

$$\frac{dy}{0.6 - 0.2y} = dt.$$

Integrating both sides gives

$$\ln |0.6 - 0.2y| = -0.2t + c$$

so that, solving for $y(t)$, we obtain

$$y(t) = 3 - Ce^{-0.2t}. \quad (1.23)$$

When t is zero, the amount of salt in the bucket is zero, that is, $y(0) = 0$. Equation (1.23) shows that, when $t = 0$, we have

$$y(0) = 3 - C$$

or $C = 3$. The value of C is now known, so that Equation (1.23) becomes

$$y(t) = 3 - 3e^{-0.2t}.$$

To find y at the end of 5 min, we simply substitute $t = 5$ so that the amount of salt in the bucket is $y(5) \approx 1.9$ kg. ■



Problems

In Problems 1–7, it will be convenient to take the velocity to be the unknown function.

1. A ball dropped from a building falls for 4.00 sec before it hits the ground. If air resistance is neglected, answer the following questions:
 - (a) What was its final velocity just as it hit the ground?
 - (b) What was the average velocity during the fall?
 - (c) How high was the building?
2. You drop a rock from a cliff, and 5.00 sec later you see it hit the ground. Neglecting air resistance, how high is the cliff?
3. A ball thrown straight up climbs for 3.0 sec before falling. Neglecting air resistance, with what velocity was the ball thrown?
4. Iron Man is flying at treetop level near Paris when he sees the Eiffel Tower elevator start to fall (the cable snapped). He knows Pepper Potts is inside. If Iron Man is 2 km away from the tower, and the elevator falls from a height of 350 m, how long does he have to save Pepper, and what must be his average velocity? Solve this problem assuming no air resistance. (Of course, Tony Stark instantly does the calculations required, as he is an expert in differential equations!)
5. The mass of a football is 0.4 kg. Air resists passage of the ball, the resistive force being proportional to the square of the velocity, and being equal to 0.004 N when the velocity is 1 m/sec. Find the height to which the ball will rise, and the time to reach that height if it is thrown upward with a velocity of 20 m/sec. How is the answer altered if air resistance is neglected?
6. The football of the preceding exercise is released (from rest) at an altitude of 17.1 m. Find its final velocity and time of fall.

7. Assume that air resistance is proportional to the square of velocity. The terminal velocity of a 75-kg human in air of standard density is 60 m/sec [38]. Neglecting the variation of air density with altitude and assuming that the 75-kg parachutist falls from an altitude of 1.8 km, find the velocity. Hint: use the terminal velocity to find the coefficient of v^2 .

Problems 8–12 concern Newton's law of cooling.

8. At the request of their children, Randy and Stephen make homemade popsicles. At 2:00 p.m., Kaelin asks if the popsicles are frozen (0°C), at which time they test the temperature of a popsicle and find it to be 5°C . If they put the popsicles with a temperature of 15°C in the freezer at 12:00 noon and the temperature of the freezer is -2°C , when will Erin, Kaelin, Robyn, Ryley, Alan, Abdi, and Avani be able to enjoy the popsicles?
9. An object cools in 10 min from 100°C to 60°C . The surroundings are at a temperature of 20°C . When will the object cool to 25°C ?
10. Determine the time of death if a corpse is 79°F when discovered at 3:00 p.m. and 68°F 3 hours later. Assume that the temperature of the surroundings is 60°F and that normal body temperature is 98.6°F .
11. A thermometer is taken from an inside room to the outside, where the air temperature is 5°F . After 1 minute the thermometer reads 55°F , and after 5 minutes it reads 30°F . Determine the initial temperature of the inside room.
12. A slug of metal at a temperature of 800°F is put in an oven, the temperature of which is gradually increased during an hour from a° to b° . Find the temperature of the metal at the end of an hour, assuming that the metal warms kT degrees per minute when it finds itself in an oven that is T degrees warmer.

In Problems 13–17, it is supposed that the amount of gas (or liquid) contained in any fixed volume is constant. Also, thorough mixing is assumed.

13. A 20-L vessel contains air (assumed to be 80% nitrogen and 20% oxygen). Suppose 0.1 L of nitrogen is added to the container per second. If continual mixing takes place and material is withdrawn at the rate at which it is added, how long will it be before the container holds 99% nitrogen?
14. A 100-L beaker contains 10 kg of salt. Water is added at the constant rate of 5 L/min with complete mixing and drawn off at the same rate. How much salt is in the beaker after 1 hour?
15. A tank contains 25 lb of salt dissolved in 50 gal of water. Brine containing 4 lb/gal is allowed to enter at a rate of 2 gal/min. If the solution is drained at the same rate, find the amount of salt as a function $S(t)$ of time t . Find the concentration of salt at time t . Suppose the rate of draining is modified to be 3 gal/min. Find the amount of salt and the concentration at time t .
16. Consider a pond that has an initial volume of $10,000\text{ m}^3$. Suppose that at time $t = 0$, the water in the pond is clean and that the pond has two streams flowing into it, stream A and stream B, and one stream flowing out, stream C. Suppose $500\text{ m}^3/\text{day}$ of water flows into the pond from stream A, $750\text{ m}^3/\text{day}$ flows into the pond from stream B, and 1250 m^3 flows out of the pond via stream C. At $t = 0$, the water flowing into the pond from stream A becomes contaminated with road salt at a concentration of $5\text{ kg}/1000\text{ m}^3$. Suppose the water in the pond is well mixed so the concentration of salt at any given time is constant. To make matters worse, suppose also that at time

$t = 0$ someone begins dumping trash into the pond at a rate of $50 \text{ m}^3/\text{day}$. The trash settles to the bottom of the pond, reducing the volume by $50 \text{ m}^3/\text{day}$. To adjust for the incoming trash, the rate that water flows out via stream C increases to $1300 \text{ m}^3/\text{day}$ and the banks of the pond do not overflow. Determine how the amount of salt in the pond changes over time. Does the amount of salt in the pond reach 0 after some time has passed?

17. A large chamber contains 200 m^3 of gas, 0.15% of which is carbon dioxide (CO_2). A ventilator exchanges $20 \text{ m}^3/\text{min}$ of this gas with new gas containing only 0.04% CO_2 . How long will it be before the concentration of CO_2 is reduced to half its original value?

Problems 18–20 concern radioactive decay. The decay law states that the amount of radioactive substance that decays is proportional at each instant to the amount of substance present.

18. The strength of a radioactive substance decreases 50% in a 30-day period. How long will it take for the radioactivity to decrease to 1% of its initial value?
19. It is experimentally determined that every gram of radium loses 0.44 mg in 1 year. What length of time elapses before the radioactivity decreases to half its original value?
20. A tin organ pipe decays with age as a result of a chemical reaction that is catalyzed by the decayed tin. As a result, the rate at which the tin decays is proportional to the product of the amount of tin left and the amount that has already decayed. Let M be the total amount of tin before any has decayed. Find the amount of decayed tin $p(t)$.

Problems 21 and 22 deal with geometric situations where the derivative arises and yields a separable equation.

21. Find a curve for which the area of the triangle determined by the tangent, the ordinate to the point of tangency, and the x -axis has a constant value equal to a^2 .
22. Find a curve for which the sum of the sides of a triangle constructed as in the previous problem has a constant value equal to b .
23. On an early Monday morning in February in rural Kentucky (not far from Western Kentucky University) it started to snow. There had been no snow on the ground before. It was snowing at a steady, constant rate so that the thickness of the snow on the ground was increasing at a constant rate. A snowplow began clearing the snow from the streets at noon. The speed of the snowplow in clearing the snow is inversely proportional to the thickness of the snow. The snowplow traveled 2 miles during the first hour after noon and traveled 1 mile during the second hour after noon. At what time did it begin snowing?

1.5 Exact Equations

We will now introduce another type of differential equation. Exact equations are not separable equations nor are they necessarily linear. They come up in higher-level math in fields such as potential theory and harmonic analysis.

Consider the first-order differential equation $\frac{dy}{dx} = f(x, y)$. We observe that it can always be expressed in the differential form

$$M(x, y) dx + N(x, y) dy = 0$$

or equivalently as

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0$$

and vice versa. We will now consider a type of differential equation that is not separable, but, nevertheless, has a solution. We need a definition from multivariable calculus to proceed:

Definition 1.5.1

Let $F(x, y)$ be a function of two real variables such that F has continuous first partial derivatives in a domain D . The total differential dF of F is defined by

$$dF(x, y) = \frac{\partial F(x, y)}{\partial x} dx + \frac{\partial F(x, y)}{\partial y} dy$$

for all $(x, y) \in D$.

Example 1 Suppose $F(x, y) = xy^2 + 2x^3y$; then

$$\frac{\partial F}{\partial x} = y^2 + 6x^2y \quad \text{and} \quad \frac{\partial F}{\partial y} = 2xy + 2x^3$$

so that the total differential dF is given by

$$\begin{aligned} dF(x, y) &= \frac{\partial F(x, y)}{\partial x} dx + \frac{\partial F(x, y)}{\partial y} dy \\ &= (y^2 + 6x^2y) dx + (2xy + 2x^3) dy. \end{aligned}$$

Definition 1.5.2

The expression

$$M(x, y) dx + N(x, y) dy \tag{1.24}$$

is called an exact differential in a domain D if there exists a function F of two real variables such that this expression equals the total differential $dF(x, y)$ for all $(x, y) \in D$. That is, (1.24) is an exact differential in D if there exists a function F such that

$$\frac{\partial F}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x, y)$$

for all $(x, y) \in D$.

If $M(x, y) dx + N(x, y) dy$ is an exact differential, then the differential equation

$$M(x, y) dx + N(x, y) dy = 0 \tag{1.25}$$

is called an *exact differential equation*. As long as $x = C$ (a constant) is not a solution, we consider the equivalent form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{1.26}$$

as the standard form for an exact equation.

Example 2 The differential equation

$$y^2 + 2xy \frac{dy}{dx} = 0$$

is exact, since, if $F(x, y) = xy^2$, then

$$\frac{\partial F}{\partial x} = y^2 \text{ and } \frac{\partial F}{\partial y} = 2xy.$$

Not all differential equations, however, are exact. Consider

$$y + 2x \frac{dy}{dx} = 0.$$

We cannot find an $F(x, y)$ so that

$$\frac{\partial F}{\partial x} = y \text{ and } \frac{\partial F}{\partial y} = 2x.$$

Numerous trials and errors may be enough to convince us that this is the case. What we really need is a method for testing a differential equation for exactness and for constructing the corresponding function $F(x, y)$. Both are contained in the following theorem and its proof.

THEOREM 1.5.1

Consider the differential equation

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (1.27)$$

where M and N have continuous first partial derivatives at all points (x, y) in a rectangular domain D . Then the differential equation (1.27) is exact in D , **if and only if**

$$\frac{\partial M(x, y)}{\partial y} = \frac{\partial N(x, y)}{\partial x} \quad (1.28)$$

for all (x, y) in D .

Remark: The proof of this theorem is rather important, as it not only provides a test for exactness, but also a method of solution for exact differential equations.

Proof: To prove one direction of the theorem, we first suppose the differential equation (1.27) is exact in D and show that (1.28) must hold as a result. If (1.27) is exact, then there is a function F such that

$$\frac{\partial F}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x, y).$$

So

$$\frac{\partial^2 F}{\partial y \partial x} = \frac{\partial M}{\partial y} \quad \text{and} \quad \frac{\partial^2 F}{\partial x \partial y} = \frac{\partial N}{\partial x}$$

by differentiation. Now we have assumed the continuity of the first partials of M and N in D , so that

$$\frac{\partial^2 F}{\partial y \partial x} = \frac{\partial^2 F}{\partial x \partial y}.$$

This means that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x},$$

which is the same as (1.28).

To prove the other direction, we assume (1.28) and show that (1.27) must be exact. (Proving this direction will also show us how to construct the solution for a given exact equation.) Thus, we assume

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

and find an F so that

$$\frac{\partial F}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x, y). \quad (1.29)$$

It is clear that we can find an F that satisfies either of these equations, but can we find an F that satisfies both? Let's proceed and see what happens. Suppose that F satisfies

$$\frac{\partial F}{\partial x} = M(x, y).$$

We can integrate both sides of this equation to get

$$F(x, y) = \int M(x, y) dx + \phi(y) \quad (1.30)$$

where $\int M(x, y) dx$ is the partial integration with respect to x , holding y constant. Note that our “constant” of integration, $\phi(y)$, is a function but is a function of y only (it might also include an additive constant, but definitely no x). This is because the expression $\partial F/\partial x$ would result in the loss of any “only y functions.” Now we need to find an $F(x, y)$ that satisfies both equations in (1.29). We thus need to make sure the $F(x, y)$ in (1.30) also satisfies $\frac{\partial F}{\partial y} = N(x, y)$. We calculate $\partial F/\partial y$ by differentiating (1.30) with respect to y :

$$\frac{\partial F}{\partial y} = \frac{\partial}{\partial y} \int M(x, y) dx + \frac{d\phi(y)}{dy}.$$

Equating with $N(x, y)$ gives

$$N(x, y) = \left(\frac{\partial}{\partial y} \int M(x, y) dx \right) + \phi'(y),$$

where $\phi'(y) = d\phi(y)/dy$. Solving for $\phi'(y)$ gives

$$\phi'(y) = N(x, y) - \frac{\partial}{\partial y} \int M(x, y) dx.$$

Since $\phi(y)$ is a function of only y , it must also be the case that $\phi'(y)$ is a function of only y . We can see this by showing

$$\frac{\partial}{\partial x} \left(N(x, y) - \frac{\partial}{\partial y} \int M(x, y) dx \right) = 0.$$

Evaluating the left-hand side and simplifying gives

$$\begin{aligned}
 \frac{\partial}{\partial x} \left(N(x, y) - \frac{\partial}{\partial y} \int M(x, y) dx \right) &= \frac{\partial N}{\partial x} - \frac{\partial^2}{\partial x \partial y} \int M(x, y) dx \\
 &= \frac{\partial N}{\partial x} - \frac{\partial^2 F}{\partial x \partial y} \text{ (by noting what } F \text{ is)} \\
 &= \frac{\partial N}{\partial x} - \frac{\partial^2 F}{\partial y \partial x} \text{ (by continuity)} \\
 &= \frac{\partial N}{\partial x} - \frac{\partial^2}{\partial y \partial x} \int M(x, y) dx \\
 &= \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \\
 &= 0,
 \end{aligned}$$

where the last equality holds since we have assumed that

$$\frac{\partial N}{\partial x} = \frac{\partial M}{\partial y}.$$

What this means is that

$$N(x, y) - \frac{\partial}{\partial y} \int M(x, y) dx$$

cannot depend on x since its derivative with respect to x is zero. Hence,

$$\phi(y) = \int \left(N(x, y) - \frac{\partial}{\partial y} \int M(x, y) dx \right) dy$$

and thus

$$F(x, y) = \int M(x, y) dx + \phi(y)$$

is a function that satisfies both

$$\frac{\partial F}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x, y).$$

Thus,

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0$$

is exact in D . ■

In short, the criterion for exactness is (1.28):

$$\frac{\partial N}{\partial x} = \frac{\partial M}{\partial y}.$$

If this equation holds, then the differential equation is exact. If this is not true, the differential equation is not exact.

Example 3 We considered the differential equation

$$y^2 + 2xy \frac{dy}{dx} = 0 \tag{1.31}$$

earlier. We see that

$$M(x, y) = y^2 \quad \text{and} \quad N(x, y) = 2xy.$$

Thus,

$$\frac{\partial M}{\partial y} = 2y = \frac{\partial N}{\partial x},$$

so that the differential equation is exact. On the other hand,

$$y + 2x \frac{dy}{dx} = 0 \tag{1.32}$$

gives $M(x, y) = y$ and $N(x, y) = 2x$ so that

$$\frac{\partial M}{\partial y} = 1 \neq 2 = \frac{\partial N}{\partial x}.$$

Hence $y + 2x \frac{dy}{dx} = 0$ is not exact.

Example 4 Consider the differential equation

$$(2x \sin y + y^3 e^x) + (x^2 \cos y + 3y^2 e^x) \frac{dy}{dx} = 0.$$

Here

$$M(x, y) = 2x \sin y + y^3 e^x \quad \text{and} \quad N(x, y) = x^2 \cos y + 3y^2 e^x;$$

hence

$$\frac{\partial M}{\partial y} = 2x \cos y + 3y^2 e^x = \frac{\partial N}{\partial x}.$$

Thus the differential equation is exact.

Remark: The test for exactness applies to equations in the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0. \tag{1.33}$$

If the left-hand side is an exact differential, then we can solve the exact differential equation (1.33) by finding a function $F(x, y)$ so that

$$\frac{\partial F(x, y)}{\partial x} dx + \frac{\partial F(x, y)}{\partial y} dy = 0.$$

More simply, using the total differential, we obtain $dF(x, y) = 0$. Thus,

$$F(x, y) = C$$

is a solution to (1.33).

THEOREM 1.5.2

Suppose the differential equation

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0$$

is exact. Then the general solution of this differential equation is given implicitly by

$$F(x, y) = C,$$

where $F(x, y)$ is a function such that

$$\frac{\partial F}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x, y).$$

Remark 1: As with separable and homogeneous equations, the constant in Theorem 1.5.2 is determined by an initial condition.

Remark 2: We have an explicit form for $F(x, y)$, namely,

$$F(x, y) = \int M(x, y) dx + \phi(y),$$

where $\phi(y) = \int \left(N(x, y) - \frac{\partial}{\partial y} \int M(x, y) dx \right) dy$. This form, however, is not always useful. We will see by example how to solve exact differential equations.

Remark 3: We integrated $\partial F/\partial x = M$ and substituted this into $\partial F/\partial y = N$. We instead could have solved $\partial F/\partial y = N$ first (by integrating with respect to y and obtaining a “constant” $\psi(x)$) and then substituted into $\partial F/\partial x = M$. The resulting F is the same but would be written

$$F(x, y) = \int N(x, y) dy + \psi(x), \quad (1.34)$$

where $\psi(x) = \int \left(M(x, y) - \frac{\partial}{\partial x} \int N(x, y) dy \right) dx$. See Problem 19 at the end of this section.

Example 5 Show that

$$(3x^2 + 4xy) + (2x^2 + 2y) \frac{dy}{dx} = 0$$

is exact and then solve it by the methods discussed in this section.

Solution

We have

$$M(x, y) = 3x^2 + 4xy \quad \text{and} \quad N(x, y) = 2x^2 + 2y$$

so that the equation is exact, since

$$\frac{\partial M}{\partial y} = 4x = \frac{\partial N}{\partial x}.$$

Our goal is to find an $F(x, y)$ that simultaneously satisfies the equations

$$\frac{\partial F}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x, y).$$

That is, F must satisfy

$$\frac{\partial F}{\partial x} = 3x^2 + 4xy \quad \text{and} \quad \frac{\partial F}{\partial y} = 2x^2 + 2y.$$

Integrating $\partial F/\partial x$ with respect to x gives

$$\begin{aligned} F(x, y) &= \int (3x^2 + 4xy) dx \\ &= x^3 + 2x^2y + \phi(y). \end{aligned}$$

This same F must also satisfy $\partial F/\partial y = N$ and we then have

$$2x^2 + \phi'(y) = \frac{\partial F}{\partial y} = 2x^2 + 2y.$$

Thus, $\phi'(y) = 2y$. Integrating with respect to y gives

$$\phi(y) = y^2 + C_0$$

so that

$$F(x, y) = x^3 + 2x^2y + y^2 + C_0.$$

Thus, a one-parameter family of solutions is given by

$$x^3 + 2x^2y + y^2 = C.$$

We now solve an exact equation by first integrating with respect to y ; see Remark 3 above.

Example 6 Show that

$$(2x \cos y + 3x^2y) + (x^3 - x^2 \sin y - y) \frac{dy}{dx} = 0$$

is exact and solve it subject to the initial condition $y(0) = 2$. Plot the solution.

Solution

We have $M(x, y) = 2x \cos y + 3x^2y$ and $N(x, y) = x^3 - x^2 \sin y - y$. The equation is exact because

$$\frac{\partial M}{\partial y} = 3x^2 - 2x \sin y = \frac{\partial N}{\partial x}.$$

Now we find an $F(x, y)$ so that

$$\frac{\partial F}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial F}{\partial y} = N(x, y).$$

This time we will integrate $\partial F/\partial y = N$ with respect to y . Thus

$$\begin{aligned} F(x, y) &= \int N(x, y) dy \\ &= \int (x^3 - x^2 \sin y - y) dy \\ &= x^3y + x^2 \cos y - \frac{y^2}{2} + \psi(x). \end{aligned}$$

This must also satisfy $\partial F/\partial x = M$. Calculating $\partial F/\partial x$ gives

$$\frac{\partial F}{\partial x} = 3x^2y + 2x \cos y + \psi'(x).$$

Substituting into

$$\frac{\partial F}{\partial x} = M(x, y)$$

gives $\psi'(x) = 0$, which is easily integrated to obtain $\psi(x) = C_1$. Thus,

$$F(x, y) = x^3y + x^2 \cos y - \frac{1}{2}y^2 + C_1,$$

and a one-parameter family of solutions is

$$x^3y + x^2 \cos y - \frac{1}{2}y^2 = C.$$

The initial condition $y(0) = 2$ gives $C = -2$. Hence

$$x^2 \cos y + x^3y - \frac{1}{2}y^2 = -2$$

is the implicit solution that satisfies the given initial condition. The solution curves can be plotted, as shown in Figure 1.8.




FIGURE 1.8: Implicit plot for Example 6. The upper curve is the solution curve because it passes through the initial condition.

Note that, although both curves in Figure 1.8 satisfy the implicit equation, only one of these curves passes through the given initial condition and thus is the correct solution. ■

Solution by Grouping

There is a much slicker method for solving exact differential equations and it is known as the **method of grouping**. For better or worse, it requires a “working knowledge” of differentials and a certain amount of ingenuity. We again consider Example 5, this time in its differential form:

$$(3x^2 + 4xy) dx + (2x^2 + 2y) dy = 0.$$

We rewrite it in the form

$$3x^2 dx + (4xy dx + 2x^2 dy) + 2y dy = 0$$

which is

$$d(x^3) + d(2x^2y) + d(y^2) = d(C).$$

That is,

$$d(x^3 + 2x^2y + y^2) = d(C)$$

so that

$$x^3 + 2x^2y + y^2 = C.$$

Clearly, this procedure is much quicker if we can find the appropriate grouping. Let's try this method one more time by again considering Example 6. We group the terms as

$$(2x \cos y \, dx - x^2 \sin y \, dy) + (3x^2y \, dx + x^3 \, dy) - y \, dy = 0.$$

Thus, we have

$$d(x^2 \cos y) + d(x^3y) - d\left(\frac{y^2}{2}\right) = d(C)$$

and so

$$x^2 \cos y + x^3y - \frac{1}{2}y^2 = C$$

is a one-parameter family of solutions.

Important Note: If we use the method of grouping, we still need to check that the equation is exact for our first step.



Problems

In Problems 1–13, check to see if the equation is exact. If it is, solve it by the methods of this section. If an initial condition is given, graph the solution.

1. $(1 + xy^2) + (1 + x^3y) \frac{dy}{dx} = 0$
2. $(1 + y^2 \sin 2x) - 2y \cos 2x \frac{dy}{dx} = 0$
3. $2xy + (x^2 - y^2) \frac{dy}{dx} = 0$
4. $(1 + y^2 \sin 2x) - y \cos 2x \frac{dy}{dx} = 0$
5. $2xy^3 + (1 + 3x^2y^2) \frac{dy}{dx} = 0$
6. $(2 + \frac{y}{x^2}) dx + (y - \frac{1}{x}) dy = 0$
7. $3x^2(1 + \ln y) = (2y - \frac{x^3}{y}) \frac{dy}{dx}$
8. $(\frac{x}{\sin y} + 2) + \frac{(x^2+1) \cos y}{\cos 2y - 1} \frac{dy}{dx} = 0$
9. $(2xy + 1) + (x^2 + 4y) \frac{dy}{dx} = 0, y(0) = 1$
10. $(2y \sin x \cos x + y^2 \sin x) + (\sin^2 x - 2y \cos x) \frac{dy}{dx} = 0, y(0) = 3$
11. $(2 - 9xy^2)x + (4y^2 - 6x^3)y \frac{dy}{dx} = 0, y(1) = 1$
12. $(y \sec^2 x + \sec x \tan x) + (\tan x + 2y) \frac{dy}{dx} = 0, y(0) = 1$
13. $e^{-y} - (2y + xe^{-y}) \frac{dy}{dx} = 0, y(1) = 3$

In Problems 14 and 15, determine the constant A such that the equation is exact. Then solve the resulting exact equation.

14. $(x^2 + 3xy) + (Ax^2 + 4y) \frac{dy}{dx} = 0$
15. $(\frac{Ay}{x^3} + \frac{y}{x^2}) + (\frac{1}{x^2} - \frac{1}{x}) \frac{dy}{dx} = 0$

In Problems 16 and 17, determine the most general function $(N(x, y) \text{ or } M(x, y))$ that makes the equation exact.

16. $M(x, y) + (2ye^x + y^2e^{3x}) \frac{dy}{dx} = 0$
17. $(x^3 + xy^2) + N(x, y) \frac{dy}{dx} = 0$

18. Let x represent the units of labor and y represent the units of capital. If $f(x, y)$ measures the number of units produced, a differential equation satisfied by a level curve of it is

$$ax^{a-1}y^{1-a} + (1-a)x^ay^{-a} \frac{dy}{dx} = 0.$$

Solve this equation as (i) a separable equation and (ii) an exact equation. In doing (ii), we obtain the well-known **Cobb–Douglas production function** $f(x, y) = Cx^ay^{1-a}$.

19. By following the proof of Theorem 1.5.1, show that an equivalent formulation of $F(x, y)$ is given by

$$F(x, y) = \int N(x, y) \frac{dy}{dx} + \int \left(M(x, y) - \frac{\partial}{\partial x} \int N(x, y) \frac{dy}{dx} \right) dx.$$

Although this could easily be obtained by rearranging the previously obtained expression for F (1.30), do *not* simply rearrange terms.

20. By using the substitution $y = vx$, show that the homogeneous equation

$$(Ax + By) + (Cx + Ey) \frac{dy}{dx} = 0,$$

where A, B, C , and E are constants, is exact if and only if $B = C$.

21. By using the substitution $y = vx$, show that the homogeneous equation

$$(Ax^2 + Bxy + Cy^2) + (Ex^2 + Fxy + Gy^2) \frac{dy}{dx} = 0,$$

where A, B, C, E, F , and G are constants, is exact if and only if $B = 2E$ and $F = 2C$.

1.6 Special Integrating Factors and Substitution Methods

Special Integrating Factors

In solving linear equations, we learned that we could multiply by an appropriate **integrating factor**, thus transforming the equation into a form we can solve. Besides the one we learned, there are other integrating factors that we will now consider.

Definition 1.6.1

If the differential equation

$$M(x, y) dx + N(x, y) dy = 0 \tag{1.35}$$

is not exact in a domain D but the differential equation

$$\mu(x, y)M(x, y) dx + \mu(x, y)N(x, y) dy = 0 \tag{1.36}$$

is exact in D , then $\mu(x, y)$ is called an integrating factor of the differential equation (1.35).

Example 1 The differential equation

$$(3y + 4xy^2) dx + (2x + 3x^2y) dy = 0 \tag{1.37}$$

is not exact since

$$\frac{\partial M}{\partial y} = 3 + 8xy \neq 2 + 6xy = \frac{\partial N}{\partial x}.$$

If we let $\mu(x, y) = x^2y$, we can use (1.36) to rewrite (1.37) as

$$(x^2y)(3y + 4xy^2) dx + (x^2y)(2x + 3x^2y) dy = 0.$$

Expanding gives

$$M = 3x^2y^2 + 4x^3y^3 \quad \text{and} \quad N = 2x^3y + 3x^4y^2.$$

Then

$$\frac{\partial}{\partial y}(3x^2y^2 + 4x^3y^3) = 6x^2y + 12x^3y^2 = \frac{\partial}{\partial x}(2x^3y + 3x^4y^2).$$

Thus the new equation is exact and hence $\mu(x, y) = x^2y$ is an integrating factor. ■

We saw above how multiplying by an appropriate integrating factor converted a linear equation into an exact equation, which we could then solve. Multiplying by an appropriate integrating factor is a technique that will work in other situations as well.

We have seen that if the equation

$$M(x, y) dx + N(x, y) dy = 0$$

is not exact and if $\mu(x, y)$ is an integrating factor, then the differential equation

$$\mu(x, y)M(x, y) dx + \mu(x, y)N(x, y) dy = 0$$

is exact. Using the criterion for exactness, we must have

$$\frac{\partial}{\partial y}(\mu(x, y)M(x, y)) = \frac{\partial}{\partial x}(\mu(x, y)N(x, y)).$$

To simplify notation, we will write M, N instead of $M(x, y), N(x, y)$ when taking the partial derivatives, even though both M and N are functions of x and y . The criterion for exactness can then be written

$$\frac{\partial \mu}{\partial y} M(x, y) + \mu(x, y) \frac{\partial M}{\partial y} = \frac{\partial \mu}{\partial x} N(x, y) + \mu(x, y) \frac{\partial N}{\partial x}.$$

Rearranging gives

$$\frac{\partial \mu}{\partial y} M(x, y) - \frac{\partial \mu}{\partial x} N(x, y) = \mu(x, y) \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right). \quad (1.38)$$

Thus $\mu(x, y)$ is an integrating factor if and only if it is a solution of the partial differential equation (1.38). We will not consider the solution of this partial differential equation. We will instead consider (1.38) in the case where μ only depends on x , i.e., $\mu(x, y) = \mu(x)$. (We can also consider the case when $\mu(x, y) = \mu(y)$ and the analogous formulation is left as one of the exercises.) In this situation, (1.38) reduces to

$$-\mu'(x) N(x, y) = \mu(x) \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right).$$

That is,

$$\frac{1}{\mu} \frac{d\mu}{dx} = \frac{1}{N(x, y)} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right). \quad (1.39)$$

If the right-hand side of (1.39) involves two dependent variables, we run into trouble. If, however, it depends only upon x , then Equation (1.39) is separable, in which case we obtain

$$\mu(x) = \exp \left[\int \frac{1}{N(x, y)} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) dx \right]$$

as an integrating factor.

Example 2 Solve the differential equation

$$(2x^2 + y) dx + (x^2y - x) dy = 0.$$

Solution

In this equation,

$$M(x, y) = 2x^2 + y \quad \text{and} \quad N(x, y) = x^2y - x$$

so that

$$\frac{\partial M}{\partial y} = 1 \neq 2xy - 1 = \frac{\partial N}{\partial x}$$

and the equation is not exact. It can also be shown (try it!) that the differential equation is not separable, homogeneous, or linear. Now

$$\begin{aligned} \frac{1}{N(x, y)} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) &= \frac{1}{x^2y - x} (1 - (2xy - 1)) \\ &= \frac{-2}{x} \end{aligned}$$

depends only upon x . Thus,

$$\mu(x) = \exp \left(- \int \frac{2}{x} dx \right) = e^{-2 \ln |x|} = \frac{1}{x^2}$$

is an integrating factor. If we multiply the equation through by this factor, we have

$$\left(2 + \frac{y}{x^2} \right) dx + \left(y - \frac{1}{x} \right) dy = 0.$$

Now this equation is exact since

$$\frac{\partial M}{\partial y} = \frac{1}{x^2} = \frac{\partial N}{\partial x}.$$

We can thus solve this differential equation using the exact method to obtain

$$2x + \frac{y^2}{2} - \frac{y}{x} = C.$$

1.6.1 Bernoulli Equation

We will now consider a class of differential equations that can be reduced to linear equations by an appropriate transformation. These equations are called *Bernoulli* equations and often arise in applications.

Definition 1.6.2

A first-order differential equation of the form

$$\frac{dy}{dx} + P(x)y = Q(x)y^n \quad n \in \mathbb{R} \quad (1.40)$$

is called a Bernoulli differential equation.

Note that, when $n = 0$ or $n = 1$, the Bernoulli equation is actually a linear equation and can be solved as such. When $n \neq 0$ or 1 , then we must consider an additional method.

THEOREM 1.6.1

Suppose $n \neq 0$ or 1 ; then the transformation

$$v = y^{1-n}$$

reduces the Bernoulli equation (1.40) to

$$\frac{dv}{dx} + (1-n)P(x)v = (1-n)Q(x), \quad (1.41)$$

which is a linear equation in v .

Proof: Multiply the Bernoulli equation by y^{-n} and thus obtain

$$y^{-n} \frac{dy}{dx} + P(x)y^{1-n} = Q(x). \quad (1.42)$$

Now let $v = y^{1-n}$ so that

$$\frac{dv}{dx} = (1-n)y^{-n} \frac{dy}{dx}.$$

Hence, Equation (1.42) becomes

$$\frac{1}{1-n} \frac{dv}{dx} + P(x)v = Q(x),$$

that is,

$$\frac{dv}{dx} + (1-n)P(x)v = (1-n)Q(x).$$

Letting

$$P_1(x) = (1-n)P(x) \quad \text{and} \quad Q_1(x) = (1-n)Q(x)$$

gives

$$\frac{dv}{dx} + P_1(x)v = Q_1(x),$$

a linear differential equation in v . ■

Example 3 Solve the differential equation

$$\frac{dy}{dx} + y = xy^3.$$

Solution

This is a Bernoulli equation with $n = 3$. We thus let $v = y^{1-3} = y^{-2}$, so that

$$\frac{dv}{dx} = -2y^{-3} \frac{dy}{dx}.$$

Using (1.41) we obtain

$$\frac{dv}{dx} - 2v = -2x. \quad (1.43)$$

This is a linear differential equation with integrating factor

$$\exp\left(\int P(x) dx\right) = \exp\left(\int -2 dx\right) = e^{-2x}.$$

We also calculate $\exp(-\int P(x) dx) = e^{2x}$. Thus the solution of (1.43) can be written

$$v = e^{2x} \left(\int -2xe^{-2x} dx \right).$$

Integrating by parts gives

$$v = e^{2x} \left(xe^{-2x} + \frac{1}{2}e^{-2x} \right) + Ce^{2x}.$$

Simplifying gives

$$v = x + \frac{1}{2} + Ce^{2x}.$$

But our original problem was in the variable y . We know $v = y^{-2}$ and thus the solution is

$$\frac{1}{y^2} = x + \frac{1}{2} + Ce^{2x}$$

which can be written as

$$y = \pm \left(\frac{1}{x + \frac{1}{2} + Ce^{2x}} \right)^{1/2}.$$

This solution is defined as long as the denominator is not equal to zero. ■

1.6.2 Homogeneous Equations of the Form $g(y/x)$

We have now been introduced to separable differential equations and their relative ease of solution. We will now consider a class of differential equations that can be reduced to separable equations by a change of variables.

Remark: Before proceeding, we alert the reader that the use of the word *homogeneous* in this section must not be confused with its use as the type of linear ordinary differential equation whose right-hand side is zero (as in Chapters 3 and 4). Its use in those chapters is more common, but both have their place.

Example 4 Consider the differential equation

$$\frac{dy}{dx} = \frac{x - y}{x + y}.$$

Solution

After a minute or so of reflection, we see that this is not a separable equation. We can, however, rewrite the equation as

$$\frac{dy}{dx} = \frac{1 - \frac{y}{x}}{1 + \frac{y}{x}} \quad (1.44)$$

so that we can isolate the fraction y/x . This suggests we consider the change of variable

$$v = \frac{y}{x}$$

or equivalently

$$y = vx.$$

Our original problem has dy/dx and thus we take the derivative of both sides of the above equation with respect to x to get

$$\frac{dy}{dx} = v + x \frac{dv}{dx}.$$

Substitution of this and $y = vx$ into (1.44) gives

$$v + x \frac{dv}{dx} = \frac{1 - v}{1 + v}.$$

Simplifying results in the separable equation

$$x \frac{dv}{dx} = \frac{1 - 2v - v^2}{1 + v},$$

and we separate its variables as

$$\frac{1 + v}{1 - 2v - v^2} dv = \frac{dx}{x},$$

and integrate to give

$$\ln |1 - 2v - v^2| = -2 \ln x + C_1.$$

Exponentiation of both sides yields

$$|1 - 2v - v^2| = e^{C_1} x^{-2} = C_2 x^{-2}.$$

But, $v = y/x$, so that substitution gives

$$1 - \frac{2y}{x} - \left(\frac{y}{x}\right)^2 = \pm C_2 x^{-2} = C x^{-2}.$$

Multiplying by x^2 to clear the fraction gives

$$x^2 - 2xy - y^2 = C$$

as the implicit solution to the differential equation.

This is an example of a general method of reducing a class of differential equations to that of a separable equation. We need some terminology.

Definition 1.6.3

The first-order differential equation

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0$$

is said to be of homogeneous type (or homogeneous) if, when written in the derivative form

$$\frac{dy}{dx} = f(x, y),$$

there exists a function g such that $f(x, y)$ can be expressed in the form $g(y/x)$.

By classifying the equation as homogeneous, we will be able to apply the above technique in order to reduce the differential equation to one that is separable. It is sometimes not obvious that a given equation can be rewritten as a homogeneous equation. We present two examples now to help clarify this concept.

Example 5 The differential equation

$$(x^2 - 3y^2) + 2xy \frac{dy}{dx} = 0$$

is homogeneous, since the equation can be written in derivative form as

$$\frac{dy}{dx} = \frac{3y^2 - x^2}{2xy},$$

and we can rearrange this as

$$\frac{3y^2 - x^2}{2xy} = \frac{3}{2} \left(\frac{y}{x} \right) - \frac{1}{2} \left(\frac{1}{y/x} \right)$$

so that

$$\frac{dy}{dx} = \frac{3}{2} \left(\frac{y}{x} \right) - \frac{1}{2} \left(\frac{1}{y/x} \right).$$

The right-hand side is of the form $g(y/x)$ for the function

$$g(z) = \frac{3z}{2} - \frac{1}{2z},$$

and so the differential equation is homogeneous. ■

Example 6 The differential equation

$$\left(y + \sqrt{x^2 + y^2} \right) dx - x dy = 0$$

can be written as

$$\frac{dy}{dx} = \frac{y + \sqrt{x^2 + y^2}}{x}.$$

For $x > 0$, we have

$$\frac{y + \sqrt{x^2 + y^2}}{x} = \frac{y}{x} + \sqrt{1 + \left(\frac{y}{x} \right)^2},$$

so

$$\frac{dy}{dx} = \frac{y}{x} + \sqrt{1 + \left(\frac{y}{x}\right)^2}$$

which is of the form $g(y/x)$ for a function of the form

$$g(z) = z + \sqrt{1 + z^2}.$$

If we had considered $x < 0$, we would have obtained

$$g(z) = z - \sqrt{1 + z^2}.$$

In either case, we see that the differential equation is homogeneous. ■

As we mentioned, we have introduced homogeneous differential equations because they are related to separable equations; in fact, we have the following theorem which formalizes the method used in Example 4.

THEOREM 1.6.2

If

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{1.45}$$

is a homogeneous equation, then the change of variables

$$y = vx$$

transforms (1.45) into a separable equation in the variables v and x .

Note that this change of variables implies that

$$y' = v + xv'$$

by the product rule.

Example 7 Solve

$$y + (x - 2y) \frac{dy}{dx} = 0.$$

Solution

We first observe that this can be rewritten as

$$\frac{dy}{dx} = \frac{y}{2y - x}.$$

Dividing numerator and denominator by x gives

$$\frac{dy}{dx} = \frac{y/x}{2y/x - 1}.$$

The right-hand side is then of the form $g(y/x)$, and making the change of variables $y = vx$ gives

$$v + x \frac{dv}{dx} = \frac{v}{2v - 1},$$

which becomes

$$x \frac{dv}{dx} = \frac{2(v - v^2)}{2v - 1}.$$

This equation is separable! Rearranging gives

$$\frac{2v - 1}{2(v - v^2)} dv = \frac{1}{x} dx,$$

and integrating both sides yields

$$-\frac{1}{2} \ln |v - v^2| = \ln |x| + C_1.$$

We then use $v = y/x$ to reintroduce the y -variable. Thus

$$-\frac{1}{2} \ln \left| \frac{y}{x} - \left(\frac{y}{x} \right)^2 \right| = \ln |x| + C_1,$$

but we can let $C_1 = \ln C$ for an arbitrary constant C , so that

$$-\frac{1}{2} \ln \left| \frac{y}{x} - \left(\frac{y}{x} \right)^2 \right| = \ln |Cx|$$

is the implicit solution. We could obtain an explicit solution with a bit more work but choose not to. Again, we could plot these solutions for various C -values with our favorite software package.



Problems

Solve Problems 1–8 by first finding an integrating factor of suitable form.

- | | |
|------------------------------------|--|
| 1. $ydx + (e^x - 1)dy = 0$ | 2. $(x^2 + y^2 + x)dx + ydy = 0$ |
| 3. $y(x + y)dx + (xy + 1)dy = 0$ | 4. $(x^2 - y^2 + y)dx + x(2y - 1)dy = 0$ |
| 5. $ydx - xdy = 2x^3 \sin x \, dx$ | 6. $(3x^2 + y)dx + (x^2y - x)dy = 0$ |
| 7. $(3x^2y - x^2)dx + dy = 0$ | 8. $(x^2 + 2x + y)dx = (x - 3x^2y)dy$ |

9. Show that, if $(\partial N / \partial x - \partial M / \partial y) / (xM - yN)$ depends only on the product xy , that is,

$$\frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} = H(xy),$$

then the equation

$$M(x, y) dx + N(x, y) dy = 0$$

has an integrating factor of the form $\mu(xy)$. Find the general formula for $\mu(xy)$.

10. We derived a formula for an integrating factor if $\mu(x, y) = \mu(x)$. If $\mu(x, y) = \mu(y)$, derive the integrating factor formula

$$\mu(y) = \exp \left[\int \frac{1}{M(x, y)} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dy \right]. \quad (1.46)$$

Solve the Bernoulli equations given in Problems **11–21**.

- | | |
|--------------------------------------|---|
| 11. $y' + y = xy^2$ | 12. $y' + 3y = y^4$ |
| 13. $y' + 2xy = 4y$ | 14. $y' - xy = xy^3$ |
| 15. $xydy = (y^2 + x)dx$ | 16. $xy' + 2y + x^5y^3e^x = 0$ |
| 17. $xy' - 2x^2\sqrt{y} = 4y$ | 18. $y' = y^4 \cos x + y \tan x$ |
| 19. $xy^2y' = x^2 + y^3$ | 20. $(x + 1)(y' + y^2) = -y$ |
- 21.** Solve the logistic equation $\frac{dN}{dt} = rN \left(1 - \frac{N}{K}\right)$.

For Problems **22–34**, solve the homogeneous differential equation analytically.

- | | |
|---|--|
| 22. $(x + y)dx - xdy = 0$ | 23. $(x + 2y)dx - xdy = 0$ |
| 24. $(y^2 - 2xy)dx + x^2dy = 0$ | 25. $2x^3y' = y(2x^2 - y^2)$ |
| 26. $2x^2\frac{dy}{dx} = x^2 + y^2$ | 27. $\frac{dy}{dx} = \frac{y^2 + 2xy}{x^2}$ |
| 28. $xy' - y = x \tan\left(\frac{y}{x}\right)$ | 29. $(x^2 + y^2)y' = 2xy$ |
| 30. $ydx = (2x + y)dy$ | 31. $(x - y)dx + (x + y)dy = 0$ |
| 32. $y' = 2\left(\frac{y}{x+y}\right)^2$ | 33. $y^2 + x^2y' = xy y'$ |
- 34.** $(x + 4y)y' = 2x + 3y$

35. A function F is called **homogeneous of degree n** if

$$F(tx, ty) = t^n F(x, y) \text{ for all } x \text{ and } y.$$

That is, if tx and ty are substituted for x and y in $F(x, y)$ and if t^n is then factored out, we are left with $F(x, y)$. For instance, if $F(x, y) = x^2 + y^2$, we note that

$$F(tx, ty) = (tx)^2 + (ty)^2 = t^2 F(x, y)$$

so that F is homogeneous of degree 2. Homogeneous differential equations and functions that are homogeneous of degree n are related in the following manner. Suppose the functions M and N in the differential equation

$$M(x, y)dx + N(x, y)dy = 0$$

are both homogeneous of the same degree n .

(a) Show, using the change of variables $t = 1/x$, that

$$M\left(1, \frac{y}{x}\right) = \left(\frac{1}{x}\right)^n M(x, y),$$

which implies that

$$M(x, y) = \left(\frac{1}{x}\right)^{-n} M\left(1, \frac{y}{x}\right).$$

(b) Show, using a similar calculation, that

$$N(x, y) = \left(\frac{1}{x}\right)^{-n} N\left(1, \frac{y}{x}\right),$$

so that the differential equation

$$M(x, y)dx + N(x, y)dy = 0$$

becomes

$$\frac{dy}{dx} = \frac{-M(x, y)}{N(x, y)} = -\frac{\left(\frac{1}{x}\right)^{-n} M\left(1, \frac{y}{x}\right)}{\left(\frac{1}{x}\right)^{-n} N\left(1, \frac{y}{x}\right)}.$$

Simplifying gives

$$\frac{dy}{dx} = -\frac{M(1, \frac{y}{x})}{N(1, \frac{y}{x})}.$$

(c) Show that both numerator and denominator of the right-hand side of

$$\frac{dy}{dx} = -\frac{M(1, \frac{y}{x})}{N(1, \frac{y}{x})}$$

are in the form $g(y/x)$ and conclude that, if M and N are both homogeneous functions of the same degree n , then the differential equation

$$M(x, y) dx + N(x, y) dy = 0$$

is a homogeneous differential equation.

- 36.** Using the idea presented in Problem **35**, show that each of the equations in Problems **22–34** are homogeneous.
- 37.** Suppose that the equation $M(x, y) dx + N(x, y) dy = 0$ is homogeneous. Show that the transformation $x = r \cos t$, $y = r \sin t$ reduces this equation to a separable equation in the variables r and t .

Use the method of Problem **37** to solve Problems **38** and **39**.

38. $(x - y)dx + (x + y)dy = 0$

39. $(x + y) dx - x dy = 0$

- 40.** (a) Solve

$$\frac{dy}{dx} = \frac{y - x}{y + x}.$$

(b) Now consider

$$\frac{dy}{dx} = \frac{y - x + 1}{y + x + 5}. \quad (1.47)$$

(i) Show that this equation is NOT homogeneous.

How can we solve this? Consider the equations $y - x = 0$ and $y + x = 0$. They represent two straight lines through the origin. The intersection of $y - x + 1 = 0$ and $y + x + 5 = 0$ is $(-2, -3)$. Check it! Let $x = X - 2$ and $y = Y - 3$. This amounts to taking new axes parallel to the old with an origin at $(-2, -3)$.

(ii) Use this transformation to obtain the differential equation

$$\frac{dY}{dX} = \frac{Y - X}{Y + X}.$$

(iii) Using the solution from part (a), obtain the solution to (1.47).

Use the technique of Problem **40** to solve Problems **41–45**.

41. $(2x + y + 1)dx - (4x + 2y - 3)dy = 0$

42. $x - y - 1 + (y - x + 2)y' = 0$

43. $(x + 4y)y' = 2x + 3y - 5$

44. $(y + 2)dx = (2x + y - 4)dy$

45. $y' = 2\left(\frac{y+2}{x+y-1}\right)^2$

Chapter 1 Review

In Problems 1–7, determine whether the statement is true or false. If it is true, give reasons for your answer. If it is false, give a counterexample or other explanation of why it is false.

1. The equation $y'' + xy' - y = x^2$ is a linear ordinary differential equation that is considered an initial-value problem.
2. The equation $y^{(4)} - y^2 = \sin x$ is a nonlinear ordinary differential equation.
3. An implicit solution to $y' = f(x, y)$ can be written in the form $y(x) = f(x)$.
4. With an appropriate substitution, any exact equation can be put in the form of a linear equation.
5. A solution to the differential equation $y' = f(x, y)$ must be defined for all x .
6. An equation of the form $M(x, y) + N(x, y)y' = 0$ is considered exact if

$$\frac{\partial M(x, y)}{\partial y} = \frac{\partial N(x, y)}{\partial x}.$$

7. Every equation of the form $y' = f(y)$ is separable.

In Problems 8–11, verify that the given function is a solution (possibly implicit) to the given differential equation. State the interval on which it is a solution and the C -values for which the solution is valid.

- | | |
|---|---|
| 8. $y(x) = 1 + Ce^{x^4}$, $y' = 4x^3(y - 1)$ | 9. $y(x) = Ce^x - x^2 - 2x - 2$, $y' = x^2 + y$ |
| 10. $y^2(x) = \frac{1}{C - 2x}$, $y' = y^3$ | 11. $y(x) = \frac{3x - 1 + 3C}{x + C}$, $y' = (y - 3)^2$ |

In Problems 12–16, solve each of the following separable differential equations.

- | | |
|--------------------------------------|----------------------------------|
| 12. $y' - xy^2 = 2xy$ | 13. $2x^2yy' + y^2 = 2$ |
| 14. $e^{-x}(1 + \frac{dx}{dt}) = 1$ | 15. $xy' + y = y^2$ $y(1) = 0.5$ |
| 16. $y' = 3\sqrt[3]{y^2}$ $y(2) = 0$ | |

In Problems 17–22, solve each of the following homogeneous differential equations.

- | | |
|--------------------------------------|---|
| 17. $xy' = y - xe^{y/x}$ | 18. $xy' - y = (x + y) \ln \frac{x+y}{x}$ |
| 19. $xy' = y \cos(\ln(\frac{y}{x}))$ | 20. $(y + \sqrt{xy})dx = xdy$ |
| 21. $xy' = \sqrt{x^2 - y^2} + y$ | 22. $(2x - 4y)dx + (x + y)dy = 0$ |

In Problems 23–25, solve each of the following exact differential equations.

23. $\frac{y}{x} + (y^3 + \ln x) \frac{dy}{dx} = 0$
24. $2x(1 + \sqrt{x^2 - y}) - \sqrt{x^2 - y} \frac{dy}{dx} = 0$
25. $\frac{3x^2 + y^2}{y^2} - \frac{2x^3 + 5y}{y^3} \frac{dy}{dx} = 0$

In Problems 26–28, solve each of the following differential equations by first finding an integrating factor of a suitable form.

- | | |
|--|--------------------------------|
| 26. $xy^2(xy' + y) = 1$ | 27. $y^2dx - (xy + x^3)dy = 0$ |
| 28. $(y - \frac{1}{x})dx + \frac{dy}{y} = 0$ | |

In Problems **29–34**, solve the following linear or Bernoulli equations.

29. $xy' + y = x^4$

30. $y' - y = \sin x$

31. $y' + 4xy = 5x^3$

32. $xy' + y = xy^4$

33. $y' - y = xy^3$

34. $2y' + xy = 4xy^3$

Solve the remaining problems using one of the methods of this chapter.

- 35.** *One Big Ol' Pot of Soup* Mike, a professor of mathematics and a part-time evening cook at a local diner, prepares a big pot of soup late at night, just before closing time. He does this so that there will be plenty of soup to feed customers the next day. Being food safety cautious, he knows that refrigeration is essential to preserve the soup overnight; however, the soup is too hot to be put directly into the fridge when it is ready. (The soup had just boiled at 100 °C, and the fridge is not powerful enough to accommodate a big pot of soup if it is any warmer than 20 °C.) Mike is resourceful (as he is a student of M.M. Rao) and discovered that by cooling the pot in a sink full of cold water (kept running, so that its temperature was roughly constant at 5 °C) and stirring occasionally, he could bring the temperature of the soup to 60 °C in 10 min. How long before closing time should the soup be ready so that Mike can put it in the fridge?
- 36.** A cup of very hot coffee from a fast food restaurant is at 120 °F when it is served. If left on the table in a room with ambient temperature of 70 °F, it is found to have cooled to 115 °F in 1 min. If burns can result from spilling coffee at a temperature greater than 100 °F, how many minutes will the coffee have to sit before the management is safe from lawsuits?
- 37.** A 30-gal tank initially has 15 gal of saltwater containing 6 lb of salt. Saltwater containing 1 lb of salt per gallon is pumped into the top of the tank at the rate of 2 gal/min, while a well-mixed solution leaves the bottom of the tank at a rate of 1 gal/min. Determine the amount of salt in the tank at time t . How long does it take for the tank to fill? What is the amount of salt in the tank when it is full?
- 38.** A 100-gal tank initially contains 100 gal of sugar-water at a concentration of 0.25 lb of sugar per gallon. Suppose sugar is added to the tank at a rate of p lb/min, sugar-water is removed at a rate of 1 gal/min, and the water in the tank is kept well mixed. Determine the concentration of sugar at time t . What value of p should be chosen so that, when 5 gal of sugar solution is left in the tank, the concentration is 0.5 lb of sugar per gallon?
- 39.** Find a curve such that the point of intersection of an arbitrary tangent with the x -axis has an abscissa half as great as the abscissa of the point of tangency.

Chapter 1 Computer Labs

The reader should only complete the computer lab(s) after reading Appendix A, where the introductory details and some basic commands and examples are given. Although every attempt has been made to make the syntax given in this book applicable to a range of versions/releases, certain syntax may be different for an earlier or later version of a given

package. We assume the reader has *no* familiarity with any package but has access to at least one of them. From this point, we will assume the reader is familiar with the relevant section of Appendix A.

Chapter 1 Computer Lab: MATLAB

MATLAB Example 1: Enter the following code, which demonstrates how to differentiate and integrate functions using the *Symbolic Math Toolbox*. It considers $f(x) = \cos x + e^{2x} + \ln x$ and calculates $f'(x)$, $f''(x)$, $\frac{df(x)}{dt}$, $\int f(x)dx$, $\int_0^\pi f(x)dx$, and $\int_0^x f(t)dt$.

```
>> clear all
>> syms x t
>> f(x)=cos(x)+exp(2*x)+log(x)
>> diff(f(x),x)
>> diff(f(x),x,2)
>> diff(f(x),t)
>> int(f(x),x)
>> int(f(x),x,0,pi)
>> quad('cos(x)+exp(2*x)+log(x)',0,pi) %definite int w/o syms
>> int(f(x),t)
>> f(t)=subs(f(x),x,t)
>> g(x)=int(f(t),0,x)
>> g(pi)
```

MATLAB Example 2: Enter the following code, which demonstrates how to verify that a given function is a solution (possibly implicit) to the differential equation. The *Symbolic Math Toolbox* is again needed. It considers $y'' + 2y' + y = 0$ with explicit solutions $y_1 = e^{-x}$ and $y_2 = xe^{-x}$, followed by $(x-4)y^4 - x^3(y^2-3)y' = 0$ with implicit solution $\frac{-1}{x} + \frac{2}{x^2} = \frac{-1}{y} + \frac{1}{y^3} + C$.

```
>> clear all
>> syms x y(x) C
>> eq1=diff(y(x),x,2)+2*diff(y(x),x)+y(x)
>> subs(eq1,y(x),exp(-x))
>> subs(eq1,y(x),x*exp(-x))
>> subs(eq1,y(x),exp(x))
>> eq2=(x-4)*y(x)^4-x^3*(y(x)^2-3)*diff(y(x),x)
>> sol2=-1/x+2/x^2+1/y(x)-1/y(x)^3-C
>> dsol2=diff(sol2,x)
>> collect(expand(y(x)^4*x^3*dsol2),diff(y(x),x))
```

MATLAB Example 3: Enter the following code, which demonstrates how to plot one implicit function (in this case a solution to a differential equation) and then multiple plots with the same function but different constants. It considers the implicit solution from MATLAB Example 2, $\frac{-1}{x} + \frac{2}{x^2} = \frac{-1}{y} + \frac{1}{y^3} + C$.

```
>> clear all
```

```
>> [X,Y]=meshgrid(-2:.01:2,-2:.01:2);
>> Z=-1./X+2./X.^2+1./Y-1./Y.^3;
>> contour(X,Y,Z,[-5 -5])
>> [C,h]=contour(X,Y,Z,[-5 -5]);
>> clabel(C,h)
>> axis([-2 2 0 0.6])
>> xlabel('x'), ylabel('y')
>> title('Implicit Plot with C=-5')
>> [C,h]=contour(X,Y,Z,[-10,2,6,50]);
>> clabel(C,h)
>> xlabel('x'), ylabel('y')
>> title('Implicit Plot with C=-10,2,6,50')
```

MATLAB Example 4: We will now numerically solve

$$(x^2 + 1)y' + 4xy = x, \quad y(0) = 10.$$

In the code below, you will first create a MATLAB function (separate file) called Example4.m. Then you will use MATLAB's ode45 to plot the approximate numerical solution.

```
%%% Create a function, Example4.m, and save it.
function dydx=Example4(xn,yn)
%
%Rewrite the original ode in the form y'=f(x,y)
%to obtain y'=(x-4xy)/(x^2+1)
%
dydx=(xn-4*xn*yn)/(xn^2+1);

% Now type the following in the command window.
>> clear all
>> [x,y]=ode45(@Example4,[0,4],10);
>> plot(x,y)
>> subplot(2,1,1),plot(x,y)
>> title('Numerical Approximation of Soln of (x^2+1)y{\prime}
+4xy=x, y(0)=10')
>> xlabel('x'), ylabel('y')
>> x1=0:.1:4;
>> y1=((1/4)*x1.^4+(1/2)*x1.^2+10)./(x1.^2+1).^2;
>> subplot(2,1,2),plot(x1,y1,'k')
>> title('Closed Form Soln of (x^2+1)y{\prime}+4xy=x,y(0)=10')
>> xlabel('x'), ylabel('y')
```

MATLAB Exercises

Turn in both the commands that you enter for the exercises below as well as the output/figures. These should all be in one document. Please highlight or clearly indicate all requested answers. Some of the questions will require you to modify the above MATLAB code to answer them.

1. Enter the commands given in MATLAB Example 1 and submit both your input and output.

2. Enter the commands given in MATLAB Example 2 and submit both your input and output.
3. Enter the commands given in MATLAB Example 3 and submit both your input and output.
4. Enter the commands given in MATLAB Example 4 and submit both your input and output.
5. Find the derivative of $t^2 + \cos t$ with respect to t .
6. Find the derivative of $x^2 + \cos x$ with respect to x .
7. Find the second derivative of $t^2 + \cos t$ with respect to t .
8. Find the second derivative of $x^2 + \cos x$ with respect to x .
9. Find the antiderivative of $t^2 + \cos t$ with respect to t .
10. Find the antiderivative of $x^2 + \cos x$ with respect to x .
11. Find the derivative of $\ln(tx) + \cos^2 x + \sin t^2$ with respect to t .
12. Find the derivative of $e^{tx} + \cos^2 x + \sin t^2$ with respect to x .
13. Find the antiderivative of $\ln(tx) + \cos^2 x + \sin t^2$ with respect to t .
14. Find the antiderivative of $e^{tx} + \cos^2 x + \sin t^2$ with respect to x .
15. Find the definite integral of $x^2 + \cos x$ from $x = 0$ to $x = 3$.
16. Find the definite integral of $\ln x$ from $x = 1$ to $x = 5$.
17. Find the definite integral of e^{-x^2} from $x = -1$ to $x = 3$. Give the answer as a decimal. You may find it easier to use **quad** rather than the *Symbolic Math Toolbox*.
18. Find the definite integral of $\frac{\sin x}{x}$ from $x = 1$ to $x = 10$. Give the answer as a decimal. You may find it easier to use **quad** rather than the *Symbolic Math Toolbox*.
19. Numerically verify that $y(x) = \sin x + 2 \cos x$ is a solution on $(-\infty, \infty)$ to $\frac{d^2 y}{dx^2} + y = 0$.
20. Numerically verify that $y_1(x) = e^x, y_2(x) = xe^x$ are both solutions on $(-\infty, \infty)$ to $y'' - 2y' + y = 0$.
21. Solve $y' = \sin(x^2), y(0) = 0$ by hand. Then use MATLAB to find $y(1), y(3)$. (Hint: remember, from calculus, that $y(x) = \int_0^x f(t)dt$ is a function of x and thus $y(1)$ is a number that can be evaluated with the above code.) Use MATLAB to plot the solution from $x = 0$ to $x = 4$.
22. Use MATLAB to numerically solve $y' = \sin(x^2), y(0) = 0$. Plot this numerical solution from $x = 0$ to $x = 4$.
23. Solve $y' = \frac{\sin(x)}{x}, y(1) = 0$ by hand. Then use MATLAB to find $y(2), y(3)$. (Hint: remember, from calculus, that $y(x) = \int_1^x f(t)dt$ is a function of x and thus $y(2)$ is a number that can be evaluated with the above code.) Use MATLAB to plot the solution from $x = 1$ to $x = 6$.
24. Use MATLAB to numerically solve $y' = \frac{\sin(x)}{x}, y(1) = 0$. Plot this numerical solution from $x = 1$ to $x = 6$.
25. Numerically verify that $y - y^6 + \cos x + x \sin x = C$ is an implicit solution to the ODE $(1 - 6y^5)y' = -x \cos x$.
26. Solve $y + xy' = \frac{y}{2y - 1}$ by hand. This solution is implicit. Choose 3 different C -values and plot the implicit solutions.