


Issues in Toxicology

Edited by Alok Dhawan and Diana Anderson

The Comet Assay in Toxicology



RSC Publishing

The Comet Assay in Toxicology

Issues in Toxicology

Series Editors:

Professor Diana Anderson, *University of Bradford, UK*

Dr Michael D Waters, *Integrated Laboratory Systems, Inc, N Carolina, USA*

Dr Timothy C Marrs, *Edentox Associates, Kent, UK*

Titles in the Series:

- 1: Hair in Toxicology: An Important Bio-Monitor
- 2: Male-mediated Developmental Toxicity
- 3: Cytochrome P450: Role in the Metabolism and Toxicity of Drugs and other Xenobiotics
- 4: Bile Acids: Toxicology and Bioactivity
- 5: The Comet Assay in Toxicology

How to obtain future titles on publication:

A standing order plan is available for this series. A standing order will bring delivery of each new volume immediately on publication.

For further information please contact:

Sales and Customer Care, Royal Society of Chemistry, Thomas Graham House, Science Park, Milton Road, Cambridge, CB4 0WF, UK

Telephone: +44 (0)1223 432360, Fax: +44 (0)1223 420247, Email: sales@rsc.org

Visit our website at <http://www.rsc.org/Shop/Books>

The Comet Assay in Toxicology

Edited by

Alok Dhawan

*Developmental Toxicology Division, Indian Institute of Toxicology Research,
Lucknow, India*

Diana Anderson

Division of Biomedical Sciences, University of Bradford, Bradford, UK

RSC Publishing

The cover image shows photomicrographs of comets from (A) *Escherichia coli* (B) *Bacopa monerii* L. (C) *Drosophila melanogaster* (D) differential DNA damage lymphocytes (E) human lymphocytes (F) irradiated diploid human lymphocyte with FISH showing double hybridisation signals indicating strand breakage (G) comets in human sperm (H) human sperm showing double breaks (I) haploid human sperm with FISH showing single hybridisation signal.

Issues in Toxicology No 5

ISBN: 978-0-85404-199-2

ISSN: 1757-7179

A catalogue record for this book is available from the British Library

© The Royal Society of Chemistry, 2009

All rights reserved

Apart from fair dealing for the purposes of research for non-commercial purposes or for private study, criticism or review, as permitted under the Copyright, Designs and Patents Act 1988 and the Copyright and Related Rights Regulations 2003, this publication may not be reproduced, stored or transmitted, in any form or by any means, without the prior permission in writing of The Royal Society of Chemistry or the copyright owner, or in the case of reproduction in accordance with the terms of licences issued by the Copyright Licensing Agency in the UK, or in accordance with the terms of the licences issued by the appropriate Reproduction Rights Organization outside the UK. Enquiries concerning reproduction outside the terms stated here should be sent to The Royal Society of Chemistry at the address printed on this page.

Published by The Royal Society of Chemistry,
Thomas Graham House, Science Park, Milton Road,
Cambridge CB4 0WF, UK

Registered Charity Number 207890

For further information see our web site at www.rsc.org

Preface

This book is the first of its kind to be devoted exclusively to the Comet assay and its applications as an important tool in current toxicology. This multi-author book will serve as both a reference and a guide for investigators in the biomedical, biochemical and pharmaceutical sciences. Specialists from the fields of genetic toxicology and human epidemiology, with first-hand knowledge of their chosen subspecialties, have contributed to this peer-reviewed scientific venture.

Simplicity, rapidity, versatility and ease of application of the Comet assay have made it a favourite amongst researchers and it is now also gaining acceptance amongst regulators. It can be used in all single cells from prokaryotes and eukaryotes, in plants and animals including humans, involving both somatic and germ cells. It is also a relatively inexpensive assay to perform.

The book is divided into different sections, reflecting the range of interest in the exploitation of this assay. It begins with an introductory section reviewing the genesis of the assay for those new to the technique, and details the various fields in which it finds wide acceptance. This sets the scene by explaining why the assay has become the most sensitive and sought after assay in modern toxicology.

There is a section that describes the protocols being followed to assess various types of DNA damage in different cell types. The third section brings together the specific applications of the assay in diverse areas ranging from genetic toxicity testing to human monitoring, and environmental toxicology. The last section considers strategies for the conduct of the assay using *in vitro* and *in vivo* systems, based on internationally accepted guidelines. The book draws to a close with an assessment of image-analysis principles and the statistics used for evaluating the data generated by the assay.

This book is a culmination of over fifteen years of active collaboration and friendship between the editors and provides a good basic understanding of issues relating to the assay.

The Editors

Issues in Toxicology No 5

The Comet Assay in Toxicology

Edited by Alok Dhawan and Diana Anderson

© Royal Society of Chemistry 2009

Published by the Royal Society of Chemistry, www.rsc.org

Contents

Section I: Genesis of Comet Assay

Chapter 1	The Comet Assay: A Versatile Tool for Assessing DNA Damage	3
	<i>Alok Dhawan, Mahima Bajpayee and Devendra Parmar</i>	
1.1	Introduction	3
1.2	Bacteria	5
1.3	Plant Models	5
1.3.1	The Comet Assay in Lower Plants	5
1.3.2	The Comet Assay in Higher Plants	18
1.4	Animal Models	19
1.4.1	Lower Animals	19
1.5	Higher Animals	24
1.5.1	Vertebrates	24
1.6	The Specificity, Sensitivity and Limitations of the Comet Assay	28
1.7	Conclusions	30
	Acknowledgements	30
	References	30

Section II: Various Procedures for the Comet Assay

Chapter 2	Detection of Oxidised DNA Using DNA Repair Enzymes	57
	<i>Amaya Azqueta, Sergey Shaposhnikov and Andrew R. Collins</i>	
2.1	Introduction	57
2.2	Methods for Measuring DNA Oxidation Damage	60

2.3	Enzyme Specificity	61
2.4	Applications	62
2.5	Protocol	64
2.5.1	Equipment	64
2.5.2	Supplies	65
2.5.3	Reagents, Buffers and Enzymes	65
2.5.4	Procedure	66
	Acknowledgment	74
	References	75
Chapter 3	Microplate-Based Comet Assay	79
	<i>Elizabeth D. Wagner and Michael J. Plewa</i>	
3.1	Introduction	79
3.2	Microplate Comet Assay	80
3.3	Drinking-Water Disinfection Byproducts	80
3.4	Chinese Hamster Ovary Cells	81
3.5	CHO Cell Microplate Comet Assay Protocol	81
3.5.1	CHO Cell Treatment	81
3.5.2	Preparation of Comet Microgels	83
3.5.3	Comet Microscopic Examination	84
3.5.4	Normalisation of CHO Cell Comet Data and Statistical Analysis	85
3.6	Utility of the Microplate Comet Assay in Comparing Classes of DBPs	87
3.6.1	Microplate Comet Analysis of the Haloacetonitriles	89
3.6.2	Microplate Comet Analysis of the Haloacetamides	90
3.6.3	Comparison of SCGE Genotoxic Potency Values of the Haloacetonitriles and Haloacetamides	91
3.7	Advantages of the Mammalian Cell Microplate Comet Assay	92
	Acknowledgements	92
	References	93
Chapter 4	The Use of Higher Plants in the Comet Assay	98
	<i>Tomas Gichner, Irena Znidar, Elizabeth D. Wagner and Michael J. Plewa</i>	
4.1	Introduction	98
4.2	Differences between the Animal and Plant Comet Assay	99

4.3	Cultivation and Treatment of Plants for the Comet Assay	99
4.3.1	Onion (<i>Allium cepa</i>)	99
4.3.2	Tobacco (<i>Nicotiana tabacum</i>)	100
4.3.3	Broad Bean (<i>Vicia faba</i>)	100
4.3.4	Plants used for <i>In-Situ</i> Studies	100
4.4	Isolation of Nuclei from Plant Tissues	101
4.4.1	Isolation of Nuclei <i>via</i> Protoplast Formation	101
4.4.2	Isolation of Nuclei by Mechanical Destruction of the Cell Wall	101
4.5	Preparation of Comet Assay Slides	101
4.6	DNA Unwinding and Electrophoresis	102
4.7	DNA Staining	103
4.8	Reading the Slides, Expressing DNA Damage, Statistics	103
4.9	Comet Assay Procedure	104
4.10	Reagents, Media, Buffers	105
4.11	Equipment and Software	107
4.12	Determination of Toxicity	107
4.13	Correlation between the DNA Damage Evaluated by the Comet Assay and Other Genetic Endpoints in Plants	108
4.14	The Utility of the Comet Assay for Genotoxic Studies in the Laboratory	109
4.15	The Utility of the Comet Assay as an <i>In Situ</i> Marker	109
4.16	Comet Assay with Irradiated Food of Plant Origin	110
4.17	Recommendations for Plant Comet Assay Users	110
	Abbreviations	114
	References	115

Chapter 5 Methods for Freezing Blood Samples at -80°C for DNA Damage Analysis in Human Leukocytes **120**

Narendra P. Singh and Henry C. Lai

5.1	Introduction	120
5.2	Materials and Methods	121
5.2.1	Protocol I	121
5.2.2	Protocol II	122
5.2.3	Fresh Blood	122
5.2.4	Fresh Blood Stored on Ice Prior to Freezing	122
5.2.5	Image and Data Analysis	123
5.3	Results and Discussion	123
	References	127

Chapter 6	Development and Applications of the Comet-FISH Assay for the Study of DNA Damage and Repair	129
	<i>Valerie J. McKelvey-Martin and Declan J. McKenna</i>	
6.1	Introduction	129
6.2	The Comet-FISH Assay Procedure	130
6.3	Applications of the Comet-FISH Assay	135
6.3.1	Discovery of the Comet-FISH Assay	135
6.3.2	Using Comet-FISH to Measure DNA Damage	135
6.3.3	Using Comet-FISH to Quantify DNA Repair	142
6.3.4	Summary of Studies	143
6.4	Limitations of Comet-FISH Assay	144
6.4.1	Practical Difficulties	144
6.4.2	Imaging Difficulties	144
6.4.3	Interpretation of Results	145
6.5	Conclusion	146
	References	146
Chapter 7	Detection of DNA Damage in <i>Drosophila</i> and Mouse	151
	<i>Alok Dhawan, Mahima Bajpayee and Devendra Parmar</i>	
7.1	General Protocol for the Assessment of DNA Damage Using the Alkaline Comet Assay	153
7.1.1	Chemicals and Materials	153
7.1.2	Preparation of Reagents	153
7.1.3	Preparation of Agarose-Coated (Base) Slides for the Comet Assay	155
7.1.4	Preparation of Microgel Slides for the Comet Assay	155
7.1.5	Electrophoresis of Microgel Slides	156
7.1.6	Evaluation of DNA Damage	157
7.2	The Alkaline Comet Assay in Multiple Organs of Mouse	157
7.2.1	Chemicals and Materials	158
7.2.2	Methodology	158
7.3	The Alkaline Comet Assay in <i>Drosophila melanogaster</i>	161
7.3.1	Chemicals and Materials	162
7.3.2	Methodology	162
7.4	Conclusions	165
	Acknowledgements	166
	References	166

Section III: Applications of Comet Assay

Chapter 8	Clinical Applications of the Comet Assay	173
	<i>S. M. Piperakis, K. Kontogianni, G. Karanastasi and M. M. Piperakis</i>	
8.1	Introduction	173
8.2	The Comet Assay Methodology	174
8.3	Clinical Studies	175
8.4	Discussion and Conclusions	195
	References	196
Chapter 9	Applications of the Comet Assay in Human Biomonitoring	201
	<i>Andrew R. Collins and Maria Dusinska</i>	
9.1	Biomonitoring and Biomarkers – An Introduction	201
9.2	The (Modified) Comet Assay	202
9.3	Guidelines for Biomonitoring Studies	202
9.4	Biomonitoring with the Comet Assay: Special Considerations	204
9.4.1	Surrogate and Target Cells; The Use of White Blood Cells	204
9.4.2	Sampling Time and Transport	205
9.4.3	Reference Standards	206
9.4.4	What Affects the Background Level of DNA Damage?	206
9.5	DNA Damage as a Marker of Environmental Exposure and Risk	207
9.6	DNA Repair as a Biomarker of Individual Susceptibility	207
9.7	Protocols	208
9.7.1	Protocol for Blood Sample Collection and Long-Term Storage of Lymphocytes for the Measurement of DNA Damage and Repair	208
9.7.2	Comet Assay – Determination of DNA Damage (Strand Breaks and Oxidised Bases)	211
9.7.3	<i>In Vitro</i> Assays for DNA Repair	214
9.8	Solutions, etc.	216
9.8.1.	Lysis Solution	216
9.8.2.	Buffer F (Enzyme Reaction Buffer for FPG, Endonuclease III, and <i>In Vitro</i> BER Assay)	217
9.8.3	Buffer F + Mg (Used for <i>In Vitro</i> NER Assay)	217
9.8.4	Buffer A (Used in <i>In Vitro</i> Repair Assays)	217
9.8.5	Triton Solution	217
9.8.6	Ro 19-8022 (Photosensitiser)	217

9.8.7	Electrophoresis Solution	217
9.8.8	Neutralising Buffer	218
9.8.9	Agarose	218
9.8.10	Enzymes	218
9.9	Analysis and Interpretation of Results	218
9.9.1	Quantitation	218
9.9.2	Calculation of Net Enzyme-Sensitive Sites	219
9.9.3	Calibration	219
9.9.4	How to Deal with Comet Assay Data Statistically	219
9.10	Conclusions	220
	Acknowledgements	221
	References	221
Chapter 10	The Comet Assay in Human Biomonitoring	227
	<i>Mahara Valverde and Emilio Rojas</i>	
10.1	Introduction	227
10.2	Human Monitoring	228
10.3	Environmental Exposure	230
10.4	Lifestyle Exposure	234
10.5	Occupational Exposure	237
10.6	Reviews	248
10.7	Usefulness of the Comet Assay in Human Monitoring	249
10.8	Conclusions	251
	References	252
Chapter 11	Comet Assays in Dietary Intervention Trials	267
	<i>Armen Nersesyan, Christine Hoelzl, Franziska Ferk, Miroslav Mišik and Siegfried Knasmueller</i>	
11.1	Introduction	267
11.2	Experimental Design of Human Studies	268
11.3	Indicator Cells and Media	269
11.4	Conventional SCGE Trials with Complex Foods and Individual Components – The Current State of Knowledge	270
11.5	Use of SCGE Trials to Detect Protection Against DNA-Reactive Carcinogens	275
11.6	Use of SCGE Experiments to Monitor Alterations of the DNA-Repair Capacity	279
11.7	What Have We Learned from Intervention Studies so Far?	281

<i>Contents</i>	xiii
11.8 Future Perspectives	282
References	284
Chapter 12 The Comet Assay for the Evaluation of Genotoxic Exposure in Aquatic Species	297
<i>G. Frenzilli and B. P. Lyons</i>	
12.1 Introduction	297
12.2 Protocols, Cell Types and Target Organs	298
12.3 Application of the Comet Assay to Invertebrate Species	299
12.3.1 Freshwater Invertebrates	299
12.3.2 Marine Invertebrates	300
12.4 Application of the Comet Assay to Vertebrate Species	301
12.4.1 Freshwater Vertebrates	301
12.4.2 Marine Vertebrates	302
12.5 Conclusions	303
References	303
Chapter 13 The Alkaline Comet Assay in Prognostic Tests for Male Infertility and Assisted Reproductive Technology Outcomes	310
<i>Sheena E. M. Lewis and Ishola M. Agbaje</i>	
13.1 Introduction	310
13.2 Sites of DNA Damage in Sperm	311
13.2.1 Oxidative Stress, a Major Cause of DNA Damage	312
13.2.2 Oxidative Stress, Antioxidant Therapies	312
13.2.3 Sperm DNA Damage Tests	313
13.2.4 Modifications to the Alkaline Comet Assay for Use with Sperm	314
13.2.5 Sperm DNA Adducts and their Relationship with DNA Fragmentation	316
13.3 Can Sperm DNA Integrity Predict Success? Relationships with Assisted Conception Outcomes	317
13.4 Clinically Induced DNA Damage	318
13.4.1 Cryopreservation	319
13.4.2 Vasectomy	319
13.5 A Major Barrier to Progress	320
13.6 Opportunities and Challenges – The Establishment of Clinical Thresholds and the Integration of DNA Testing into Clinical Practice	320

Acknowledgements	321
References	321

Chapter 14 The Comet Assay in Sperm – Assessing Genotoxins in Male Germ Cells 331

Adolf Baumgartner, Eduardo Cemeli, Julian Laubenthal and Diana Anderson

14.1	Introduction	331
14.2	Single-Cell Gel Electrophoresis	332
14.3	The Use of Sperm with the Comet Assay	333
14.3.1	Human Sperm	333
14.3.2	Modifying Existing Comet Protocols for the Use of Sperm	333
14.3.3	Sperm DNA and the Comet Assay	334
14.3.4	The Sperm Comet Assay and the Use of Repair Enzymes	335
14.3.5	Assessing the Sperm Comet	336
14.3.6	Comet-FISH on Sperm	337
14.3.7	Cryopreserved <i>versus</i> Fresh Sperm	338
14.3.8	Viability Considerations	338
14.3.9	Statistical Analysis	339
14.4	Utilising Male Germ Cells with the Comet Assay	339
14.4.1	<i>In Vivo</i> Comet Assay	348
14.4.2	<i>In Vitro</i> Comet Assay	349
14.5	The Sperm Comet Assay <i>versus</i> Other Assays Used in Reproductive Toxicology	350
14.6	Conclusions	351
	Acknowledgements	351
	References	351

Section IV: Regulatory, Imaging and Statistical Considerations

Chapter 15 Comet Assay – Protocols and Testing Strategies 373

Andreas Hartmann and Günter Speit

15.1	Introduction	373
15.2	Applications of the <i>In Vivo</i> Comet Assay for Regulatory Purposes	374
15.3	Recommendations for Test Performance	375
15.3.1	Genetic Endpoint of the Comet Assay	375
15.3.2	Basic Considerations for Test Protocol	376
15.3.3	Selection of Tissues and Cell Preparation	377
15.3.4	Image Analysis	378

15.3.5	Assessment of Cytotoxicity – A Potential Confounding Factor	378
15.3.6	Ongoing Validation Exercises	379
15.4	Applications of the <i>In Vivo</i> Comet Assay for Regulatory Purposes	380
15.4.1	Follow-Up Testing of Positive <i>In Vitro</i> Cytogenetics Assays	380
15.4.2	Follow-Up Testing of Tumourigenic Compounds	381
15.4.3	Assessment of Local Genotoxicity	382
15.4.4	Assessment of Germ Cell Genotoxicity	382
15.4.5	Assessment of Photogenotoxicity	382
15.4.6	Genotoxicity Testing of Chemicals	384
15.5	Conclusions	384
	References	385

Chapter 16 Imaging and Image Analysis in the Comet Assay **390**

Mark Browne

16.1	Introduction	390
16.1.1	Experimental Design and Applications	390
16.2	Comet Sample Preparation	391
16.3	Comet Fluorescence Staining and Visualisation	392
16.4	Fluorescence Microscopy for Comet Imaging	395
16.4.1	Light Sources	396
16.4.2	Epifluorescence Light Path	398
16.4.3	Fluorescence Filter Sets	399
16.4.4	Microscope Objectives	401
16.4.5	Beam-Splitter and C-Mount Adapter	402
16.5	Image Detection – CCD, EMCCD and CMOS Cameras	403
16.5.1	Practical Matters	407
16.6	Image Processing and Comet Scoring	408
16.6.1	Image Analysis	409
16.6.2	Segmentation	410
16.6.3	Further Segmentation – Identifying Head and Tail of the Comet	413
16.6.4	Analysis of the Comet, Head and Tail Distributions	413
16.6.5	Comet Analysis – Other Approaches	416
16.7	How Many Cells, How Many Replicates?	417
16.7.1	Data Presentation and Preparation for Analysis	418
16.7.2	Statistical Analyses	419
16.7.3	Data Storage and Management	420

16.8	Conclusions	421
	References	421
Chapter 17	Statistical Analysis of Comet Assay Data	424
	<i>David P. Lovell</i>	
17.1	Introduction	424
17.2	Experimental Design and Statistical Analysis	425
17.3	Study Design	425
17.4	Endpoints	427
17.5	The Experimental Unit and Experimental Design	431
17.6	Statistical Methods	432
17.7	Use of Control Groups	436
17.8	Assessment of Results	437
17.9	Multiple Comparison Issues	438
17.10	Power and Sample Size	441
17.11	Human Studies	443
17.12	Standardisation and Interlaboratory Comparisons	445
	References	446
Subject Index		451

SECTION I: GENESIS OF COMET ASSAY

CHAPTER 1

The Comet Assay: A Versatile Tool for Assessing DNA Damage

ALOK DHAWAN*, MAHIMA BAJPAYEE AND
DEVENDRA PARMAR

Developmental Toxicology Division, Indian Institute of Toxicology Research
(Formerly Industrial Toxicology Research Centre), P.O. Box 80, M.G. Marg,
Lucknow, 226 001, India

1.1 Introduction

New chemicals are being added each year to the existing burden of toxic substances in the environment. This has led to increased pollution of ecosystems as well as deterioration of the air, water and soil quality. Excessive agricultural and industrial activities adversely affect biodiversity, threatening the survival of species in a particular habitat as well as posing disease risks to humans. Some of the chemicals, *e.g.* pesticides and heavy metals, may be genotoxic to the sentinel species and/or to nontarget species, causing deleterious effects in somatic or germ cells. Test systems that help in hazard prediction and risk assessment are important to assess the genotoxic potential of chemicals before their release into the environment or for commercial use as well as DNA damage in flora and fauna affected by contaminated/polluted habitats. The Comet assay has been widely accepted as a simple, sensitive and rapid tool for assessing DNA

*Corresponding author

damage and repair in individual eukaryotic as well as some prokaryotic cells, and it has increasingly found application in diverse fields ranging from genetic toxicology to human epidemiology.

This review is an attempt to comprehensively encase the use of the Comet assay in different models from bacteria to man, employing diverse cell types to assess the DNA-damaging potential of chemicals and/or environmental conditions. Sentinel species are the first to be affected by adverse changes in their environment. Determination of DNA damage using the Comet assay in these indicator organisms would thus provide information about the genotoxic potential of their habitat at an early stage. This would allow for intervention strategies to be implemented for prevention or reduction of deleterious health effects in the sentinel species as well as in humans.

Ostling and Johanson¹ were the first to quantify DNA damage in cells using a microgel electrophoresis technique, known as the single-cell gel electrophoresis (SCGE) or Comet assay. However, the neutral conditions that they used allowed the detection of only double strand breaks in the DNA. Later, the assay was adapted under alkaline conditions by Singh *et al.*,² which led to a sensitive version of the assay that could assess both double- and single-strand DNA breaks as well as alkali-labile sites expressed as frank strand breaks in the DNA. Since its inception, however, the assay has been modified at various steps (lysis, electrophoresis) to make it suitable for various kinds of damage in different cells.^{3,4} The assay is now a well-established, simple, versatile, rapid, visual, and a sensitive, extensively used tool to assess DNA damage and repair, quantitatively as well qualitatively in individual cell populations.⁵ Some other lesions of DNA damage such as DNA cross-linking (*e.g.* thymidine dimers) and oxidative DNA damage may also be assessed using lesion specific antibodies or specific DNA repair enzymes in the Comet assay. It has gained wide acceptance as a valuable tool in fundamental DNA damage and repair studies,⁴ genotoxicity testing⁶ and human biomonitoring.^{7,8}

Relative to other genotoxicity tests, such as chromosomal aberrations, sister chromatid exchanges, alkaline elution and the micronucleus assays, the advantages of the Comet assay include its demonstrated sensitivity for detecting low levels of DNA damage (one break per 10^{10} Daltons of DNA⁹), requirement for small number of cells ($\sim 10\,000$) per sample, flexibility to use proliferating as well as nonproliferating cells, low cost, ease of application, and the short time needed to complete a study. It can be conducted on cells that are the first site of contact with mutagenic/carcinogenic substances (*e.g.* oral and nasal mucosal cells). The data generated at the single-cell level allow for robust types of statistical analysis.

A limitation of the Comet assay is that aneugenic effects, which may be a possible mechanism for carcinogenicity,¹⁰ and epigenetic mechanisms (indirect) of DNA damage such as effects on cell-cycle checkpoints are not detected. The other drawbacks such as single cell data (which may be rate limiting), small cell sample (leading to sample bias), technical variability and interpretation are some of its disadvantages. However, its advantages far outnumber the

disadvantages and hence it has been widely used in fields ranging from molecular epidemiology to genetic toxicology.

The present review deals with various models ranging from bacteria to man used in the Comet assay for assessing DNA damage (Figure 1.1).

1.2 Bacteria

The first study to assess the genetic damage in bacteria treated with 12.5–100 rad of X-rays, using the Comet assay was conducted by Singh *et al.*¹¹ In the study, the neutral Comet assay was used for direct (visual) determination of DNA double-strand breaks in the single electrostretched DNA molecule of *Escherichia coli* JM101. A significant increase in DNA breaks was induced by a dose as low as 25 rad, which was directly correlated to X-ray dosage. The study supported a hypothesis that the strands of the electrostretched human DNA in the Comet assay represented individual chromosomes.

1.3 Plant Models

Plant bioassays are important tests that help detect genotoxic contamination in the environment.¹² Plant systems can provide information about a wide range of genetic damage, including gene mutations and chromosome aberrations. The mitotic cells of plant roots have been used for the detection of clastogenicity of environmental pollutants, especially for *in situ* monitoring of water contaminants. Roots of *Vicia faba* and *Allium cepa* have long been used for assessment of chromosome aberrations¹³ and micronuclei.¹⁴ During the last decade, the Comet assay has been extensively applied to plants (leaves, shoots, and roots) to detect DNA damage arising due to chemicals and heavy metals in polluted soil (Table 1.1).

1.3.1 The Comet Assay in Lower Plants

1.3.1.1 Fungi

Schizosaccharomyces pombe has been used as a model organism to investigate DNA damage due to chlorinated disinfectant, alum and polymeric coagulant mixture in drinking-water samples.¹⁵ The authors observed a significantly higher ($P < 0.001$) DNA damage in chlorinated water (*i.e.* tap water) when compared to untreated (negative control) or distilled water (laboratory control). Hahn and Hock¹⁶ used mycelia of *Sordaria macrospora* grown and treated with a variety of DNA-damaging agents directly on agarose minigels for the assessment of genotoxicity using the Comet assay. DNA-strand breaks were detected by an increase in the DNA migration from the nucleus. This model allowed for the rapid and sensitive detection of DNA damage by a number of chemicals simultaneously. *Saccharomyces cerevisiae* has also been employed for successful investigation of DNA damage at low concentrations of chemicals.²⁰²




Figure 1.1 Schematic diagram of the use of the Comet assay in assessing DNA damage in different models from bacteria to humans.

Table 1.1 Comet assay for assessment of DNA damage – bacteria to humans.

<i>Model</i>	<i>Agent tested</i>	<i>Cell used</i>	<i>DNA damage</i>	<i>Ref.</i>
Bacteria				
<i>Escherichia coli</i> JM101	X-rays	Whole organism <i>in vivo</i>	↑	11
<i>Euglena gracilis</i>	1-Methyl-3-nitro-1-nitrosoguanidine (MNNG), benzo[a]pyrene, mitomycin C and actinomycin D.	Whole organism <i>in vivo</i>	↑	18
<i>Chlamydomonas reinhardtii</i>	4-Nitroquinoline-1-oxide (4-NQO), N-nitrosodimethylamine, and hydrogen peroxide	Whole organism <i>in vivo</i>	↑	17
<i>Rhodomonas</i>	UV (UVA + UVB) radiation	Whole organism <i>in vivo</i>	↑	19
<i>Vicia faba</i>	N-methyl-N-nitrosourea (MNU) and methyl methanesulfonate (MMS)	Root tip meristematic cells	↑	21
Tobacco (<i>Nicotiana tabacum</i> L)	Ethyl methanesulfonate	Nuclei from leaf tissue	↑	22
	Age	Leaf nuclei		23
	Kinetics of DNA repair	Leaf nuclei		24
	Ethyl methanesulfonate (EMS) and N-ethyl-N-nitrosourea (ENU), maleic hydrazide (MH)	Whole roots <i>in vivo</i>	↑	25,26
	O-phenylenediamine (o-PDA), hydrogen peroxide and ethyl methanesulfonate (EMS).	Isolated root nuclei	–	27
	Heavy metal (Cd, Cu, Pb, and Zn)	Leaf nuclei	↑	28
	Polychlorinated biphenyls	Leaf nuclei	↑	29
Potato plants (<i>Solanum tuberosum</i> var. <i>Korela</i>)	Heavy metal (Cd, Cu, Pb, and Zn)	Nuclei from leaf tissue	↑	28
<i>Phaeoecolus vulgaris</i>	Uranium	Root or shoot cells	–	30
<i>Impatiens balsamina</i>	Cr ⁶⁺ and airborne particulate	Stem, root and leaves	↑	31
<i>Bacopa momieri</i> L.	Ethyl methanesulfonate, methyl methanesulfonate, cadmium	Nuclei isolated from roots and leaves	↑ dose- and time-dependent roots > leaves	32
Animal models				
<i>Tetrahymena thermophila</i>	Phenol, hydrogen peroxide, and formaldehyde, influent and effluent water samples	Whole animal <i>in vivo</i>	↑	33

Table 1.1 (Continued).

Model	Agent tested	Cell used	DNA damage	Ref.
Invertebrates – Bivalves				
Freshwater bivalve zebra mussel (<i>Dreissena polymorpha</i>)	Polybrominated diphenyl ethers (pbdes)	Haemocytes	↑ ↑	34
	Sodium hypochlorite, chlorine dioxide and peracetic acid		↑	35
	Pentachlorophenol		↑	36
	Varying temperatures		↑	37
<i>Mytilus edulis</i>	Polluted waters		↑	38
	Cadmium (Cd) and chromium (Cr)		—	39
	Styrene	Gills	↑	40
	Tritium	Haemolymph cells	↑	41
	Marine waters (Denmark), French Atlantic Coast	Haemocytes	↑	42
		Gill and haemolymph	↑	
	Polycyclic aromatic hydrocarbons	Gill and haemocytes	↑	44
	Seasonal variation	Haemocytes	↑	45
Freshwater mussels (<i>Unio tumidus</i>)	Polyphenols	Digestive gland cells	↑	46
Golden mussel (<i>Limno-perna fortunei</i>)	Guaíba Basin water	Haemocytes	↑	47
Bivalve mollusc (<i>Scapharca inaequivalvis</i>)	Organotin compounds (MBTC, DBTC and TBTC)	Erythrocytes	↑	48
<i>Mytilus galloprovincialis</i>	Environmental stress	Haemocytes	↑	49
	Heavy oil spill	Gills		50
	Cadmium	Digestive gland cells		51
	Hydrostatic pressure change	Haemocytes and gill tissues	↑	52,53
Vent mussels (<i>Bathymodiolus azoricus</i>)	Benzo[a]pyrene	Haemocytes	↑	54
Green-lipped mussel (<i>Perna viridis</i>)	Chemicals used in lawn care (atrazine, glyphosate, carbaryl, and copper)	Glochidia	↑	55
Freshwater mussel (<i>Urtterbackia imbecillis</i>)	Cryopreservation	Spermatozoa	↑	56

Manila clam (<i>Tapes semidecussatus</i>)	Sediment-bound contaminants	Haemolymph, gill and digestive gland	↑	57,58
Clams (<i>Mya arenaria</i>)	Petroleum hydrocarbons	Haemocytes and digestive gland cells	—	59
Invertebrates – Earthworms				
<i>Eisenia foetida</i>	Chemical-treated soil	Coelomocytes	↑ dose dependent	60
	Soil from coke ovens	Coelomocytes	↑	61
	Soil from industrialised contaminated areas	Coelomocytes	↑	62
	Sediment from polluted river	Coelomocytes	↑	63
	Wastewater-irrigated soil	Coelomocytes	↑	64
	Commercial parathion	Coelomocytes	↑	65
	Imidacloprid and RH-5849	Sperm cells	↑	66
	PAH-contaminated soil and hydrogen peroxide, cadmium (<i>in vitro</i>)	Eleocytes	↑	67
	Nickel chloride	Coelomocytes	↑	68
<i>Aporrectodea longa</i> (Ude)	Soil samples spiked with benzo[a]pyrene (B[a]P) and/or lindane	Intestine and crop/gizzard cells	↑ intestine > crop	69
Other Invertebrates				
Fruit fly (<i>Drosophila melanogaster</i>)	Ethyl methanesulfonate (EMS), methyl methanesulfonate (MMS), N-ethyl-N-nitrosourea (ENU) and cyclophosphamide (CP)	Gut and brain cells of first instar larvae	↑	70,71
	Cypermethrin		↑	72
	Leachates of industrial waste		↑	71
	Cisplatin		↑	73
	Dispersed crude oil	Coelomocytes	↑ concentration-dependent	74
Sea urchins (<i>Strongylocentrotus droebachiensis</i>)	UV, benzo[a]pyrene, and cadmium	Embryos	↑ damage and decreased repair	75
Grass shrimp, (<i>Palaemonetes pugio</i>)	Estuarine sediments	Hepatopancreas	↑	76
	Coal combustion residues			77
Sea anemone (<i>Anthopleura elegantissima</i>)	Hydrogen peroxide ethylmethanesulfonate (EMS) or benzo[a]pyrene (B[a]P)	Blood cells	↑ dose response	78

Table 1.1 (Continued).

Model	Agent tested	Cell used	DNA damage	Ref.
Vertebrates – Fishes				
Chub (<i>Leuciscus cephalus</i>)	PAHs, PCBs, organochlorine pesticides (OCPs), as well as heavy metals	Hepatocytes	↑	79
Estuarine mullet (<i>Mugil sp.</i>) and sea catfish (<i>Netuma sp.</i>)	Exhaustive exercise Organochlorine pesticides and heavy metals High temperature	Erythrocytes Erythrocytes	↑ ↑	80 81,82
Fresh water teleost fish (<i>Mystus vittatus</i>)	Endosulfan	Gill, kidney, and erythrocytes	↑ in all cells	83
Eastern mudminnow (<i>Umbra pinnacea</i> L.)	Rhine water for 11 days	Blood erythrocytes	↑	84
Neotropical fish (<i>Prochilodus lineatus</i>)	Diesel water soluble fraction acute (6, 24 and 96 h) and subchronic (15 days) exposures,	Erythrocytes	↑	85
Freshwater goldfish (<i>Carassius auratus</i>)	Technical herbicide Roundup containing glyphosphate salt ADDB and PBTA-6 Sediment collected from polluted sites in Cork Harbour (Ireland)	Erythrocytes	↑↑ dose dependent	86
Turbot (<i>Scophthalmus maximus</i> L.)	Contaminated estuary waters	Hepatocytes	↑	87 88
Brazilian flounder (<i>Paralichthys orbignyanus</i>)		Blood cells	↑↑	89
Bullheads (<i>Ameiurus nebulosus</i>)	Polycyclic aromatic hydrocarbon (PAH) and polychlorinated biphenyl (PCB) polluted waters	Erythrocytes	↑	90
Carp (<i>Cyprinus carpio</i>)	Polycyclic aromatic hydrocarbon (PAH) and polychlorinated biphenyl (PCB) polluted waters	Erythrocytes	↑	90
Brown trout (<i>Salmo trutta fario</i>)	PCB77 (3,3',4,4'-tetrachlorobiphenyl)	Erythrocytes	–	91
Marine flatfish	Ethyl methanesulfate	Blood, gill, liver and kidney	↑ in all tissues	92
Trout (<i>Oncorhynchus mykiss</i>)	Cryopreservation (freeze–thawing)	Spermatozoa	Slight ↑	93

European eel (<i>Anguilla anguilla</i>)	Benzo[a]pyrene, Arochlor 1254, 2,3-7,8-tetrachlorodibenzo-p-dioxin and beta-naphthoflavone	Erythrocytes	↑	94
Eelpout (<i>Zoarces viviparus</i>)	Oil spill (PAH)	Nucleated erythrocytes	↑	95
Gilthead sea bream (<i>Sparus aurata</i>)	Copper	Erythrocytes	↑ ↑	96
Dab (<i>Limanda limanda</i>)	PAHs and PCBs polluted waters of English Channel	Blood cells	↑ in adults and males	43
Hornyhead turbot (<i>Pleuronichthys verticalis</i>)	Gender and age Sediments collected from a natural petroleum seep (pahs)	Liver cells	↑	97
In vitro				
Carp (<i>Cyprinus carpio</i>)	Organic sediment extracts from the North Sea (Scotland)	Leukocytes	↑	98
Trout (<i>Oncorhynchus mykiss</i>)	Cadmium Oxidative stress and its prevention by indolinic and quinolinic nitroxide radicals	Hepatocytes Erythrocytes	↑ ↑	99 100
	Tannins		↓	101
Zebrafish (<i>Danio rerio</i>)	Diaryl tellurides and ebselen (organoselenium)		↓	102
Rainbow trout hepatoma cell line (RTH-149)	Surface waters of German rivers, Rhine and Elbe		↓	103
Rainbow trout gonad (RTG-2) cell line	Water samples from the polluted Kishon river (Israel)	Hepatocytes and gill cells Liver	↑ ↑	104
	4-Nitroquinoline-N-oxide N-methyl-N'-nitro-N-nitrosoguanidine, benzo[a]pyrene, nitrofurantoin, 2-acetylaminofluorene, and dimethylnitrosamine, and surface waters	Gonad	↑ dose-dependent response	105
		Epitheloid liver		
Vertebrates – Amphibians				
Amphibian larvae (<i>Xenopus laevis</i> and <i>Pleurodeles waltli</i>)	Cadmium (CdCl ₂) Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	Erythrocytes	↑ concentration and time dependent	106,107
Amphibian larva (<i>Xenopus laevis</i>)	Benzo[a]pyrene, ethyl methanesulfonate methyl methanesulfonate, aqueous extracts of five sediments from French channels	Erythrocytes	– ↑	108,109

Table 1.1 (Continued).

Model	Agent tested	Cell used	DNA damage	Ref.
Toad (<i>Bufo raddei</i>)	Petrochemical (mainly oil and phenol) polluted area	Liver cells and erythrocytes	↑	110
Toad (<i>Xenopus laevis</i> , and <i>Xenopus tropicalis</i>)	Bleomycin-induced DNA damage and repair	Splenic lymphocytes	↑ DNA damage <i>X. tropicalis</i> > <i>X. laevis</i> DNA repair in <i>X. laevis</i> > <i>X. tropicalis</i>	111
Tadpoles of <i>Rana N. Hallowell</i>	Imidacloprid [1-(6-chloro-3-pyridylmethyl)-N-nitro-imidazolidin-2-ylideneamine] and RH-5849 [2'-benzoyl-1'-tert-butylbenzoylhydrazinell]	Erythrocytes	↑	112
<i>Rana hexadactyla</i> tadpoles	Sulfur dyes (Sandopel Basic Black BHLN, Negrosine, Dermalpel Black FNI, and Turquoise Blue) used in the textile and tannery industries	Erythrocytes	↑↑	113
Bullfrog (<i>Rana catesbeiana</i>) tadpoles	Herbicides AAtrex Nine-O (atrazine), Dual-960E (metolochlor), Roundup (glyphosate), Sencor-500F (metribuzin), and Amsol (2,4-D amine)	Erythrocytes	↑↑	114
Tadpole <i>Rana clamitans</i> <i>Rana pipiens</i>	Agricultural regions, Industrial regions	Erythrocytes	↑ industrial regions > agricultural regions	115
<i>In vitro</i>				
<i>Xenopus laevis</i>	High peak-power pulsed electromagnetic field	Erythrocytes	↑ due to rise in temperature	116
Wild nesting white storks (<i>Ciconia ciconia</i>)	Heavy metals and arsenic	Blood cells	↑ correlated with arsenic	117
Black kites (<i>Milvus migrans</i>)	Toxic acid mining waste rich in heavy metals	Blood cells	↑↑	118,119,120
	Heavy metals and arsenic	Blood cells	↑↑ correlated with copper and cadmium	117
Turkey	Toxic acid mining waste rich in heavy metals	Sperm	↑ (2-10 fold)	118,119,120
	Short-term storage		↑	121

Chicken	T-2 toxin and deoxynivalenol (DON) Storage conditions (4 °C)	Spleen leukocytes Liver and breast muscle cells	↑ ↑ liver cells > breast muscle cells	122 123
Vertebrates – Rodents				
Aldh2 knockout mice P53(+/-) mice	Ethanol Melfhalan	Hepatic cells Liver, bone marrow, peripheral blood and the distal intestine Epidermal cells	↑ oxidative damage DNA cross-links in all cells tested	124 125
SKH-1 mice	UV A + Fluoroquinolones (clinafloxacin, lomefloxacin, ciprofloxacin) UVA + 8-methoxypsoralene (8-MOP) Ageing Diesel exhaust particles <i>Trypanosoma cruzi</i> infection Lead acetate	Aorta, liver, and lung Peripheral blood, liver, heart and spleen cells Nasal epithelial cells, lung, whole blood, liver, kidney, bone marrow, brain and testes Blood, bone marrow cells and liver Brain, liver, kidney, bone marrow, blood, spleen Stomach cells, hepatocytes, kidney and testicle cells Brain cells Brain cells	↑ ↑ for fluoroquinolones ↓ for MOP ↑ Oxidative damage in liver – in lung or aorta ↑ in heart and spleen ↑ in all organs on prolonged exposure – in testes	126 127, 128 129 130
Swiss albino mice	Sanguinarine alkaloid, argemone oil Cypermethrin Steviol		↑ dose dependent in blood and bone marrow ↑ ↑	131, 132 133 134
	Apomorphine, 8-oxo-apomorphine-semiquinone Ethanol, grape seed oligomer and polymer pro-cyanidin fractions		– ↑ ↓ ethanol-induced protection by grape seed	135 136
Male CBA mice	Pesticide formulations (Bravo and Gesaprim)	Hepatic cells, bone marrow cells, spleen cells	↑ ↑	137
Isogenic mice	Sulfonamide, protozoan parasite <i>Toxoplasma gondii</i>	Peripheral blood cells, liver cells and brain cells	↑ in peripheral blood cells	138
Cirrhotic rats	Rutin and quercetin	Bone marrow cells	↑ ↑	139

Table 1.1 (Continued).

Model	Agent tested	Cell used	DNA damage	Ref.
<i>In vitro</i>				
FE1 muta mouse lung epithelial cell line	Carbon black	Lung epithelial cell line	↑	140
L5178Y mouse lymphoma cells	Ketoprofen, promazine, chlorpromazine, dacarbazine, acridine, lomefloxacin, 8-methoxypsoralen, chlorhexidine, titanium dioxide, octylmethoxycinnamate	Lymphoma cells	Positive with phototoxic compound	141
Murine primary cultures of brain cells and a continuous cell line of astrocytes	Xanthine/xanthine oxidase, hydrogen peroxide superoxide dismutase, catalase, or ascorbic acid	Brain cells	↓ by antioxidants	142
Chinese hamster ovary cell line (CHO)	Endosulfan Cypermethrin, pendimethalin, dichlorovous	Ovary cells	↑	143 144
Humans – Clinical				
Breast cancer patients and controls	Radiosensitivity	Peripheral blood mononuclear cells	↑	145
Breast cancer patients and controls	Radiosensitivity	Peripheral blood mononuclear cells	↑ and reduced DNA repair	146
Normal individuals	Chlorhexidine	Buccal epithelial cells and peripheral blood lymphocytes	↑	147
Transitional cell carcinoma patients and controls	DNA-strand breaks	Exfoliated cells extracted from bladder washing	↑ in patients	148
<i>Axvia telangiectasia</i> heterozygote	X-irradiation	Peripheral leukocytes	↑ (~3 times high) in patients	149

Nijmegen breakage syndrome (NBS) patients	X-irradiation	Peripheral blood mononuclear cells	↑ in patients	150
Alzheimer disease patients	–	Peripheral blood mononuclear cells	↑ in patients	151
Breast cancer patients	–	Peripheral blood mononuclear cells	↑ in patients	152
Type 2 diabetes mellitus	Oxidative DNA damage	Peripheral blood cells	↑	153
Cancer (testicular cancer, lymphoma and leukemia) patients	DNA integrity	Spermatozoa	Decreased DNA integrity	154
Humans – Dietary intervention				
Healthy subjects	Tomato drink	Blood lymphocytes	↓	155
	Green vegetables			156
	Grape juice			157
Smokers	Vitamin C supplementation	Blood lymphocytes	↓	158
Technical anesthesiology staff	Vitamin E and vitamin C	Blood lymphocytes	↓ in oxidative damage	159
Humans – Occupational				
Airport personnel	Jet fuel vapours, jet fuel combustion products	Exfoliated buccal cells and lymphocytes	↑	160
Agricultural workers	Pesticides	Lymphocytes	–	161
				162, 163
Rubber factory workers	Substances used in the rubber industry	Peripheral blood	↑ in exposed population	164
Outdoor workers in Mexico cities	Air pollutants	Blood lymphocytes	↑	165
Rickshaw pullers	Exhaustive exercise	Lymphocytes	↑	166
Nuclear medicine personnel	Ionising radiation	Peripheral blood leukocytes	↑	167
Workers	Polycyclic aromatic hydrocarbons (PAH)	Human T- and B-lymphocytes, and granulocytes	↑ B-lymphocytes > T-lymphocytes > granulocytes	168
	Benzene in printing	"	↑	169
	Lead (Pb) and cadmium (Cd)	"	↑	170
	Asbestos cement plant	Peripheral lymphocytes	↑	171
	Fenvalerate (FE) exposure	Sperm	↑	172
	Organic solvents	Peripheral blood	↑	173

Table 1.1 (Continued).

Model	Agent tested	Cell used	DNA damage	Ref.
Nurses	Coke oven emissions (coe)	Blood lymphocytes		174
	Welders (Cd, Co, Cr, Ni, and Pb)	Lymphocytes		175
	Pesticide formulators (organophosphorus pesticides)	Lymphocytes	↑	176
	Copper smelters (inorganic arsenic)	Leukocytes	↑	177
	Chrome-plating workers (chromium VI)	Lymphocytes	↑↑	178
	Workers in foundry and pottery (silica)	Lymphocytes	↑	179
Nurses	5-Fluorouracil, cytarabine, gemcitabine, cyclophosphamide, and ifosfamide	Lymphocytes	Slight ↑	180
Humans – Lifestyle				
Normal individuals Active and passive smokers	Endurance exercise	Lymphocytes	↑	181
	Smoking	Lymphocytes	↑	182
Normal individuals	Smoking	Lymphocytes	↑	183–186
	Diet (vegetarian or non-vegetarian)	Lymphocytes		
Rural Indian women	Biomass fuels	Lymphocytes	↑	187
Normal individuals	Benzo[a]pyrene, beta-naphthoflavone (BNF)	Human umbilical vein endothelial cells (HUEVC)	↑	188
In vitro				
Episkin	UV, Lomefloxacin and UV or 4-nitroquinoline-N-oxide (4NQO) and protection by Mexoryl	Skin fibroblast cells	↑ reduced by Mexoryl	189
Sperms	Reproductive toxins	Male germ cells	↑	190, 191
	2-Amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP), its N-hydroxy metabolite (N-OH-PhIP) and benzo[a]pyrene (B[a]P)	Prostate cells	↑ dose related	192
Human keratinocytes MCF-7 cells JM1 cells HepG2 cells	UVA or UVB	Skin cells	↑	193
	Estradiol	Breast cells	↑	194
	Estradiol	Lymphoblast cells	↑ concentration dependent	194
	Endosulfan	Liver cells	–	195
	Indirect acting genotoxins (cyclophosphamide)			196

Miniorgan cultures of human inferior nasal turbinate epithelia	Sodium dichromate, N-nitrosodiethylamine (NDEA) and N-methyl-N-nitroso-guanidine (MNNG) Mono(2-ethylhexyl) phthalate (MEHP), benzo[a]pyrene-7,8-diol-9,10-epoxide (BPDE), or N-methyl-N'-nitro-N-nitrosoguanidine (MNNG).	Nasal cells	↑ with sodium dichromate and MNNG – with NDEA ↑ with BPDE and MNNG – with MEHP	197 198
Human lymphocytes	Heterocyclic amine and prevention by monomeric and dimeric flavanols and black tea polyphenols	Lymphocyte	↓ in oxidative damage ↑ ↑	199
	C ₆₀ Fullerenes			200
	Municipal sludge leachates			201

↑ significant increase in DNA damage; ↑↑ highly significant increase in DNA damage; ↓ decrease in DNA damage; – no DNA damage reported

1.3.1.2 Algae

Aquatic unicellular plants like algae provide information on the potential genotoxicity of the water in which they grow. Being single celled they can be used as a model for assessment of DNA damage and monitoring of environmental pollution utilising the Comet assay. Unicellular green alga *Chlamydomonas reinhardtii* was used for evaluation of DNA damage due to known genotoxic chemicals and also demonstrated that oxidative stress was better managed by the algal cells under light rather than dark conditions.¹⁷ The Comet assay was found to be useful for evaluating chemically induced DNA damage and repair in *Euglena gracilis* and responses were more sensitive than those of human lymphocytes under the same treatment conditions.¹⁸ The ease of culturing and handling *E. gracilis* as well as its sensitivity, makes it a useful tool for testing the genotoxicity of chemicals and monitoring environmental pollution. A modified version of the Comet assay was used as an alternative technique to assess DNA damage due to UV radiation in *Rhodomonas* sp. (*Cryptophyta*), a marine unicellular flagellate.¹⁹

1.3.2 The Comet Assay in Higher Plants

Vicia faba has been widely used for the assessment of DNA damage using the Comet assay. Strand breaks and abasic (AP) sites in meristematic nuclei of *V. faba* root tips were studied by the neutral and alkaline Comet assay.^{20,21} The alkaline electrophoresis procedure was found to be most sensitive at low doses, while the neutral electrophoresis procedure yielded an optimal dose–response curve within a wider dose range. Angelis *et al.*²⁰ also suggested that the Comet assay was able to detect a phenomenon resembling clastogenic adaptation at the molecular level. Gichner and Plewa²² developed a sensitive method for isolation of nuclei from leaf tissue of *Nicotiana tabacum*. The method resulted in high resolution and constant low tail moment values for negative controls, and hence it could be incorporated as a test for *in situ* plant environmental monitoring.²²

The Comet assay has also been used to study the effect of age of plant on DNA integrity²³ as well as the kinetics of DNA repair²⁴ in isolated nuclei from leaves of tobacco plants. A small but significant increase in DNA damage compared to controls was noted in heterozygous tobacco and potato plants grown on soil contaminated with heavy metals.²⁸ The tobacco and potato plants with increased DNA damage were also found to be severely injured (inhibited growth, distorted leaves), which may be associated with necrotic or apoptotic DNA fragmentation. No DNA damage was observed in the root or shoot cells of *Phaseolus vulgaris* treated with different concentrations of uranium.³⁰ The ornamental plant *Impatiens balsamina* was used as a model to understand the genotoxic effect of Cr⁶⁺ and airborne particulate matter,³¹ which produced increased strand breaks in plant parts (stem, root and leaves). Thus, this plant could be used for environmental biomonitoring studies involving air pollution and heavy metals.

The major drawback with plant models was the fact that exposure needs to be given in the soil and it is difficult to say whether the result demonstrates synergies with other chemicals in the soil or nonavailability of the toxicant due to its soil binding affinity. Therefore, Vajpayee *et al.*³² used *Bacopa monnieri* L., a wetland plant, as a model for the assessment of ecogenotoxicity using the Comet assay. *In vivo* exposure to cadmium (0.01–500 μ M) for 2, 4, and 18 h resulted in dose- and time-dependent increases in DNA damage in the isolated roots and leaf nuclei, with roots showing greater DNA damage than leaves. *In vitro* (acellular) exposure of nuclei from leaves of *B. monnieri* to 0.001–200 μ M cadmium resulted in significant ($P < 0.05$) levels of DNA damage.

These studies revealed that DNA damage measured in plants using the Comet assay is a good model for assessment of genotoxicity of polluted environments since *in situ* monitoring and screening can be accomplished. Higher plants can be used as an alternative first-tier assay system for the detection of possible genetic damage resulting from polluted waters/effluents due to industrial activity or agricultural run offs.

1.4 Animal Models

To assess safety/toxicity of chemicals/finished products, animal models have long been used. With the advancements in technology, knockouts and transgenic models have become common to mimic the effects in humans. The Comet assay has globally been used for assessment of DNA damage in various animal models (Table 1.1).

1.4.1 Lower Animals

Tetrahymena thermophila is a unicellular protozoan, widely used for genetic studies due to its well-characterised genome. Its uniqueness lies in the fact that it has a somatic and a germ nucleus in the same cell. Therefore it has been validated as a model organism for assessing DNA damage using a modified Comet assay protocol standardised with known mutagens such as phenol, hydrogen peroxide, and formaldehyde.³³ The method was then used for the assessment of genotoxic potential of influent and effluent water samples from a local municipal wastewater treatment plant.³³ The method provided an excellent, low-level detection of genotoxins and proved to be a cost-effective and reliable tool for genotoxicity screening of wastewater.

1.4.1.1 Invertebrates

Studies have been carried out on various aquatic (marine and freshwater) and terrestrial invertebrates (Table 1.1). The genotoxicity assessment in marine and freshwater invertebrates using the assay has been reviewed.^{203–205} Cells from haemolymph, embryos, gills, digestive glands and coelomocytes from mussels (*Mytilus edulis*⁴²), zebra mussel (*Dreissena polymorpha*), clams

(*Mya arenaria*), and polychaetes (*Nereis virens*), have been used for ecogenotoxicity studies using the Comet assay. DNA damage has also been assessed in earthworms^{61,63} and fruit flies, *Drosophila*.^{72,206} The Comet assay has been employed to assess the extent of DNA damage in organisms at polluted sites in comparison to those at reference sites in the environment. In the laboratory it has been widely used as a mechanistic tool to determine pollutant effects and mechanisms of DNA damage.⁷⁸

1.4.1.2 The Comet Assay in Mussels

Freshwater and marine mussels have been used to study the adverse effect of contaminants in the aquatic environment as they are important pollution-indicator organisms. These sentinel species are adversely affected by the pollution of the water bodies and thus provide the potential for environmental biomonitoring. The Comet assay in mussels has been used to detect a reduction in water quality caused by chemical pollution.^{41,42,49,207} *Mytilus edulis* has been widely used for Comet assay studies to evaluate DNA-strand breaks in gill and digestive gland nuclei due to polycyclic aromatic hydrocarbons (PAHs) including benzo[a]pyrene (B[a]P),⁴⁴ and oil spills with petroleum hydrocarbons.⁵⁹ The DNA damage was found to be elevated in the exposed mussels. However, the damage returned to normal levels, after continued exposure to a high dose (20 ppb-exposed diet) of B[a]P for 14 days. This was attributed to an adaptive response in mussels to prevent the adverse effects of DNA damage.⁴⁴ The green lipped mussels (*Perna viridis*) also showed a similar result on exposure to B[a]P in water.⁵⁴

Significant levels of interindividual variability, including seasonal variations in DNA damage have been reported from some studies, both laboratory and field.^{45,49,208,209} Baseline monitoring thus has to be carried out over long time intervals. Temperature-dependent DNA damage was observed in haemocytes of freshwater mussel *Dreissena polymorpha*³⁷ showing that the mussels are sensitive towards change in water temperatures. Thus, monitoring ecogenotoxicity with these species should take into account variations in temperatures. Findings have also suggested that antioxidant supplementation can improve the sensitivity of the Comet assay by lowering the baseline damage in untreated animals.²⁰⁸

Villela *et al.*²¹⁰ used the golden mussel (*Limnoperna fortunei*) as a potential indicator organism for freshwater ecosystems due to its sensitivity to water contaminants. The Comet assay in haemocytes of freshwater Zebra mussel, *D. polymorpha* Pallas, was used as a tool in determining the potential genotoxicity of water pollutants.^{34–36,38} Klobucar *et al.*³⁸ suggested the use of the Comet assay in haemocytes from caged, nonindigenous mussels as a sensitive tool for monitoring genotoxicity of freshwater. DNA damage and repair studies in vent mussels, *Bathymodiolus azoricus*, have been carried out to study the genotoxicity of a naturally contaminated deep-sea environment.^{52,53} The vent mussels demonstrated similar sensitivity to environmental mutagens as that of coastal mussels and thus could be used for ecogenotoxicity studies of deep sea waters using the Comet assay.

In vitro Comet assay has also been used in cells of mussels. Dose–response increases in DNA-strand breakages were recorded in digestive gland cells²¹¹ haemocytes²¹² and gill cells^{208,212} of *M. edulis* exposed to both direct (hydrogen peroxide and 3-chloro-4-(dichloromethyl)-5-hydroxy-2[5H]-furanone) and indirect (B[a]P, 1-nitropyrene, nitrofurantoin and *N*-nitrosodimethylamine) acting genotoxicants. Digestive gland cells of *Unio tumidus* were also used for *in vitro* studies of DNA damage and repair due to pro-oxidative effect of poly-phenolic compounds.^{46,213} Wilson *et al.*²⁰⁸ demonstrated the potential application of the Comet assay to the gill cells of *M. edulis* as a potential *in vitro* screen for agents destined for release or disposal into the marine environment.

1.4.1.3 The Comet Assay in Other Bivalves

Coughlan *et al.*⁵⁷ showed that the Comet assay could be used as a tool for the detection of DNA damage in clams (*Tapes semidecussatus*) as biomonitor organisms for sediments. Significant DNA-strand breaks were observed in cells isolated from haemolymph, gill and digestive gland from clams exposed to polluted sediment.^{57,58} The Comet assay was used for the assessment of sperm DNA quality of cryopreserved semen in Pacific oysters (*Crassostrea gigas*) as it is widely used for artificial fertilisation.⁵⁶ Gielazyn *et al.*²¹⁴ demonstrated the use of lesion-specific DNA repair enzyme formamidopyrimidine glycosylase (Fpg) to enhance the usefulness and sensitivity of the Comet assay in studying oxidative DNA damage in isolated haemocytes from oysters (*Crassostrea virginica*) and clams (*Mercenaria mercenaria*).

The studies in mussels have shown the Comet assay to be a sensitive, but nonspecific, molecular biomarker of genotoxicity. One of the drawbacks when applying single-cell gel electrophoresis to field populations may be the adaptability of the animals to high concentrations of contaminants (*e.g.* B[a]P), which may pose a major problem.⁴⁴ Also, seasonal variation and temperature altered both DNA damage baseline levels in untreated animals and cell sensitivity towards environmental pollutants under *in vitro* conditions.^{37,58} The Comet assay detecting DNA-strand breaks has demonstrated that higher basal levels of DNA damage are observed in marine invertebrates, hence the protocol followed in these animals should be considered for biomonitoring the ecogenotoxicity of a region.²¹⁵

1.4.1.4 The Comet Assay in Earthworms

The Comet assay applied to earthworms is a valuable tool for monitoring and detection of genotoxic compounds in terrestrial ecosystems^{61,66} (Table 1.1). Since the worms feed on the soil they live in, they are a good indicator of the genotoxic potential of the contaminants present in the soil and thus used as a sentinel species. Verschaeye *et al.*⁶⁰ demonstrated a dose–response effect with the extent of DNA damage in coelomic leucocytes (coelomocytes) of

earthworms (*Eisenia foetida*) from soil treated with different chemicals as an indicator of soil pollution.

Coelomocytes from *E. foetida* demonstrated increased DNA damage when worms were exposed to soil samples from polluted coke oven sites,⁶¹ or industrialised contaminated areas⁶² and even sediment samples from polluted river system.⁶³ An insecticide, parathion, produced DNA-strand breaks at all time points and doses in the sperm cells of *E. foetida*⁶⁵ while dose-effect relationships were displayed by two pesticides, Imidacloprid and RH-5849 in the same species,⁶⁶ showing that pesticides could also have adverse effects on nontarget species. *In vitro* exposure of coelomocytes primary cultures to nickel chloride as well as whole animals either in spiked artificial soil water or in spiked cattle manure substrates exhibited increased DNA-strand breaks due to the heavy metal.⁶⁸ The leucocytes cells, a subset of coelomocytes produced increased DNA-strand breaks under both *in vitro* and *in vivo* conditions and could be used a sensitive biomarker for genotoxicity in earthworms.⁶⁷ Another earthworm, *Aporrectodea longa* (Ude), when exposed to soil samples spiked with B[a]P and/or lindane demonstrated genotoxicity in the intestinal cells to be more sensitive to the effect of the toxicants than the crop/gizzard cells.⁶⁹

Fourie *et al.*²¹⁶ used five earthworm species (*Amyntas diffringens*, *Aporrectodea caliginosa*, *Dendrodrilus rubidus*, *Eisenia foetida* and *Microchaetus benhami*) to study genotoxicity of sublethal concentrations of cadmium sulfate, with significant DNA damage being detected in *E. foetida* followed by *D. rubidus* and *A. caliginosa*. The study showed the difference in sensitivity of species present in an environment and its influence on the genotoxicity risk assessment. Hence, for environmental biomonitoring, specific species have to be kept in mind to reduce false-negative results.

1.4.1.5 The Comet Assay in *Drosophila*

The simple genetics and developmental biology of *Drosophila melanogaster* has made it the most widely used insect model and has been recommended as an alternate animal model by the European Centre for the Validation of Alternative Methods.²¹⁷ Recently, *Drosophila* has evolved into a model organism in toxicological studies.^{218,219} *D. melanogaster* has also been used as an *in vivo* model for assessment of genotoxicity using the Comet assay^{70–72,206} (Table 1.1). Neuroblast cells of third instar larvae, DNA repair deficient in nucleotide excision repair (mus201) and a mechanism of damage bypass (mus308), have been used for mechanistic studies.²⁰⁶

Third instar larvae of *D. melanogaster* (Oregon R+) were validated for genotoxicity assessment using a modified Comet assay.^{70,71} Since the cells of *Drosophila* are smaller than mammalian cells, modifications in the Comet assay were done, *e.g.* higher concentration of agarose (for the smaller size of *Drosophila* cells), removal of DMSO from lysing solution (DMSO is toxic to the cells) and lower electrophoresis time (for improved performance of the assay). This modified protocol was validated in gut and brain cells using

well-known alkylating agents, *i.e.* ethyl methanesulfonate (EMS), methyl methanesulfonate (MMS), N-ethyl-N-nitrosourea (ENU) and cyclophosphamide (CP) that were mixed in standard *Drosophila* diet and produced a significant dose-dependent response.^{70,71} Cypermethrin, a synthetic pyrethroid, even at low concentrations (at 0.002 ppm) and leachates of industrial waste produced significant dose-dependent increases in DNA damage in the brain ganglia and anterior mid gut of *D. melanogaster*.^{71,72} Results from the Comet assay have also shown a direct correlation between the concentrations of cisplatin adducts and DNA damage in somatic cells of *D. melanogaster*.⁷³

In vitro studies using *Drosophila* S2 cells demonstrated that the ectopically expressed DNA glycosylases (dOgg1 and RpS3) reduced the oxidised guanosine (8-OxoG), but contributed to increased DNA degradation due to one of the constituents of the DNA repair system.²²⁰

The studies in *Drosophila* have shown it to be a good alternative to animal models for the assessment of *in vivo* genotoxicity of chemicals using the Comet assay.

1.4.1.6 The Comet Assay in Other Invertebrates

Nereis virens, a polychaete, plays an important role in the distribution of pollutants in sediments due to their unique property of bioturbation. These worms are similar to earthworms in soil and can be used for genotoxicity assessment of sediments. Intracoelomic injection of B[a]P was given to the worms and the Comet assay was conducted on coelomocytes.²²¹ *Nereis* species was, however, not found to be suitable for assessing PAH genotoxicity probably due to its lack of metabolic capability to convert B[a]P to its toxic metabolite.²²¹

DNA damage was assessed in neuroblast cells of brains of 1st instars of grasshoppers (*Chorthippus brunneus*) exposed to various doses of zinc from a polluted site to understand the mechanism of toxicity in insects due to industrial pollutants.²²²

The estuarine grass shrimp, *Palaemonetes pugio*, exposed to coal-combustion residues from coal-fired electrical generation, were studied for DNA damage using the Comet assay. Chronic exposure caused DNA damage in hepatopancreatic cells of adult shrimps as compared to the reference shrimp.⁷⁷ The Comet assay in planarians is an important test for environmental monitoring studies since these are simple organisms with high sensitivity, low cost and a high proliferative rate.²²³ The genotoxic potential of water from Diluvio's Basin was evaluated in planarians, where an increase in pollutants towards the basin led to an increase in the DNA damage in these species.²²³ A significant increase of primary DNA damage was observed in planarian cells due to a Norflurazon, a bleaching herbicide²²⁴ and copper sulfate,²²⁵ when compared to the control animals.

These studies have also shown the use of the Comet assay in biomonitoring diverse environmental conditions utilising sentinel species.

1.5 Higher Animals

1.5.1 Vertebrates

Studies of vertebrate species where the Comet assay is used include fishes, amphibians, birds and mammals. Cells (blood, gills, kidneys and livers) of different fishes, tadpoles and adult frogs, as well as rodents have been used for assessing *in vivo* and *in vitro* genotoxicity of chemicals, and human biomonitoring has also been carried out employing the Comet assay (Table 1.1).

1.5.1.1 The Comet Assay in Fishes

Various fishes (freshwater and marine) have been used for environmental biomonitoring, as they are endemic organisms, which serve as sentinel species for a particular aquatic region to the adverse effects of chemicals and environmental conditions. The Comet assay has found wide application as a simple and sensitive method for evaluating *in vivo* as well as *in vitro* DNA damage in different tissues (gills, liver, blood) of fishes exposed to various xenobiotics in the aquatic environment (Table 1.1).

Environmental biomonitoring to assess the water quality in rivers has been carried out in hepatocytes of chub,⁷⁹ erythrocytes of mullet (*Mugil* sp.), sea catfish (*Netuma* sp.^{81,82}), bullheads (*Ameiurus nebulosus*) and carp (*Cyprinus carpio*^{90,226}). The basal level of DNA damage has been shown to be influenced by various factors, such as the temperature of water in erythrocytes of mullet and sea catfish,^{81,82} age and gender in dab (*Limanda limanda*⁴³), and exhaustive exercise in chub.⁸⁰ Therefore, these factors should be accounted for during environmental biomonitoring studies. The sensitivity of the assay may be affected by high intraindividual variability.⁴³ The protocol and experimental conditions used for the Comet assay for monitoring marine ecosystems may lead to differences in the results obtained.⁹² The use of chemical and mechanical procedures to obtain cell suspensions may also lead to DNA damage.²²⁷ Anesthesia did not contribute towards DNA damage *in vivo* in methyl methanesulfonate (MMS) treated fishes and the anesthetic benzocaine did not alter the DNA damage in erythrocytes after *in vitro* exposure to MMS or H₂O₂.²²⁸ Hence keeping in mind animal welfare, multi sampling in the same fish can be conducted.

In vitro studies on fish hepatocytes,⁹⁹ primary hepatocytes and gill cells¹⁰³ as well as established cell lines (with metabolic competence²²⁹) using the Comet assay have also been conducted to assess the genotoxicity of chemicals in water samples. The antioxidant potential of indolinic and quinolinic nitroxide radicals,¹⁰⁰ tannins¹⁰¹ and low concentrations (<10 µM) of diaryl tellurides and ebselen – an organoselenium compound¹⁰² – in oxidative DNA damage has been studied in nucleated trout (*Oncorhynchus mykiss*) erythrocytes for use of these compounds in biological systems. Kammann *et al.*⁹⁸ demonstrated the Comet assay in isolated leukocytes of carp as an *in vitro* model for evaluating genotoxicity of marine sediment extracts and increased sensitivity of the

method with use of the DNA repair inhibitor, 1-beta-D-arabinofuranosyl-cytosine (ara C). The Comet assay with fish cell lines may be a suitable tool for *in vitro* screening of environmental genotoxicity, however, the metabolising capabilities of the cell line need to be taken into account.

Cryopreservation has been shown to induce DNA-strand breaks in spermatozoa of trout,^{93,230} sea bass (*Dicentrarchus labrax*²³¹) and gilthead sea bream (*Sparus aurata*²³⁰). The DNA damage was prevented by the addition of cryopreservants such as BSA and dimethyl sulfoxide.²³¹ These studies have demonstrated the sperm Comet assay as a useful model in determining the DNA integrity in frozen samples for commercially cultured species.

The above studies have shown the usefulness of the Comet assay in fishes as a model for monitoring genotoxicity of aquatic habitats.

1.5.1.2 The Comet Assay in Amphibians

The Comet assay in amphibians has been carried out at adult and larval stages for ecogenotoxicity of aquatic environments and studies since 1999 have been well reviewed by Cotelle and Ferard.²⁰³ The animals chosen for the Comet assay act as sensitive bioindicators of aquatic and agricultural ecosystems (Table 1.1). The animals were either collected from the site (*in situ*) or exposed to chemicals under laboratory/natural conditions.

Erythrocytes from tadpoles of two species *Rana clamitans* and *Rana pipiens* have been used for the assessment of genotoxicity of water bodies as *in situ* sentinel organisms for environmental biomonitoring.¹¹⁵ *R. clamitans* tadpoles collected from agricultural regions showed significantly higher ($P < 0.001$) DNA damage than tadpoles collected from sites of little or no agriculture. Similarly *R. pipiens* tadpoles collected from industrial sites showed significantly higher ($P < 0.001$) DNA-strand breaks than samples from agricultural areas. The higher levels of DNA damage may be due to the pesticides used in the agricultural region. Variation in DNA damage due to sampling time¹¹⁵ and during various metamorphosis states²³² was also observed. Hence, for biomonitoring environmental genotoxicity using the Comet assay, pooling of early tadpole phases could be helpful. Studies have also been conducted on caged tadpoles in areas where the indigenous population is not present, due to ecological imbalance from pollution. *Rana clamitans* and the American toad (*Bufo americanus*) tadpoles were caged at the polluted reference site and demonstrated significant ($P < 0.05$) increases in DNA damage, relative to control tadpoles in the laboratory.²³³ These results demonstrated that caged tadpoles could be used for monitoring genotoxicity of water habitats that do not support the survival of tadpoles, e.g. large lakes and aquatic areas near high industrial activity.

Huang *et al.*¹¹⁰ have shown the genotoxicity of petrochemicals in liver and erythrocytes of toad *Bufo raddeis*. DNA damage was found to be positively correlated to the concentration of petrochemicals in liver, pointing to the fact that liver is the site for metabolism and may be a good marker for studying genotoxicity of compounds that require metabolic activation. The effect of

polyploidy on bleomycin-induced DNA damage and repair in *X. laevis* (pseudotetraploid) and *Xenopus tropicalis* (diploid) was studied using the Comet assay.¹¹¹ The *X. tropicalis* was more sensitive with a lower capacity for repair than *X. laevis*, showing that polyploidy protects DNA damage and allows rapid repair, and hence these species may be used as a good model for DNA damage and repair studies.

1.5.1.3 The Comet Assay in Birds

There are few studies involving the Comet assay in birds (Table 1.1). Genetic damage due to a mining accident involving heavy metals has been reported in free-living, nestling white storks (*Ciconia ciconia*) and black kites (*Milvus migrans*) from southwestern Spain,^{117–120} however, species-specific and intra-species differences were observed. Faullimel *et al.*¹²³ showed that the neutral Comet assay could be used to study the impact of freezing and thawing on DNA integrity in breast fillets and liver cells of frozen chicken. Frankic *et al.*¹²² reported that T-2 toxin and deoxynivalenol (DON) induced DNA fragmentation in chicken spleen leukocytes that was abrogated by dietary nucleotides. Kotlowska *et al.*¹²¹ have demonstrated increased DNA fragmentation in turkey sperm after 48 h of liquid storage which might be helpful in evaluating the DNA integrity for artificial insemination.

1.5.1.4 The Comet Assay in Rodents

Mice and rats have been widely used as animal models for the assessment of *in vivo* genotoxicity of chemicals using the Comet assay (Table 1.1). The *in vivo* Comet assay has been accepted by the UK Committee on Mutagenicity Testing of Chemicals in Food, Consumer Products and Environment¹⁰ as a test for assessing DNA damage, and is recommended for follow-up testing of positive *in vitro* findings. A positive result in the *in vivo* Comet assay assumes significance if mutagenic potential of a chemical has already been demonstrated *in vitro*. Within a battery of tests, the Comet assay finds a place as a supplemental *in vivo* test that has been accepted by international guidelines.²³⁴ There are specific guidelines for the performance of the Comet assay *in vivo* for reliable results.^{235–237}

Multiple organs of mouse/rat including brain, blood, kidney, lungs, liver, bone marrow have been utilised for the comprehensive understanding of the systemic genotoxicity of chemicals.^{133,134,238,239} The most important advantage of the use of Comet assay is that DNA damage in any organ can be evaluated without the need for mitotic activity and DNA damage in target as well as nontarget organs can also be seen.²³⁹ Comprehensive data on chemicals representing different classes, *e.g.* PAHs, alkylating compounds, nitroso compounds, food additives, *etc.* that caused DNA-strand breaks in various organs of mice was compiled by Sasaki *et al.*^{239,240} The mouse or rat organs exhibiting

increased levels of DNA damage were not necessarily the target organs for carcinogenicity. Therefore, for the prediction of carcinogenicity of a chemical, organ-specific genotoxicity was necessary but not sufficient.²⁴⁰ The Comet assay can be used as an *in vivo* test apart from the cytogenetic assays in haematopoietic cells and also for those compounds that have poor systemic bioavailability.

Different routes of exposure in rodents have been used, *e.g.* intraperitoneal,^{131,133} oral^{241,242} and inhalation^{130,243} to study the genotoxicity of different chemicals. The route of exposure is an important determinant of the genotoxicity of a chemical due to its mode of action.¹³⁴ The *in vivo* Comet assay helps in hazard identification and assessment of dose–response relationships as well as the mechanistic understanding of a substance’s mode of action. Besides being used for testing the genotoxicity of chemicals in laboratory-reared animals, the Comet assay in wild mice can be used as a valuable test in pollution monitoring and environmental conservation.²⁴⁴

The *in vivo* Comet assay in rodents is an important test model for genotoxicity studies, since many rodent carcinogens are also human carcinogens, and hence this model not only provides an insight into the genotoxicity of human carcinogens but is also suited for studying their underlying mechanisms.

1.5.1.5 The Comet Assay in Humans

The Comet assay is a valuable method for detection of occupational and environmental exposures to genotoxicants in humans and can be used as a tool in risk assessment for hazard characterisation^{6,8,245,246} (Table 1.1). The DNA-damage assessed by the Comet assay gives an indication of recent exposure and at an early stage where it could also undergo repair²⁴⁷ and thus it provides an opportunity for intervention strategies to be implemented in a timely manner. The assay can be conducted in the same population after removal of genotoxicant/dietary intervention to detect the extent of reduction in DNA damage. The assay is a noninvasive technique compared to other DNA-damage techniques (chromosomal aberrations, micronucleus), which require larger samples (~2–3 ml) as well as a proliferating cell population (or cell culture). Human biomonitoring using the Comet assay is advantageous since it is rapid, cost effective, with easy compilation of data and concordance with cytogenetic assays.²⁴⁸

The assay has been widely used in studying DNA damage and repair in healthy individuals,^{3,194,249,250} in clinical studies^{31,251,252} as well as in dietary intervention studies,^{155,158,253–255} and in monitoring the risk of DNA damage resulting from occupational,^{161,256–258} environmental,^{187,259} oxidative DNA damage,^{177,260} exposures or lifestyle.^{185,261} White blood cells or lymphocytes are the most frequently used cell type for the Comet assay in human biomonitoring studies.^{248,262,263} However, other cells have also been used, *e.g.* buccal cells,²⁶⁴ nasal,²⁶⁵ sperm,^{191,266–268} epithelial^{269–271} and placental cells.²⁷²

The Comet assay has been used as a test to predict the risk for development of diseases (renal cell carcinoma, cancers of the bladder, oesophagus and lung) due to susceptibility of the individual to DNA damage.^{149,273–275} The *in vitro* Comet assay is proposed as an alternative to cytogenetic assays in early genotoxicity/photogenotoxicity screening of drug candidates²⁷⁶ as well for neurotoxicity. Certain factors like age, diet, lifestyle (alcohol and smoking) as well as diseases have been shown to influence the Comet assay parameters and for interpretation of responses these factors need to be accounted for during monitoring human genotoxicity.^{277,278}

Human biomonitoring studies using the Comet assay provide an efficient tool for measuring human exposure to genotoxicants, thus helping in risk assessment and hazard identification.

1.6 The Specificity, Sensitivity and Limitations of the Comet Assay

The Comet assay has found worldwide acceptance for detecting DNA damage and repair in prokaryotic and eukaryotic cells. However, there are issues relating to the specificity, sensitivity and limitations of the assay that need to be addressed by genetic toxicologists before it gets accepted in the regulatory framework including interlaboratory validation of *in vitro* and *in vivo* Comet assay.

The variability in the results of the Comet assay is largely due to its sensitivity and minor differences in the conditions of various laboratories as well as the effect of confounding factors in human studies (lifestyle, age, diet, inter-individual and seasonal variation). Prospective cohort studies have not been conducted to find the predictive value of the Comet assay in human biomonitoring, further limiting its application.⁸ Cell to cell, gel to gel, culture to culture, animal to animal variability as well as use of various image-analysis systems or visual scoring²⁷⁹ and use of different Comet parameters, *e.g.* Olive tail moment and tail (%) DNA, are the other factors contributing to inter-laboratory differences in the results.

The limitation of the Comet assay is that it only detects DNA damage in the form of strand breaks. The alkaline (pH > 13) version of the assay assesses direct DNA damage or alkali-labile sites, while specific classes of DNA damage including base oxidation DNA adduct formation cannot be measured. The specific and sensitive detection of these lesions requires the use of lesion-specific enzymes.³ These enzymes are bacterial glycosylase/endonuclease enzymes, which recognise a particular type of damage and convert it into a break that can then be measured in the Comet assay. Hence, broad classes of oxidative DNA damage, alkylations, and ultraviolet light-induced photoproducts can be detected as an increased amount of DNA in the tail.⁸ Oxidised pyrimidines are detected with use of endonuclease III, while oxidised purines are detected with formamidopyrimidine DNA glycosylase (FPG). Modifications have been made in the protocol to specifically detect double-strand breaks (neutral Comet

assay²⁸⁰), single-strand breaks (at pH 12.1,²⁸¹), DNA crosslinking (decrease in DNA migration due to crosslinks²⁸⁰) and apoptosis.²⁸⁰ The neutral Comet assay also helps to distinguish apoptosis from necrosis as evidenced by the increased Comet score in apoptotic cells and the almost zero Comet score in necrotic cells.²⁸² An adaptation of the Comet assay was also developed that enables the discrimination of viable, apoptotic and necrotic single cells.²⁸³ Use of proteinase-K specifically removes DNA–protein crosslinking, leading to increased migration but would not affect the DNA–DNA crosslinking, thereby indicating a specific type of lesion.²⁸⁰

Tail (%) DNA and Olive tail moment give a good correlation in genotoxicity studies and since most studies have reported these Comet parameters, it has been recommended that both these parameters should be applied for routine use. Since the OTM is reported as arbitrary units and different image-analysis systems give different values, tail (%) DNA is considered a better parameter.²⁸⁵

It is therefore required that the *in vitro* and *in vivo* testing be conducted according to the Comet assay guidelines, and appropriately designed multi-laboratory international validation studies be carried out.

Guidelines for the *in vitro* as well as *in vivo* Comet assay have been formulated.^{235,236} Recently, issues relating to study design and data analysis in the Comet assay were discussed by the International Workgroup on Genotoxicity Testing (IWGT), where particular attention was given to the alkaline version (pH > 13) of the *in vivo* Comet assay and recommendations were made for a standardised protocol, which would be acceptable to international agencies.²³⁷ It was decided that a single dose should be replaced with multiple dosing to avoid misinterpretation of data, isolated cells or nuclei could be used for the studies, cytotoxicity should be tested in the cells to prevent mechanisms of apoptosis/necrosis from interfering with the results, and scoring of comets could be carried out both manually as well as with image-analysis systems. Consensus was also reached on the need for an international validation study to stringently evaluate the reliability and accuracy of the *in vivo* Comet assay (as well as *in vitro* versions). These recommendations are also aimed at reducing the variability arising in interlaboratory studies.

Since *in vivo* Comet assay has been accepted as the first tier screening assay for assessment of DNA damage in rodents by the Committee on Mutagenicity, UK,¹⁰ international validation studies are underway supported by the European Centre for Validation of Alternative Methods (ECVAM), Japanese Centre for Validation of Alternative Methods (JaCVAM), US Interagency Coordinating Committee on Validation of Alternative Methods (ICCVAM), US National Toxicology Program Interagency Centre for Evaluation of Alternative Toxicological Methods (NICEATM) and Japanese Environmental Mutagen Society.²³⁷

There has been only one multilaboratory validation study in the European countries that has been conducted to study the FPG sensitive sites and background level of base oxidation in DNA using the Comet assay, in human lymphocytes.²⁸⁴ It was found that half of the laboratories demonstrated a

dose–response effect. However, many laboratories have carried out their own validation studies for DNA damage to optimise their research work.⁸ Moller²⁶³ has critically evaluated the published Comet assay data on human biomonitoring studies using blood cells from 22 countries and has established reference values for DNA damage. The large number of biomonitoring studies has indicated that the Comet assay is a useful tool for detecting exposure and its validation status as a biomarker in biomonitoring is dependent on its performance in cohort studies.⁸

1.7 Conclusions

The Comet assay is now well established and its versatility has imparted a sensitive tool to the toxicologists for assessing DNA damage. This has been demonstrated with its wide applications in assessing genotoxicity in plant and animal models, both aquatic as well as terrestrial, in a variety of organisms, tissues and cell types. *In vitro*, *in vivo*, *in situ* and biomonitoring studies using the Comet assay have proved it to be a “Rossetta Stone” in the garden of genetic toxicology.

Acknowledgments

The authors wish to thank the Council of Scientific and Industrial Research (CSIR), New Delhi, India for funding through the Networked Projects (CMM0018 and NWP34) as well as the support from the UK-India Education and Research Initiative (UKIERI).

References

1. O. Ostling and K. J. Johanson, Microelectrophoretic study of radiation-induced DNA damages in individual mammalian cells, *Biochem. Biophys. Res. Commun.*, 1984, **123**(1), 291–8.
2. N. P. Singh, M. T. McCoy, R. R. Tice and E. L. Schneider, A simple technique for quantitation of low levels of DNA damage in individual cells, *Exp. Cell. Res.*, 1988, **175**(1), 184–91.
3. A. R. Collins, The Comet assay for DNA damage and repair principles, applications, and limitations, *Mol. Biotechnol.*, 2004, **26**, 249–60.
4. G. Speit and A. Hartmann, The Comet assay: a sensitive genotoxicity test for the detection of DNA damage, *Methods Mol. Biol.*, 2005, **291**, 85–95.
5. P. L. Olive and J. P. Banath, The Comet assay: a method to measure DNA damage in individual cells, *Nature Protocols*, 2006, **1**(1), 23–9.
6. P. Moller, Genotoxicity of environmental agents assessed by the alkaline Comet assay, *Basic Clin. Pharmacol. Toxicol.*, 2005, **96**, 1–42.
7. F. Kassie, W. Parzefall and S. Knasmüller, Single-cell gel electrophoresis assay: a new technique for human biomonitoring studies, *Mutat. Res.*, 2000, **463**(1), 13–31.

8. P. Moller, The alkaline Comet assay: towards validation in biomonitoring of DNA damaging exposures, *Basic Clin. Pharmacol. Toxicol.*, 2006a, **98**(4), 336–45.
9. C. Gedik, S. Ewen and A. Collins, Single-cell gel electrophoresis applied to the analysis of UV-C damage and its repair in human cells, *Int. J. Radiat. Biol.*, 1992, **62**, 313–20.
10. Committee on Mutagenicity of Chemicals in Food, Consumer Products and the Environment, COM guidance on a strategy for testing of chemicals for mutagenicity, United Kingdom, December 2000.
11. N. P. Singh, R. E. Stephens, H. Singh and H. Lai, Visual quantification of DNA double-strand breaks in bacteria, *Mutat. Res.*, 1999, **429**, 159–68.
12. J. Maluszynska and J. Juchimiuk, Plant genotoxicity: a molecular cytogenetic approach in plant bioassays, *Arh. Hig. Rada. Toksikol.*, 2005, **56**(2), 177–84.
13. W. F. Grant, Higher plant assays for the detection of chromosomal aberrations and gene mutations—a brief historical background on their use for screening and monitoring environmental chemicals, *Mutat. Res.*, 1999, **426**, 107–12.
14. T. Ma, Z. Xu, C. Xu, H. McConnell, E. V. Rabago and G. A. Arreola, *et al.*, The improved *Allium/Vicia* root tip micronucleus assay for clastogenicity of environmental pollutants, *Mutat. Res.*, 1995, **334**, 185–95.
15. P. Banerjee, S. N. Talapatra, N. Mandal, G. Sundaram, A. Mukhopadhyay and D. Chattopadhyay, *et al.*, Genotoxicity study with special reference to DNA damage by Comet assay in fission yeast, *Schizosaccharomyces pombe* exposed to drinking water, *Food Chem. Toxicol.*, 2008, **46**(1), 402–7.
16. A. Hahn and B. Hock, Assessment of DNA damage in filamentous fungi by single-cell gel electrophoresis, Comet assay, *Environ. Toxicol. Chem.*, 1999, **18**, 1421–24.
17. M. Erbes, A. Wessler, U. Obst and A. Wild, Detection of primary DNA damage in *Chlamydomonas reinhardtii* by means of modified microgel electrophoresis, *Environ. Mol. Mutagen.*, 1997, **30**(4), 448–58.
18. K. Aoyama, K. Iwahori and N. Miyata, Application of *Euglena gracilis* cells to Comet assay: evaluation of DNA damage and repair, *Mutat. Res.*, 2003, **538**(1–2), 155–62.
19. M. P. Sastre, M. Vernet and S. Steinert, Single-cell gel/Comet assay applied to the analysis of UV radiation-induced DNA damage in *Rhodomonas* sp. (Cryptophyta), *Photochem. Photobiol.*, 2001, **74**(1), 55–60.
20. K. J. Angelis, M. McGuffie, M. Menke and I. Schubert, Adaptation to alkylation damage in DNA measured by the Comet assay, *Environ. Mol. Mutagen.*, 2000, **36**(2), 146–50.
21. M. Menke, A. Meister and I. Schubert, N-Methyl-N-nitrosourea-induced DNA damage detected by the Comet assay in *Vicia faba* nuclei during all interphase stages is not restricted to chromatid aberration hot spots, *Mutagenesis*, 2000, **15**(6), 503–6.