

London Mathematical Society
Lecture Note Series 190

Polynomial Invariants of Finite Groups

D. J. Benson

CAMBRIDGE

*London
Mathematical
Society*



CAMBRIDGE

more information – www.cambridge.org/9780521458863

LONDON MATHEMATICAL SOCIETY LECTURE NOTE SERIES

Managing Editor: Professor J.W.S. Cassels, Department of Pure Mathematics and Mathematical Statistics, University of Cambridge, 16 Mill Lane, Cambridge CB2 1SB, England

The books in the series listed below are available from booksellers, or, in case of difficulty, from Cambridge University Press.

- 34 Representation theory of Lie groups, M.F. ATIYAH *et al*
- 36 Homological group theory, C.T.C. WALL (ed)
- 39 Affine sets and affine groups, D.G. NORTHCOTT
- 46 p-adic analysis: a short course on recent work, N. KOBLITZ
- 49 Finite geometries and designs, P. CAMERON, J.W.P. HIRSCHFELD & D.R. HUGHES (eds)
- 50 Commutator calculus and groups of homotopy classes, H.J. BAUES
- 57 Techniques of geometric topology, R.A. FENN
- 59 Applicable differential geometry, M. CRAMPIN & F.A.E. PIRANI
- 66 Several complex variables and complex manifolds II, M.J. FIELD
- 69 Representation theory, I.M. GELFAND *et al*
- 74 Symmetric designs: an algebraic approach, E.S. LANDER
- 76 Spectral theory of linear differential operators and comparison algebras, H.O. CORDES
- 77 Isolated singular points on complete intersections, E.J.N. LOOIJENGA
- 79 Probability, statistics and analysis, J.F.C. KINGMAN & G.E.H. REUTER (eds)
- 80 Introduction to the representation theory of compact and locally compact groups, A. ROBERT
- 81 Skew fields, P.K. DRAXL
- 82 Surveys in combinatorics, E.K. LLOYD (ed)
- 83 Homogeneous structures on Riemannian manifolds, F. TRICERRI & L. VANHECKE
- 86 Topological topics, I.M. JAMES (ed)
- 87 Surveys in set theory, A.R.D. MATHIAS (ed)
- 88 FPF ring theory, C. FAITH & S. PAGE
- 89 An F-space sampler, N.J. KALTON, N.T. PECK & J.W. ROBERTS
- 90 Polytopes and symmetry, S.A. ROBERTSON
- 91 Classgroups of group rings, M.J. TAYLOR
- 92 Representation of rings over skew fields, A.H. SCHOFIELD
- 93 Aspects of topology, I.M. JAMES & E.H. KRONHEIMER (eds)
- 94 Representations of general linear groups, G.D. JAMES
- 95 Low-dimensional topology 1982, R.A. FENN (ed)
- 96 Diophantine equations over function fields, R.C. MASON
- 97 Varieties of constructive mathematics, D.S. BRIDGES & F. RICHMAN
- 98 Localization in Noetherian rings, A.V. JATEGAONKAR
- 99 Methods of differential geometry in algebraic topology, M. KAROUBI & C. LERUSTE
- 100 Stopping time techniques for analysts and probabilists, L. EGGHE
- 101 Groups and geometry, ROGER C. LYNDON
- 103 Surveys in combinatorics 1985, I. ANDERSON (ed)
- 104 Elliptic structures on 3-manifolds, C.B. THOMAS
- 105 A local spectral theory for closed operators, I. ERDELYI & WANG SHENGWANG
- 106 Syzygies, E.G. EVANS & P. GRIFFITH
- 107 Compactification of Siegel moduli schemes, C-L. CHAI
- 108 Some topics in graph theory, H.P. YAP
- 109 Diophantine analysis, J. LOXTON & A. VAN DER POORTEN (eds)
- 110 An introduction to surreal numbers, H. GONSHOR
- 111 Analytical and geometric aspects of hyperbolic space, D.B.A. EPSTEIN (ed)
- 113 Lectures on the asymptotic theory of ideals, D. REES
- 114 Lectures on Bochner-Riesz means, K.M. DAVIS & Y-C. CHANG
- 115 An introduction to independence for analysts, H.G. DALES & W.H. WOODIN
- 116 Representations of algebras, P.J. WEBB (ed)
- 117 Homotopy theory, E. REES & J.D.S. JONES (eds)
- 118 Skew linear groups, M. SHIRVANI & B. WEHRFRITZ
- 119 Triangulated categories in the representation theory of finite-dimensional algebras, D. HAPPEL
- 121 Proceedings of *Groups - St Andrews 1985*, E. ROBERTSON & C. CAMPBELL (eds)
- 122 Non-classical continuum mechanics, R.J. KNOPS & A.A. LACEY (eds)
- 124 Lie groupoids and Lie algebroids in differential geometry, K. MACKENZIE
- 125 Commutator theory for congruence modular varieties, R. FREESE & R. MCKENZIE
- 126 Van der Corput's method of exponential sums, S.W. GRAHAM & G. KOLESNIK
- 127 New directions in dynamical systems, T.J. BEDFORD & J.W. SWIFT (eds)

- 128 Descriptive set theory and the structure of sets of uniqueness, A.S. KECHRIS & A. LOUVEAU
- 129 The subgroup structure of the finite classical groups, P.B. KLEIDMAN & M.W. LIEBECK
- 130 Model theory and modules, M. PREST
- 131 Algebraic, extremal & metric combinatorics, M-M. DEZA, P. FRANKL & I.G. ROSENBERG (eds)
- 132 Whitehead groups of finite groups, ROBERT OLIVER
- 133 Linear algebraic monoids, MOHAN S. PUTCHA
- 134 Number theory and dynamical systems, M. DODSON & J. VICKERS (eds)
- 135 Operator algebras and applications, 1, D. EVANS & M. TAKESAKI (eds)
- 136 Operator algebras and applications, 2, D. EVANS & M. TAKESAKI (eds)
- 137 Analysis at Urbana, I, E. BERKSON, T. PECK, & J. UHL (eds)
- 138 Analysis at Urbana, II, E. BERKSON, T. PECK, & J. UHL (eds)
- 139 Advances in homotopy theory, S. SALAMON, B. STEER & W. SUTHERLAND (eds)
- 140 Geometric aspects of Banach spaces, E.M. PEINADOR and A. RODES (eds)
- 141 Surveys in combinatorics 1989, J. SIEMONS (ed)
- 142 The geometry of jet bundles, D.J. SAUNDERS
- 143 The ergodic theory of discrete groups, PETER J. NICHOLLS
- 144 Introduction to uniform spaces, I.M. JAMES
- 145 Homological questions in local algebra, JAN R. STROOKER
- 146 Cohen-Macaulay modules over Cohen-Macaulay rings, Y. YOSHINO
- 147 Continuous and discrete modules, S.H. MOHAMED & B.J. MÜLLER
- 148 Helices and vector bundles, A.N. RUDAKOV et al
- 149 Solitons, nonlinear evolution equations and inverse scattering, M.J. ABLOWITZ & P.A. CLARKSON
- 150 Geometry of low-dimensional manifolds 1, S. DONALDSON & C.B. THOMAS (eds)
- 151 Geometry of low-dimensional manifolds 2, S. DONALDSON & C.B. THOMAS (eds)
- 152 Oligomorphic permutation groups, P. CAMERON
- 153 L-functions and arithmetic, J. COATES & M.J. TAYLOR (eds)
- 154 Number theory and cryptography, J. LOXTON (ed)
- 155 Classification theories of polarized varieties, TAKAO FUJITA
- 156 Twistors in mathematics and physics, T.N. BAILEY & R.J. BASTON (eds)
- 157 Analytic pro- p groups, J.D. DIXON, M.P.F. DU SAUTOY, A. MANN & D. SEGAL
- 158 Geometry of Banach spaces, P.F.X. MÜLLER & W. SCHACHERMAYER (eds)
- 159 Groups St Andrews 1989 volume 1, C.M. CAMPBELL & E.F. ROBERTSON (eds)
- 160 Groups St Andrews 1989 volume 2, C.M. CAMPBELL & E.F. ROBERTSON (eds)
- 161 Lectures on block theory, BURKHARD KÜLSHAMMER
- 162 Harmonic analysis and representation theory for groups acting on homogeneous trees, A. FIGA-TALAMANCA & C. NEBBIA
- 163 Topics in varieties of group representations, S.M. VOVSI
- 164 Quasi-symmetric designs, M.S. SHRIKANDE & S.S. SANE
- 165 Groups, combinatorics & geometry, M.W. LIEBECK & J. SAXL (eds)
- 166 Surveys in combinatorics, 1991, A.D. KEEDWELL (ed)
- 167 Stochastic analysis, M.T. BARLOW & N.H. BINGHAM (eds)
- 168 Representations of algebras, H. TACHIKAWA & S. BRENNER (eds)
- 169 Boolean function complexity, M.S. PATERSON (ed)
- 170 Manifolds with singularities and the Adams-Novikov spectral sequence, B. BOTVINNIK
- 172 Algebraic varieties, GEORGE R. KEMPF
- 173 Discrete groups and geometry, W.J. HARVEY & C. MACLACHLAN (eds)
- 174 Lectures on mechanics, J.E. MARSDEN
- 175 Adams memorial symposium on algebraic topology 1, N. RAY & G. WALKER (eds)
- 176 Adams memorial symposium on algebraic topology 2, N. RAY & G. WALKER (eds)
- 177 Applications of categories in computer science, M.P. FOURMAN, P.T. JOHNSTONE, & A.M. PITTS (eds)
- 178 Lower K- and L-theory, A. RANICKI
- 179 Complex projective geometry, G. ELLINGSRUD, C. PESKINE, G. SACCHIERO & S.A. STRØMME (eds)
- 180 Lectures on ergodic theory and Pesin theory on compact manifolds, M. POLLICOTT
- 181 Geometric group theory I, G.A. NIBLO & M.A. ROLLER (eds)
- 182 Geometric group theory II, G.A. NIBLO & M.A. ROLLER (eds)
- 183 Shintani zeta functions, A. YUKIE
- 184 Arithmetical functions, W. SCHWARZ & J. SPILKER
- 185 Representations of solvable groups, O. MANZ & T.R. WOLF
- 186 Complexity: knots, colourings and counting, D.J.A. WELSH
- 187 Surveys in combinatorics, 1993, K. WALKER (ed)
- 190 Polynomial invariants of finite groups, D.J. BENSON

London Mathematical Society Lecture Note Series. 190

Polynomial Invariants of Finite Groups

D. J. Benson
University of Oxford

 **CAMBRIDGE**
UNIVERSITY PRESS

CAMBRIDGE UNIVERSITY PRESS

Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore,
São Paulo, Delhi, Dubai, Tokyo, Mexico City

Cambridge University Press
The Edinburgh Building, Cambridge CB2 8RU, UK

Published in the United States of America by
Cambridge University Press, New York

www.cambridge.org

Information on this title: www.cambridge.org/9780521458863

© Cambridge University Press 1993

This publication is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without the written
permission of Cambridge University Press.

First published 1993

A catalogue record for this publication is available from the British Library

Library of Congress cataloguing in publication data available

ISBN 978-0-521-45886-3 Paperback

Cambridge University Press has no responsibility for the persistence or
accuracy of URLs for external or third-party internet websites referred to in
this publication, and does not guarantee that any content on such websites is,
or will remain, accurate or appropriate. Information regarding prices, travel
timetables, and other factual information given in this work is correct at
the time of first printing but Cambridge University Press does not guarantee
the accuracy of such information thereafter.

Contents

Introduction	vii
1 Finite Generation of Invariants	1
1.1 The basic object of study	1
1.2 Noetherian rings and modules	3
1.3 Finite groups in arbitrary characteristic	5
1.4 Krull dimension and going up and down	5
1.5 Noether's bound in characteristic zero	9
1.6 Linearly reductive algebraic groups	10
2 Poincaré series	13
2.1 The Hilbert–Serre theorem	13
2.2 Noether normalization	15
2.3 Systems of parameters	18
2.4 Degree and ψ	20
2.5 Molien's theorem	21
2.6 Reflecting hyperplanes	23
3 Divisor Classes, Ramification and Hyperplanes	25
3.1 Divisors	25
3.2 Primes of height one	27
3.3 Duality	30
3.4 Reflexive modules	31
3.5 Divisor classes and unique factorization	32
3.6 The Picard group	33
3.7 The trace	34
3.8 Ramification	35
3.9 $\text{Cl}(K[V]^G)$	37
3.10 The different	39
3.11 The homological different	40
3.12 A ramification formula	42
3.13 The Carlisle–Kropholler conjecture	43

4	Homological Properties of Invariants	45
4.1	Minimal resolutions	45
4.2	Hilbert's syzygy theorem	47
4.3	Depth and Cohen–Macaulay rings	50
4.4	Homological characterization of depth	52
4.5	The canonical module and Gorenstein rings	55
4.6	Watanabe's theorem	59
5	Polynomial tensor exterior algebras	61
5.1	Motivation and first properties	61
5.2	A variation on Molien's theorem	62
5.3	The invariants are graded Gorenstein	63
5.4	The Jacobian	64
6	Polynomial rings and regular local rings	69
6.1	Regular local rings	69
6.2	Serre's converse to Hilbert's syzygy theorem	70
6.3	Uniqueness of factorization	73
6.4	Reflexive modules	74
7	Groups Generated by Pseudoreflections	79
7.1	Reflections and pseudoreflections	79
7.2	The Shephard–Todd theorem	82
7.3	A theorem of Solomon	86
8	Modular invariants	89
8.1	Dickson's theorem	89
8.2	The special linear group	91
8.3	Symplectic invariants	92
A	Examples over the complex numbers	99
B	Examples over finite fields	103
	Bibliography	109
	Index	116

Introduction

Invariant theory is a subject with a long history, a subject which in the words of Dieudonné and Carrell [32] “has already been pronounced dead several times, and like the Phoenix it has been again and again arising from the ashes.”

The starting point is a linear representation of a linear algebraic group on a vector space, which then induces an action on the ring of polynomial functions on the vector space, and one looks at the ring consisting of those polynomials which are invariant under the group action. The reason for restricting ones attention to linear algebraic group representations, or equivalently to Zariski closed subgroups of the general linear group on the vector space, is that the polynomial invariants of an arbitrary subgroup of the general linear group are the same as the invariants of its Zariski closure.

In the nineteenth century, attention focused on proving finite generation of the algebra of invariants by finding generators for the invariants in a number of concrete examples. One of the high points was the proof of finite generation for the invariants of $SL_2(\mathbb{C})$ acting on a symmetric power of the natural representation, by Gordan [39] (1868). The subject generated a language all its own, partly because of the influence of Sylvester, who was fond of inventing words to describe rather specialized concepts.

In the late nineteenth and early twentieth century, the work of David Hilbert and Emmy Noether in Göttingen clarified the subject considerably with the introduction of abstract algebraic machinery for addressing questions like finite generation, syzygies, and so on. This was the beginning of modern commutative algebra. Hilbert proved finite generation for the case of $GL_n(\mathbb{C})$ acting on a symmetric power of the natural representation [45] (1890), using the fact that a polynomial ring is Noetherian (Hilbert’s basis theorem).

Hilbert’s fourteenth problem, posed at the 1900 International Congress of Mathematicians in Paris, asks whether the invariants are always finitely generated. Well, actually, this is false. He was under the impression that L. Maurer had just proved this, and he goes on to ask a more complicated question which generalizes this. In fact, Maurer’s proof contained a mistake, and Nagata [65] (1959) found a counterexample to finite generation. An account of the Hilbert problems and the progress made on them up to the mid nineteen seventies is contained in the A.M.S. publication [6].

For reductive algebraic groups, however, finite generation does hold. This was first proved over the complex numbers by Weyl [110] (1926), using his “unitarian trick”, which amounts to proving that reductive groups are linearly reductive. In characteristic p , reductive groups are no longer necessarily linearly reductive (the only connected linearly reductive groups in characteristic p are the tori), but Mumford [64] conjectured and Haboush [41] (1975) proved that reductive groups are “geometrically reductive”, and Nagata [67] had already proved that this is enough to ensure finite generation for the invariants.

The question of when the ring of invariants of representation of a finite group over the complex numbers is a polynomial ring was solved by Shephard and Todd [95] (1954). Their theorem states that the ring of invariants is a polynomial ring if and only if the group is generated by elements fixing a hyperplane. Such elements are called pseudoreflections. Their proof involved classifying the finite groups generated by pseudoreflections, and is a tour de force of combinatorics and geometry. The classification was used to prove that groups generated by pseudoreflections have invariant rings that are polynomial rings, and this first implication together with Molien's theorem was then used in the proof of the converse. Later, Chevalley [26] (1955) provided a proof of the first implication for a real reflection group, which avoided classifying the groups. This proof involved the combinatorics of differential operators. Serre later showed how this proof could be adapted to the complex case. A more homological proof of this first implication was found by Larry Smith [96] (1985), and a proof of the converse which works in arbitrary characteristic appeared in Bourbaki [17] Chapter 5 §5, Exercise 7 (1981). The first implication is false in arbitrary characteristic.

The next interesting ring theoretic property of the ring of invariants of a reductive group is that it is Cohen–Macaulay. This was proved in the case of a finite group by Hochster and Eagon [47] (1971). Watanabe [107, 108] (1974) proved that in this case the ring of invariants is Gorenstein provided the group acts as matrices with determinant one. In the general reductive case, both these statements were proved by Hochster and Roberts [48] (1974).

The question of when unique factorization holds in rings of invariants was first attacked by Samuel [86] (1964). He developed the framework for defining the ideal class group of a Krull domain. A finitely generated Krull domain is the same thing as a normal domain (i.e., a Noetherian integrally closed domain). Using Samuel's ideas, Nakajima proved that the ring of invariants of a finite group is a unique factorization domain if and only if there are no non-trivial homomorphisms from the group to the multiplicative group of the field, taking every pseudoreflection to the identity element.

A lot of the ideas involved in invariant theory in the reductive case are already present in the finite case; and the latter has the advantage that one does not have to spend an inordinate amount of time setting up the machinery of algebraic groups before getting to the interesting theorems. For this reason, I decided to restrict the scope of this book to the case of finite group representations.

This book is based on a lecture course I gave at Oxford in the spring of 1991. My starting point for these lectures was the excellent survey article of Richard Stanley [102], which I strongly recommend to anyone wishing to get an overview of the subject. The influence of this article will be apparent in almost every part of this book. The main direction in which this book differs from that article, apart from length and amount of detail, is that I have tried to indicate how much is true in arbitrary characteristic, whereas Stanley restricts his discussion to characteristic

zero. Also, Stanley does not discuss divisor classes and unique factorization.

At Larry Smith's invitation, I spent two months in Göttingen in the summer of 1991, and he persuaded me that we should write a book together on invariant theory of finite groups. Differences in personal style prevented this from being realized as a joint project, but I would like to take this opportunity to thank him for discussions which had a considerable effect on the presentation of the material in Sections 7.2 and 8.1, and indeed for persuading me to embark on this project in the first place.

I would also like to thank Bill Crawley-Boevey for taking the trouble to make many constructive criticisms, and for suggesting the inclusion of the appendices; and David Tranah of Cambridge University Press for his usual endless supply of patience.

Chapter 1

Finite Generation of Invariants

1.1 The basic object of study

Let G be a finite group, K a field of coefficients and V a finite dimensional K -vector space on which G acts by linear substitutions (or equivalently a finitely generated KG -module). We write $K[V]$ for the ring of polynomial functions on V . In other words, if V has dimension n as a vector space, and x_1, \dots, x_n is a basis for the dual space $V^* = \text{Hom}_K(V, K)$, then

$$K[V] = K[x_1, \dots, x_n] = K \oplus V^* \oplus S^2(V^*) \oplus S^3(V^*) \oplus \dots$$

Here, $S^m(V^*)$ denotes the symmetric m th power of V^* , which consists of the homogeneous polynomials of degree m in x_1, \dots, x_n . Thus for example $S^2(V^*)$ has a basis consisting of the monomials $x_i x_j$ for the $\binom{n+1}{2}$ choices of i and j . In general, the dimension of $S^m(V^*)$ as a vector space is $\binom{n+m-1}{m} = \binom{n+m-1}{n-1}$. We regard $K[V]$ as a graded ring by putting each x_i in degree one. (If you are a topologist, then you may wish to double all the degrees.) The group G acts on $K[V]$ via $(gf)(v) = f(g^{-1}v)$, and the basic object of study of invariant theory is the set of all fixed points of this action, the **ring of invariants** $K[V]^G$.

Warning If K is a finite field \mathbb{F}_q , then we should not really regard $K[V]$ as a ring of functions on V , since for example x_i and x_i^q take the same value at all points of V . Rather, we regard $K[V]$ as a ring of functions on $\bar{K} \otimes_K V$ fixed by the Galois automorphisms $\text{Gal}(\bar{K}/K)$, where \bar{K} is an algebraic closure of K .

If $\chi : G \rightarrow K^\times$ is a 1-dimensional representation of G , the module of χ -relative invariants, denoted by $K[V]_\chi^G$ is defined by

$$K[V]_\chi^G = \{f \in K[V] \mid g(f) = \chi(g)f \forall g \in G\}.$$