

Phosphorus response and amino acid composition of
different green gram (*Vigna radiata* (L.) K.Wilczek)
cultivars and green gram rotation effects on maize
(*Zea mays* L.) growth in Myanmar



Department of Organic Agriculture and Agroecosystems Research in the Tropics
and Subtropics
University of Kassel
Germany

Phosphorus response and amino acid composition of different green
gram (*Vigna radiata* (L.) K.Wilczek) cultivars and green gram rotation
effects on maize (*Zea mays* L.) growth in Myanmar

Doctoral Dissertation
submitted for the degree of Doctor of Agricultural Sciences
of the Faculty of Ecological Agricultural Sciences
University of Kassel

Presented by
Myo Kywe
M.Agr.St. (Qld), Australia

Witzenhausen, 20th February, 2006

Bibliografische Information Der Deutschen Bibliothek

Die Deutsche Bibliothek verzeichnet diese Publikation in der Deutschen Nationalbibliografie; detaillierte bibliografische Daten sind im Internet über <http://dnb.ddb.de> abrufbar.

1. Aufl. - Göttingen : Cuvillier, 2006

Zugl.: Kassel, Univ., Diss., 2006

ISBN 3-86537-788-2

1. Referee: Prof. Dr. Andreas Bürkert

2. Co-referee: Prof. Dr. Maria Renate Finckh

Date of examination: 20th February, 2006

Gedruckt mit Unterstützung des Deutschen Akademischen Austauschdienstes

© CUVILLIER VERLAG, Göttingen 2006

Nonnenstieg 8, 37075 Göttingen

Telefon: 0551-54724-0

Telefax: 0551-54724-21

www.cuvillier.de

Alle Rechte vorbehalten. Ohne ausdrückliche Genehmigung des Verlages ist es nicht gestattet, das Buch oder Teile daraus auf fotomechanischem Weg (Fotokopie, Mikrokopie) zu vervielfältigen.

1. Auflage, 2006

Gedruckt auf säurefreiem Papier

ISBN 3-86537-788-2

TABLE OF CONTENTS

CONTENT	Pages
LIST OF TABLES	iii
LISTS OF FIGURES	v
LIST OF APPENDICES	vii
SUMMARY	x
ZUSAMMENFASSUNG	xii
 Chapter 1.....	 1
General introduction.....	1
1.1. Maize, green gram and rotation systems in Myanmar	1
1.2. Aims and hypotheses	4
1.3. References	5
Chapter 2.....	6
Literature review.....	6
2.1. Nutritional constraints in tropical soils of Myanmar.....	6
2.1.1. Availability and management of phosphorus in tropical soils.....	6
2.1.2. Nitrogen availability and management	7
2.1.3. Soil organic matter, soil pH.....	8
2.1.4. The role of legumes in a rotation	8
2.1.5. Farmyard manure management in tropical soils	10
2.2. The nutritional value of legume proteins	11
2.3. References	12
Chapter 3.....	20
Phosphorus response and amino acid composition of different green gram (<i>Vigna radiata</i> (L.) R. Wilczek) cultivars from Myanmar	20
3.1. Introduction	20
3.2. Materials and methods	21
3.2.1. Site conditions and experimental set up.....	21
3.2.2. Pot experiment.....	24
3.2.3. Data analysis	25
3.3. Results.....	25
3.3.1. Field experiments at Yezin and Nyaung Oo.....	25
3.3.2. Pot experiment.....	34
3.4. Discussion	35
3.5. Conclusions	36
3.6. References	37
Chapter 4.....	39
Green gram rotation effects on maize growth parameters and soil quality in Myanmar	39
4.1. Introduction	39
4.2. Materials and methods	40
4.2.1. Experimental details.....	40
4.2.2. Plant sampling and analysis	42
4.2.3. Measurements of mycorrhizae and nematodes	43
4.2.4. Data analysis	43
4.3. Results.....	43
4.3.1. Green gram.....	43
4.3.2. Maize.....	51
4.4. Discussion	59

4.4.1. Green gram.....	59
4.4.2. Maize.....	59
4.5. Conclusions	61
4.6. References	62
Chapter 5.....	66
General discussion.....	66
5.1. General findings	66
5.1.1. P application on green gram.....	66
5.1.2. Amino acid composition.....	66
5.1.3. Maize green gram rotation.....	67
5.1.4. Manure.....	67
5.1.5. Phosphorus application on maize.....	67
5.1.6. Legume cultivars and management.....	67
5.2. Final Recommendations.....	68
Appendices.....	69
Acknowledgements	92
Curriculum Vitae.....	94
Erklärung	95

LIST OF TABLE	Pages
Table 1.1. Land utilization in Myanmar in 2003-2004.....	1
Table 1.2. Area sown and production of various groups of crops in Myanmar, 2003-2004 (MOAI, 2004).....	3
Table 2.1. N and P contents of different animal manures	11
Table 3.1. List of improved cultivars and landraces from Myanmar of green gram tested for their growth response to phosphorus and for their amino acid composition in a field experiment in Myanmar, 2001.	22
Table 3.2. Combined analysis of variance (ANOVA) table for phosphorus (P) effects on total dry matter (TDM) and grain yield of different green gram cultivars grown in field experiments at Yezin and Nyaung Oo, Myanmar (2001-2002).....	27
Table 3.3. Effects of phosphorus (P) application at 0 and 15 kg P ha ⁻¹ on pest damage (%) and virus score in different green gram cultivars at Nyaung Oo, Myanmar, 2002.	30
Table 3.4. Effects of phosphorus (P) application at 0 and 15 kg P ha ⁻¹ on pest damage (%) and virus incidence (scored from 0 to 2) in different green gram cultivars at Yezin, Myanmar, 2003.....	31
Table 3.5. Effects of phosphorus (P) application at 0 and 15 kg P ha ⁻¹ on nitrogen and phosphorus concentration (mg g ⁻¹) in different green gram cultivars at Yezin, Myanmar, 2003.	32
Table 3.6. Amino acid concentrations (μmol essential amino acid g ⁻¹ TDM) in seeds of three improved and seven landraces green gram cultivars from Myanmar, 2001.	33
Table 3.7. Effects of phosphorus (P) application at an equivalent rate of 0 and 15 kg P ha ⁻¹ on total dry matter (TDM), grain yield and root length density (RLD) of four green gram cultivars grown in a pot experiment.	34
Table 3.8. Effects of phosphorus (P) application at an equivalent rate of 0 and 15 kg P ha ⁻¹ on shoot and seed nitrogen (N) concentration (mg g ⁻¹) of four green gram cultivars grown in a pot experiment.	35
Table 4.1. Cropping system in field experiments at Yezin, Myanmar	41
Table 4.2. Combined analysis of variance (ANOVA) table on manure (M) and phosphorus (P) application effects on green gram grown in a cropping systems (CS) experiment with continuous maize and maize in rotation with green gram at Yezin, Myanmar (five seasons from 2002-2004).....	46
Table 4.3. Effects of manure application (0 and 2000 kg ha ⁻¹), banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) and cultivar on nitrogen (N) and P concentrations of green gram shoots in 2003 at Yezin, Myanmar (2 nd rainy season).....	48
Table 4.4. Effects of manure application (0 and 2000 kg ha ⁻¹), banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) and cultivar on nitrogen (N) and P concentrations of green gram shoots in 2003 at Yezin, Myanmar (3 rd dry season). Means with different letters within columns are significantly different at P < 0.05.....	49
Table 4.5. Effects of manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on nitrogen (N) and P concentrations of green gram shoots in 2004 at Yezin, Myanmar (4 th rainy season).....	50
Table 4.6. Effects of manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on nitrogen (N) and P concentrations of green gram shoots in 2004 at Yezin, Myanmar (5 th dry	

season). Means with different letters within columns are significantly different at $P < 0.05$ 50

Table 4.7. Analysis of variance (ANOVA) table for the time dependent effects (five seasons from 2002-2004) of cropping system (continuous maize *versus* green gram-maize rotation), manure application (0 and 2000 kg ha⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha⁻¹) on total dry matter (TDM), grain yield and root length density (RLD) of maize at Yezin, Myanmar. 55

Table 4.8. Decline of nitrogen (N) and increase of phosphorus (P) concentrations in maize shoots in a cropping system field experiment at Yezin, Myanmar across five seasons (2002-2004). Data show treatments means; (n.a. not available). . 56

LIST OF FIGURE	Pages
Figure 3.1. Effects of phosphorus (P) application at 0 and 15 kg P ha ⁻¹ as triple superphosphate on total dry matter and grain yield of 20 green gram cultivars grown in a field experiment at Yezin, Myanmar, 2001. Cultivar ID are given in Table 3.1 Columns marked with different letters are significantly different at P < 0.05	26
Figure 3.2. Effects of phosphorus (P) application at 0 and 15 kg P ha ⁻¹ as triple superphosphate on total dry matter and grain yield of 15 green gram cultivars grown in a field experiment at Nyaung Oo, Myanmar, 2002. Cultivar ID are given in Table 3.1 Columns marked with different letters are significantly different at P < 0.05.	27
Figure 3.3. Root length density of 15 green gram cultivars grown in a field experiment at Nyaung Oo, Myanmar, 2002 (0 and 15 kg P ha ⁻¹ as TSP).....	28
Figure 3.4. Effect of phosphorus (P) application at 0 and 15 kg P ha ⁻¹ as triple superphosphate on total dry matter and grain yield of six green gram cultivars grown at Yezin, Myanmar, 2003.	29
Figure 3.5. Effect of phosphorus (P) application at 0 and 15 kg P ha ⁻¹ as triple superphosphate (TSP) on root length density of six green gram cultivars grown at Yezin, Myanmar, 2003.	29
Figure 4.1. Effects of manure and phosphorus (P) application on total dry matter of green gram (cv. V-3726) in a field trial in Myanmar (2002-2004). M0P0 = no manure, no mineral P; M0P1 = no manure, 15 kg P ha ⁻¹ ; M1P0 = 2 t ha ⁻¹ manure, no mineral P; M1P1 = 2 t ha ⁻¹ manure, 15 kg P ha ⁻¹ . The Lsd _{0.05} values across treatments are 1650, 1208, 720, 1175 and 480, in season 1-5, respectively. Columns within season marked with different letters are significantly different at P < 0.05.	44
Figure 4.2. Effects of manure and phosphorus (P) application on grain yield of green gram (cv. V-3726) in a field trial at Yezin, Myanmar (2002-2004). M0P0 = no manure, no mineral P; M0P1 = no manure, 15 kg P ha ⁻¹ ; M1P0 = 2 t ha ⁻¹ manure, no mineral P; M1P1 = 2 t ha ⁻¹ manure, 15 kg P ha ⁻¹ . The Lsd _{0.05} values are 201, 143, 199, 95 and 80, in season 1-5, respectively. Columns within seasons marked with different letters are significantly different at P < 0.05.	45
Figure 4.3. Effects of manure and phosphorus (P) application on root length density of greengram (cv. V-3726) in a field trial at Yezin, Myanmar (2002-2004). M0P0 = no manure, no mineral P; M0P1 = no manure, 15 kg P ha ⁻¹ ; M1P0 = 2 t ha ⁻¹ manure, no mineral P; M1P1 = 2 t ha ⁻¹ manure, 15 kg P ha ⁻¹ . The Lsd _{0.05} values are 1.88, 1.09, 0.64, 0.46 and 0.15, in season 1-5, respectively. Columns within seasons marked with different letters are significantly different at P < 0.05.	47
Figure 4.4. Effects of cropping system, manure and phosphorus (P) application on total dry matter of maize in a field trial at Yezin, Myanmar (2002-2004). C = continuous maize, R = maize-green gram rotation, M0P0 = no manure, no mineral P; M0P1 = no manure, 15 kg P ha ⁻¹ ; M1P0 = 2 t ha ⁻¹ manure, no mineral P; M1P1 = 2 t ha ⁻¹ manure, 15 kg P ha ⁻¹ . The Lsd _{0.05} values are 1040, 503, 643, 582 and 107, respectively. Columns within seasons marked with different letters are significantly different at P < 0.05.....	51
Figure 4.5. Effects of cropping system, manure and phosphorus (P) application on grain yield of maize in a field trial at Yezin, Myanmar (2002-2004). C = continuous maize, R = maize-green gram rotation, M0P0 = no manure, no mineral P; M0P1 = no manure, 15 kg P ha ⁻¹ ; M1P0 = 2 t ha ⁻¹ manure, no mineral P; M1P1 = 2 t ha ⁻¹ manure, 15 kg P ha ⁻¹ . The Lsd _{0.05} values are 450, 568, 743,	

- 477 and 106, respectively. Columns within seasons marked with different letters are significantly different at $P < 0.05$ 52
- Figure 4.6.** Effects of cropping system, manure and phosphorus (P) application on root length density maize in a field trial at Yezin, Myanmar (2002-2004). C = continuous maize, R = maize-green gram rotation, M0P0 = no manure, no mineral P; M0P1 = no manure, 15 kg P ha⁻¹; M1P0 = 2 t ha⁻¹ manure, no mineral P; M1P1 = 2 t ha⁻¹ manure, 15 kg P ha⁻¹. The $lsd_{0.05}$ values are 0.28, 0.65, 0.21, 0.39 and 0.29, respectively. Columns within seasons marked with different letters are significantly different at $P < 0.05$ 53
- Figure 4.7.** Effect of phosphorus (P) application (kg ha⁻¹) and legume residue management (removal *versus* incorporation) on grain yield of maize in the 5th season at Yezin, Myanmar (2004 dry season). The $lsd_{0.05}$ value was 42. Columns within seasons marked with different letters are significantly different at $P < 0.05$ 54
- Figure 4.8.** Effects of cropping system and manure application on mycorrhizal infection of maize in the 2004 rainy and dry season at Yezin, Myanmar (2004). M-M = continuous maize, M-G = maize-green gram rotation; R = legume straw removed; I = legume straw incorporated. The $lsd_{0.05}$ values 22.6 and 31.0, in season 4 and 5, respectively. Columns within seasons marked with different letters are significantly different at $P < 0.05$ 57
- Figure 4.9.** Effect of cropping system and manure application on plant-parasitic nematodes in soils of a maize cropping systems experiment in the 4th and 5th season at Yezin, Myanmar (2004). The data are back transformend after log-transformation for statistical analysis; $lsd_{0.05}$ values of transformed values are 0.26 and 0.64, respectively. Columns within season marked with different letters are significantly different at $P < 0.05$ 58
- Figure 4.10.** Soil mineral N dynamics (0-15 cm) in plots planted to continuous maize and plots under a green gram-maize rotation where legume residues had been either removed or incorporated before the onset of the maize season in 2004 at Yezin, Myanmar (5th dry season). 58

LISTS OF APPENDICES

Table 1. Analysis of variance table showing the effects of manure (M) and P application on three greengram cultivars with at Yezin, Myanmar (1 st rainy season, 2002).	69
Table 2. Analysis of variance table showing the effects of manure (M) and P application on three greengram cultivars with at Yezin, Myanmar (2 st rainy season, 2003).	69
Table 3. Analysis of variance table showing the effects of manure (M) and P application on three greengram cultivars with at Yezin, Myanmar (3 rd dry season, 2003).	70
Table 4. Analysis of variance table showing the effects of manure (M) and P application on three greengram cultivars with at Yezin, Myanmar (4 th rainy season, 2004).	70
Table 5. Analysis of variance table showing the effects of manure (M) and P application on three greengram cultivars with at Yezin, Myanmar (5 th dry season, 2004).	71
Table 6. Analysis of variance table for maize grown in a cropping system (CS) experiment with continuous maize and maize in rotation with greengram and manure (M) and P application at Yezin, Myanmar (2 nd rainy season, 2003).	71
Table 7. Analysis of variance table for maize grown in a cropping system (CS) experiment with continuous maize and maize in rotation with greengram and manure (M) and P application at Yezin, Myanmar (3 rd dry season, 2003).	72
Table 8. Analysis of variance table for maize grown in a cropping system (CS) experiment with continuous maize and maize in rotation with greengram and manure (M) and P application at Yezin, Myanmar (4 th rainy season, 2004).	72
Table 9. Analysis of variance table for maize grown in a cropping system (CS) experiment with continuous maize and maize in rotation with greengram and manure (M) and P application at Yezin, Myanmar (5 th dry season, 2004).	73
Table 10. Analysis of variance table for maize grown in a cropping system (CS) experiment with continuous maize and maize in rotation with greengram and manure (M), P application, and Legume management (LM) at Yezin, Myanmar (5 th dry season, 2004).	73
Table 11. Effects of manure (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on leaf area index of maize at Yezin, Myanmar (1 st rainy season, 2002). Means with different letters within columns are significantly different at $P < 0.05$	74
Table 12. Effects of cropping system (continuous maize <i>versus</i> green gram maize rotation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on leaf area index and ear number of maize at Yezin, Myanmar (2 nd rainy season, 2003). Means with different letters within columns are significantly different at $P < 0.05$	75
Table 13. Effects of cropping system (continuous maize <i>versus</i> green gram maize rotation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on leaf area index and ear number of maize at Yezin, Myanmar (3 rd dry season, 2003). Means with different letters within columns are significantly different at $P < 0.05$	76
Table 14. Effects of cropping system (continuous maize <i>versus</i> green gram maize rotation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on leaf area index and ear number of maize at Yezin, Myanmar (4 th rainy season, 2004). Means with different letters within columns are significantly different at $P < 0.05$	77

Table 15. Effects of cropping system (continuous maize <i>versus</i> green gram maize rotation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on leaf area index and ear number per plot of maize at Yezin, Myanmar (5 th dry season, 2004).	78
Table 16. Effects of cropping system (continuous maize <i>versus</i> green gram maize rotation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on plant height and stem diameter of maize at Yezin, Myanmar (2 nd rainy season, 2003).	79
Table 17. Effects of cropping system (continuous maize <i>versus</i> green gram maize rotation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on plant height and stem diameter of maize at Yezin, Myanmar (3 rd dry season, 2003).	80
Table 18. Effects of cropping system (continuous maize <i>versus</i> green gram maize rotation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on plant height of maize at Yezin, Myanmar (4 th rainy season, 2004). Means with different letters within columns are significantly different at $P < 0.05$	81
Table 19. Effects of cropping system (continuous maize <i>versus</i> green gram maize rotation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on plant height and stem diameter of maize at Yezin, Myanmar (5 th dry season, 2004).	82
Table 20. Treatment means and probabilities of treatment effects on nitrogen and phosphorus content of maize at Yezin, Myanmar (2 nd rainy season, 2003).	83
Table 21. Treatment means and probabilities of treatment effects on nitrogen and phosphorus content of maize at Yezin, Myanmar (3 rd dry season, 2003).	84
Table 22. Treatment means and probabilities of treatment effects on nitrogen and phosphorus content of maize at Yezin, Myanmar (4 th rainy season, 2004).	85
Table 23. Effects of legume residue management (removal <i>versus</i> incorporation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) application (0 and 15 kg P ha ⁻¹) on nitrogen and P concentration of maize shoots at Yezin, Myanmar (5 th dry season, 2004).	86
Table 24. Effects of cropping system (continuous maize <i>versus</i> green gram maize rotation), manure application (0 and 2000 kg ha ⁻¹) and banded phosphorus (P) fertilization (0 and 15 kg P ha ⁻¹) on nitrogen and concentration of maize shoots at Yezin, Myanmar (5 th dry season, 2004).	87
Figure 1. Effects of manure and phosphorus (P) application on grain yield of three green gram cultivars in a field trial at Yezin, Myanmar (2002). M0P0 = no manure, no mineral P; M0P1 = no manure, 15 kg P ha ⁻¹ ; M1P0 = 2 t ha ⁻¹ manure, no mineral P; M1P1 = 2 t ha ⁻¹ manure, 15 kg P ha ⁻¹	88
Figure 2. Effect of cropping system, manure and phosphorus (P) application (kg ha ⁻¹) on the grain yield of three green gram cultivars during the 2 nd (rainy) and 3 rd (dry) season in a field trial at Yezin in Myanmar (2003). M0P0 = no manure, no mineral P; M0P1 = no manure, 15 kg P ha ⁻¹ ; M1P0 = 2 t ha ⁻¹ manure, no mineral P; M1P1 = 2 t ha ⁻¹ manure, 15 kg P ha ⁻¹	88
Plate 1. Green gram protein experiment at Yezin University Farm, 2001.	89
Plate 2. Green gram protein experiment at Nyaung Oo Experimental Farm, 2002.	89
Plate 3. Maize-green gram rotation experiment at Yezin University Farm, 2003 rainy season.	90
Plate 4. Maize-green gram rotation experiment at Yezin University Farm, 2004 dry season.	90

Plate 5. Maize-green gram rotation experiment at Yezin University Farm, 2004 rainy season.....	91
--	----