

# CLINICAL BIOCHEMISTRY

Lecture Notes



Peter Rae  
Mike Crane  
Rebecca Pattenden

10th Edition

LN



WILEY Blackwell



# **Clinical Biochemistry**

## Lecture Notes



# Clinical Biochemistry

## Lecture Notes

---

### **Peter Rae**

BA, PhD, MBChB, FRCPE, FRCPath  
Consultant Clinical Biochemist  
Royal Infirmary of Edinburgh  
Honorary Senior Lecturer in Clinical Biochemistry  
University of Edinburgh

### **Mike Crane**

BSc, PhD, MSc, FRCPath  
Consultant Clinical Biochemist  
Royal Hospital for Sick Children & Royal Infirmary of Edinburgh  
Honorary Lecturer in Clinical Biochemistry  
University of Edinburgh

### **Rebecca Pattenden**

BSc, MSc, FRCPath  
Consultant Clinical Biochemist  
Western General Hospital, Edinburgh

**Tenth Edition**

**WILEY** Blackwell

This edition first published 2018 © 2018 by John Wiley & Sons Ltd

#### *Edition History*

John Wiley & Sons Ltd (9e, 2013)

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by law. Advice on how to obtain permission to reuse material from this title is available at <http://www.wiley.com/go/permissions>.

The right of Peter Rae, Mike Crane, Rebecca Pattenden to be identified as the authors of the editorial material in this work has been asserted in accordance with law.

#### *Registered Office(s)*

John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, USA

John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

#### *Editorial Office*

9600 Garsington Road, Oxford, OX4 2DQ, UK

For details of our global editorial offices, customer services, and more information about Wiley products visit us at [www.wiley.com](http://www.wiley.com).

Wiley also publishes its books in a variety of electronic formats and by print-on-demand. Some content that appears in standard print versions of this book may not be available in other formats.

#### *Limit of Liability/Disclaimer of Warranty*

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting scientific method, diagnosis, or treatment by physicians for any particular patient. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. While the publisher and authors have used their best efforts in preparing this work, they make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives, written sales materials or promotional statements for this work. The fact that an organization, website, or product is referred to in this work as a citation and/or potential source of further information does not mean that the publisher and authors endorse the information or services the organization, website, or product may provide or recommendations it may make. This work is sold with the understanding that the publisher is not engaged in rendering professional services. The advice and strategies contained herein may not be suitable for your situation. You should consult with a specialist where appropriate. Further, readers should be aware that websites listed in this work may have changed or disappeared between when this work was written and when it is read. Neither the publisher nor authors shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

#### *Library of Congress Cataloging-in-Publication Data*

Names: Rae, Peter, 1953– author. | Crane, Mike, 1979– author. |

Pattenden, Rebecca, 1974– author.

Title: Lecture notes. Clinical biochemistry/Peter Rae, Mike Crane, Rebecca Pattenden.

Other titles: Clinical biochemistry

Description: Tenth edition. | Hoboken, NJ : Wiley, 2018. | Preceded by Lecture notes.

Clinical biochemistry. 9th ed./Geoffrey Beckett ... [et al.]. 2013. |

Includes bibliographical references and index.

Identifiers: LCCN 2017013974 (print) | LCCN 2017014660 (ebook) | ISBN 9781119248699 (pdf) |

ISBN 9781119248637 (epub) | ISBN 9781119248682

Subjects: | MESH: Biochemical Phenomena | Clinical Laboratory Techniques |

Clinical Chemistry Tests | Pathology, Clinical-methods

Classification: LCC RB40 (ebook) | LCC RB40 (print) | NLM QU 34 | DDC 616.07/56–dc23

LC record available at <https://lccn.loc.gov/2017013974>

Cover design by Wiley

Cover image: © Miguel Malo/gettyimages

Set in 8.5/11pt Utopia by SPi Global, Pondicherry, India

# Contents

Preface, vi

List of abbreviations, vii

About the companion website, x

- 1 Requesting and interpreting tests, 1
- 2 Disturbances of water, sodium and potassium balance, 12
- 3 Acid–base balance and oxygen transport, 30
- 4 Renal disease, 43
- 5 Disorders of calcium, phosphate and magnesium metabolism, 60
- 6 Diabetes mellitus and hypoglycaemia, 77
- 7 Disorders of the hypothalamus and pituitary, 91
- 8 Abnormalities of thyroid function, 105
- 9 Disorders of the adrenal cortex and medulla, 121
- 10 Investigation of gonadal function infertility, menstrual irregularities and hirsutism, 139
- 11 Pregnancy and antenatal screening, 158
- 12 Cardiovascular disorders, 167
- 13 Liver disease, 183
- 14 Gastrointestinal tract disease, 198
- 15 Nutrition, 208
- 16 Inflammation, immunity and paraproteinaemia, 224
- 17 Malignancy and tumour markers, 234
- 18 Disorders of iron and porphyrin metabolism, 241
- 19 Uric acid, gout and purine metabolism, 252
- 20 Central nervous system and cerebrospinal fluid, 259
- 21 Therapeutic drug monitoring and chemical toxicology, 263
- 22 Clinical biochemistry in paediatrics and the elderly, 276

Index, 295

# Preface

This is the tenth edition of the book originally written by Professor Gordon Whitby, Dr Alistair Smith and Professor Iain Percy-Robb in 1975. It remains an Edinburgh-based book, but both the content and the authorship continue to evolve.

Ever since the first edition this book has been primarily aimed at medical students and junior doctors, but we also believe that it will be of value to specialist registrars, clinical scientists and biomedical scientists pursuing a career in clinical biochemistry and metabolic medicine, and studying for higher qualifications. It has continued to develop in line with changes that have both reshaped the undergraduate curriculum and taken place in medical practice.

Over the course of the book's existence changes in medical education have tended to reduce or abolish courses exclusively covering laboratory medicine disciplines, with their content being integrated into the relevant parts of a systems-based curriculum. This clearly places the laboratory disciplines at the heart of medical teaching in the diagnosis and management of patients, but risks losing the opportunity to take a closer view of the principles behind the use of diagnostic investigations. This book aims to focus on the choice and interpretation of investigations in the diagnosis and management of conditions where biochemical testing plays a key role, with a view to understanding not only their uses but also developing an appreciation of their limitations. This is underpinned by brief summaries of the relevant pathophysiology. There is an emphasis on commonly requested tests and commonly occurring pathology, but less common tests and disorders are also described.

We have reviewed and updated all chapters to ensure that they reflect current clinical practice, the availability of new tests, and where relevant the latest versions of national guidelines, with an emphasis on those published in the UK. Planning this new edition benefited from helpful feedback from a number of sources, including groups of both students and their teachers, commissioned by Wiley, and in response to this we have among other changes increased the numbers of diagrams and tables where these help to summarise useful information. We have also increased the numbers of clinical cases, as these remain a popular feature. Multiple choice questions with an explanation of the answers, and key learning points for each chapter are available as an on-line resource for revision.

Since the last edition, Geoff Beckett, Simon Walker and Peter Ashby have all retired. They were authors since the fourth, fifth and seventh editions, respectively, and have had an enormous effect on the development and success of this book. Their places have been ably taken by Mike Crane and Rebecca Pattenden, who have brought a fresh perspective to many of the topics covered. As ever, we are also indebted to a number of colleagues who read various chapters and provided valuable comment and advice, in particular Catriona Clarke and Jonathan Malo. We remain grateful for the continued interest and support provided by the staff at Wiley towards this title since its first appearance over forty years ago.

*Peter Rae  
Mike Crane  
Rebecca Pattenden*



# List of abbreviations

$\alpha$ -MSH	$\alpha$ -melanocyte stimulating hormone	CKD	chronic kidney disease
AAT	$\alpha_1$ -antitrypsin	CNS	central nervous system
ABP	androgen-binding protein	CoA	coenzyme A
A&E	accident and emergency	COC	combined oral contraceptive
ACE	angiotensin-converting enzyme	COHb	carboxyhaemoglobin
ACTH	adrenocorticotrophic hormone	CRH	corticotrophin-releasing hormone
ADH	antidiuretic hormone	CRP	C-reactive protein
AFP	$\alpha$ -foetoprotein	CSF	cerebrospinal fluid
AI	angiotensin I	CT	computed tomography
AIi	angiotensin II	CV	coefficient of variation
AIii	angiotensin III	DDAVP	1-deamino,8-D-arginine vasopressin
AIP	acute intermittent porphyria	DHEA	dehydroepiandrosterone
AIS	androgen insensitivity syndrome	DHEAS	dehydroepiandrosterone sulphate
ALA	aminolaevulinic acid	DHCC	dihydrocholecalciferol
ALP	alkaline phosphatase	DHT	dihydrotestosterone
ALT	alanine aminotransferase	DIT	di-iodotyrosine
AMA	anti-mitochondrial antibodies	DKA	diabetic ketoacidosis
AMH	anti-Mullerian hormone	DPP	4 dipeptidyl peptidase-4
AMP	adenosine 5-monophosphate	DSD	disorder of sexual differentiation
ANP	atrial natriuretic peptide	DVT	deep venous thrombosis
AST	aspartate aminotransferase	ECF	extracellular fluid
ATP	adenosine triphosphate	ECG	electrocardiogram/electrocardiography
AT	Pase adenosine triphosphatase	ED	erectile dysfunction
$\beta$ -LPH	$\beta$ -lipotrophin	EDTA	ethylenediamine tetraacetic acid
BChE	butylcholinesterase	eGFR	estimated glomerular filtration rate
BMI	body mass index	EPH	electrophoresis
BMR	basal metabolic rate	EPP	erythropoietic protoporphyria
BNP	B-type natriuretic peptide	ERCP	endoscopic retrograde cholangiopancreatography
CABG	coronary artery bypass grafting	ESR	erythrocyte sedimentation rate
CAH	congenital adrenal hyperplasia	FAD	flavin adenine dinucleotide
cAMP	cyclic adenosine monophosphate	FAI	free androgen index
CBG	cortisol-binding globulin	FBHH	familial benign hypocalcaemic hypercalcaemia
CCK	cholecystokinin	FIT	faecal immunochemical test
CCK-PZ	cholecystokinin-pancreozymin	FMN	flavin mononucleotide
CDT	carbohydrate-deficient transferrin	FOB	faecal occult blood
CEA	carcinoembryonic antigen	FPP	free protoporphyrin
CFT	calculated free testosterone	FSH	follicle-stimulating hormone
ChE	cholinesterase		
CK	creatin kinase		

FT3	free tri-iodothyronine	IFG	impaired fasting glucose
FT4	free thyroxine	Ig	immunoglobulin
GAD	glutamic acid decarboxylase	IGF	insulin-like growth factor
Gal-1-PUT	galactose-1-phosphate uridylyl-transferase	IGFBP	insulin-like growth factor-binding protein
GDM	gestational diabetes mellitus	IGT	impaired glucose tolerance
GFR	glomerular filtration rate	IM	intramuscular
GGT	$\gamma$ -glutamyltransferase	INR	international normalised ratio
GH	growth hormone	IV	intravenous
GHD	growth hormone deficiency	LCAT	lecithin cholesterol acyltransferase
GHRH	growth hormone-releasing hormone	LDH	lactate dehydrogenase
GI	gastrointestinal	LDL	low-density lipoprotein
GIP	glucose-dependent insulinotropic peptide/ gastric inhibitory polypeptide	LH	luteinising hormone
GLP-1	glucagon-like polypeptide-1	Lp(a)	lipoprotein (a)
GnRH	gonadotrophin-releasing hormone	LSD	lysergic acid diethylamide
GP	general practitioner	MCAD	medium chain acyl-CoA dehydrogenase
GSH	glucocorticoid-suppressible hyperaldosteronism	MCV	mean cell volume
GTT	glucose tolerance test	MDRD	Modification of Diet in Renal Disease
Hb	haemoglobin	MEGX	monoethylglycinexylidide
HC	hereditary coproporphyrin	MEN	multiple endocrine neoplasia
HCC	hydroxycholecalciferol	MGUS	monoclonal gammopathy of unknown significance
hCG	human chorionic gonadotrophin	MI	myocardial infarction
HDL	high-density lipoprotein	MIT	mono-iodotyrosine
HGPRT	hypoxanthine-guanine phosphoribosyltransferase	MODY	maturity onset diabetes of the young
HHS	hyperosmolar hyperglycaemic state	MOM	multiples of the median
5-HIAA	5-hydroxyindoleacetic acid	MRCP	magnetic resonance cholangiopancreatography
HIV	human immunodeficiency virus	MRI	magnetic resonance imaging
HLA	human leucocyte antigen	MTC	medullary thyroid cancer
HMG-CoA	$\beta$ -hydroxy- $\beta$ -methylglutaryl-coenzyme A	MUST	Malnutrition Universal Screening Tool
HMMA	4-hydroxy-3-methoxymandelic acid	NABQI	<i>N</i> -acetyl- <i>p</i> -benzoquinoneimine
HNF	hepatic nuclear factor	NAC	<i>N</i> -acetylcysteine
HPA	hypothalamic–pituitary–adrenal	NAD	nicotinamide–adenine dinucleotide
HPLC	high-performance liquid chromatography	NADP	NAD phosphate
HRT	hormone replacement therapy	NAFLD	nonalcoholic fatty liver disease
hsCRP	highly sensitive C-reactive protein	NASH	nonalcoholic steatohepatitis
5-HT	5-hydroxytryptamine	NICE	National Institute for Health and Clinical Excellence
5-HTP	5-hydroxytryptophan	NIPT	noninvasive prenatal testing
IBS	irritable bowel syndrome	NSAID	nonsteroidal anti-inflammatory agent
ICF	intracellular fluid	NTD	neural tube defect
ICU	intensive care unit	NTI	nonthyroidal illness
IDL	intermediate-density lipoprotein	OCP	oral contraceptive pill
IFCC	International Federation for Clinical Chemistry	OGTT	oral glucose tolerance test

PAPP-A	pregnancy-associated plasma protein A	SGLT	sodium-glucose cotransporter
PBG	porphobilinogen	SUR	sulphonylurea receptor
PCI	percutaneous coronary intervention	T3	tri-iodothyronine
PCOS	polycystic ovarian syndrome	T4	thyroxine
PCSK9	proprotein convertase subtilisin/kexin type 9	TBG	thyroxine-binding globulin
PCT	porphyria cutanea tarda	TDM	therapeutic drug monitoring
PE	pulmonary embolism	TDP	thiamin diphosphate
PEG	percutaneous endoscopic gastrostomy	TGN	6-thioguanine nucleotide
PEM	protein-energy malnutrition	THR	thyroid hormone resistance
PIIINP	pro-collagen type III	TIBC	total iron-binding capacity
PKU	phenylketonuria	TNF	tumour necrosis factor
PLP	pyridoxal 5'-phosphate	TPMT	thiopurine S-methyltransferase
POCT	point of care testing	TPN	total parenteral nutrition
POP	progestogen-only pill	TPOAb	thyroid peroxidase antibody
PRPP	5-phosphoribosyl-1-pyrophosphate	TPP	thiamin pyrophosphate
PSA	prostate-specific antigen	TRAb	thyrotrophin receptor antibody
PT	prothrombin time	TRH	thyrotrophin-releasing hormone
PTH	parathyroid hormone	TSH	thyroid-stimulating hormone
PTHrP	PTH-related protein	TSI	thyroid-stimulating immunoglobulin
RBP	retinol-binding protein	tTG	tissue transglutaminase
RDA	recommended dietary allowance	U&Es	urea and electrolytes
RF	rheumatoid factor	UFC	urinary free cortisol
RMI	risk of malignancy index	UV	ultraviolet
ROC	receiver operating characteristic	VIP	vasoactive intestinal peptide
SAAG	serum-ascites albumin gradient	VLDL	very low density lipoprotein
SAH	subarachnoid haemorrhage	VMA	vanillylmandelic acid
SD	standard deviation	VP	variegate porphyria
SHBG	sex hormone-binding globulin	WHO	World Health Organization
SIADH	inappropriate secretion of ADH	XO	xanthine oxidase
		ZPP	zinc protoporphyrin

# About the companion website

This book is accompanied by a companion website:



[www.lecturenoteseries.com/clinicalbiochemistry](http://www.lecturenoteseries.com/clinicalbiochemistry)

The website includes:

- Interactive multiple-choice questions
- Key revision points for each chapter

# Requesting and interpreting tests

---

## Learning objectives

To understand:

- ✓ how sample handling, analytical and biological factors can affect test results in health and disease and how these relate to the concept of a test reference range;
- ✓ the concepts of accuracy, precision, test sensitivity, test specificity in the quantitative assessment of test performance.

---

## Introduction

Biochemical tests are crucial to modern medicine. Most biochemical tests are carried out on blood using plasma or serum, but urine, cerebrospinal fluid (CSF), faeces, kidney stones, pleural fluid, etc. are sometimes required. Plasma is obtained by collecting blood into an anticoagulant and separating the fluid, plasma phase from the blood cells by centrifugation. Serum is the corresponding fluid phase when blood is allowed to clot. For many (but not all) biochemical tests on blood, it makes little difference whether plasma or serum is used.

There are many hundreds of tests available in clinical biochemistry but a core of common tests makes up the majority of tests requested. These core tests are typically available from most clinical laboratories throughout the 24-h period. Tests are sometimes brought together in profiles, especially when a group of tests provides better understanding of a problem than a single test (e.g. the liver function test profile). More specialist tests may be restricted to larger laboratories or specialist centres offering a national or regional service.

In dealing with the large number of routine test requests, the modern clinical biochemistry laboratory depends heavily on automated instrumentation linked to a laboratory computing system. Test results are assigned to electronic patient files that allow maintenance of a cumulative patient record. Increasingly, test requests can be electronically booked at the ward, clinic or in General Practice via a terminal linked to the main laboratory computer. Equally, the test results can be displayed on computer screens at distant locations, removing the need to issue printed reports.

In this first chapter, we set out some of the principles of requesting tests and of the interpretation of results. The effects of analytical errors and of physiological factors, as well as of disease, on test results are stressed. Biochemical testing in differential diagnosis and in screening is discussed.

---

## Collection of specimens

Test requests require unambiguous identification of the patient (patient's name, sex, date of birth and, increasingly, a unique patient identification number), together with the location, the name of the

requesting doctor and the date and time of sampling. Each test request must specify the analyses requested and provide details of the nature of the specimen itself and relevant clinical diagnostic information. This may be through a traditional request form and labelled specimen or be provided electronically in which case only the sample itself need be sent to the laboratory with its own unique identifier (typically a bar code which links it to the electronic request).

Clinical laboratories have multiple procedures at every step of sample processing to avoid errors. Regrettably, errors do occur and these arise at different stages between the sample being taken and the result being received:

- Pre-analytical. These arise prior to the actual test measurement and can happen at the clinical or laboratory end. Most errors fall into this category (see Table 1.1).
- Analytical. Laboratory based analytical errors are rare but may occur, e.g. reagent contamination, pipetting errors related to small sample volumes, computing errors.
- Post-analytical. These are increasingly rare because of electronic download of results from the analyser but include, for example, transcription errors when entering results from another laboratory into the

computer manually; results misheard when these are telephoned to the clinician.

Despite the scale of requesting of biochemical tests, errors are fortunately very rare. However, occasional blunders do arise and, if very unexpected results are obtained, it is important that the requesting doctor contacts the laboratory immediately to check whether the results are indeed correct or whether some problem may have arisen. Occasionally this reveals that more than one problem has occurred, for example two samples were labelled with each other's details on the ward, so querying the results can have wider benefits.

## The use of clinical biochemistry tests

Biochemical tests are most often *discretionary*, meaning that the test is requested for defined diagnostic purposes. The justification for discretionary testing is well summarised by Asher (1954):

- 1 Why do I request this test?
- 2 What will I look for in the result?
- 3 If I find what I am looking for, will it affect my diagnosis?

**Table 1.1 Some more common causes of pre-analytical errors arising from use of the laboratory.**

Error	Consequence
Crossover of addressograph labels between patients	This can lead to two patients each with the other's set of results. Where the patient is assigned a completely wrong set of results, it is important to investigate the problem in case there is a second patient with a corresponding wrong set of results.
Timing error	There are many examples where timing is important but not considered. Sending in a blood sample too early after the administration of a drug can lead to misleadingly high values in therapeutic monitoring. Interpretation of some tests (e.g. cortisol) is critically dependent on the time of day when the blood was sampled.
Sample collection tube error	For some tests the nature of the collection tube is critical, which is why the Biochemistry Laboratory specifies this detail. For example, using a plasma tube with lithium-heparin as the anti-coagulant is not appropriate for measurement of a therapeutic lithium level. Electrophoresis requires a serum sample rather than plasma so that fibrinogen does not interfere with the detection of any monoclonal bands. Topping up a biochemistry tube with a haematology (potassium ethylenediamine tetraacetic acid [EDTA]) sample will lead to high potassium and low calcium values in the biochemistry sample.
Sample taken from close to the site of an intravenous (IV) infusion	The blood sample will be diluted so that all the tests will be correspondingly low with the exception of those tests that might reflect the composition of the infusion fluid itself. For example, using normal saline as the infusing fluid would lead to a lowering of all test results, but with sodium and chloride results that are likely to be raised.

- 4 How will this investigation affect my management of the patient?
- 5 Will this investigation ultimately benefit the patient?

The main reasons for this type of testing are summarised in Table 1.2. Tests may also be used to help evaluate the future risk of disease (e.g. total cholesterol and HDL-cholesterol levels contribute to assessment of an individual's risk of cardiovascular disease), or in disease prognosis (e.g. biochemical tests to assess prognosis in acute pancreatitis or liver failure), or to screen for a disease, without there being any specific indication of its presence in the individual (e.g. maternal screening for foetal neural tube defects).

*Screening* may take several forms:

- In well-population screening a spectrum of tests is carried out on individuals from an apparently healthy population in an attempt to detect pre-symptomatic or early disease. It is easy to miss significant abnormalities in the large amount of data provided by the laboratory, even when the abnormalities are highlighted in some way. For these and other reasons, the value of well-population screening has been called into question and certainly should only be initiated under certain specific circumstances (Table 1.3).

**Table 1.2 Test selection for the purposes of discretionary testing.**

Category	Example
To confirm a diagnosis	Serum free T4 and thyroid-stimulating hormone (TSH) in suspected hyperthyroidism
To aid differential diagnosis	To distinguish between different forms of jaundice
To refine a diagnosis	Use of adrenocorticotrophic hormone (ACTH) to localise Cushing's syndrome
To assess the severity of disease	Serum creatinine or urea in renal disease
To monitor progress	Plasma glucose and serum K <sup>+</sup> to follow treatment of patients with diabetic ketoacidosis (DKA)
To detect complications or side effects	Alanine aminotransferase (ALT) measurements in patients treated with hepatotoxic drugs
To monitor therapy	Serum drug concentrations in patients treated with anti-epileptic drugs

**Table 1.3 Requirements for well-population screening.**

- The disease is common or life-threatening
- The tests are sensitive and specific
- The tests are readily applied and acceptable to the population to be screened
- Clinical, laboratory and other facilities are available for follow-up
- Economics of screening have been clarified and the implications accepted

**Table 1.4 Examples of tests used in case-finding programmes.**

Programmes to detect diseases in	Chemical investigations
Neonates	
Phenylketonuria (PKU)	Serum phenylalanine
Hypothyroidism	Serum TSH
Adolescents and young adults	
Substance abuse	Drug screen
Pregnancy	
Diabetes mellitus in the mother	Plasma glucose
Open neural tube defect (NTD) in the foetus	Maternal serum $\alpha$ -foetoprotein
Industry	
Industrial exposure to lead	Blood lead
Industrial exposure to pesticides	Serum cholinesterase activity
Elderly	
Malnutrition	Serum vitamin D levels
Thyroid dysfunction	Serum TSH and thyroxine

- In case-finding screening programmes appropriate tests are carried out on a population sample known to be at high risk of a particular disease. These are inherently more selective and yield a higher proportion of useful results (Table 1.4).

## Point of care testing (POCT)

These are tests conducted close to the patient, for example in the emergency department, an out-patient clinic, or a general practitioner's surgery.

**Table 1.5** Examples of POCT that are in common use.

Common POCT in blood	Common POCT in urine
Blood gases	Glucose
Glucose	Ketones
Urea and creatinine	Red cells/haemoglobin
Na, K and Ca	Bilirubin
Bilirubin	Protein
Alcohol	hCG

The instrumentation used is typically small and fits on a desk or may even be handheld. This approach can be helpful where there is a need to obtain a result quickly (e.g. blood gas results in the emergency department in a breathless patient), or where a result can be used to make a real-time clinical management decision (e.g. whether to adjust someone's statin dose on the basis of a cholesterol result). A further attraction is the immediate feedback of clinical information to the patient. POCT can be used to monitor illness by the individual patient and help identify if a change in treatment is needed (e.g. blood glucose monitoring in a diabetic patient). There is also an increasing number of urine test sticks that are sold for home use (e.g. pregnancy and ovulation testing by measuring human chorionic gonadotrophin (hCG) and luteinising hormone (LH), respectively). Table 1.5 shows examples of POCT tests in common use.

The introduction of POCT methodology requires attention to cost, ease of use, staff training, quality, health and safety as well as need. The advantages and disadvantages of POCT are summarised in Table 1.6.

## Interpretation of clinical biochemistry tests

Most reports issued by clinical biochemistry laboratories contain numerical measures of concentration or activity, expressed in the appropriate units. Typically, the result is interpreted in relation to a reference range (see Chapter 1: Reference ranges) for the analyte in question. Results within and outside the reference range may be subject to variation caused by a number of factors. These include analytical variation, normal biological variation, and the influence of pathological processes.

**Table 1.6** Advantages and disadvantages of point-of-care testing (POCT).

Advantages	Disadvantages
Rapid results on acutely ill patients	More expensive than centralised tests
Allows more frequent monitoring	Wide staff training may be needed
Immediate patient feedback	Nontrained users may have access, with potential for errors
Available 24 h if required	Calibration and quality control may be less robust
	Health and Safety may be less well monitored
	Results less often integrated into patient electronic record

## Sources of variation in test results

### Analytical sources of variation

Analytical results are subject to error, no matter how good the laboratory and no matter how skilled the analyst. The words "accuracy" and "precision" have carefully defined meanings in this context.


An *accurate* method will, on average, yield results close to the true value of what is being measured. It has no systematic bias. Lack of accuracy means that results will always tend to be either high or low.

A *precise* method yields results that are close to one another (but not necessarily close to the true value) on repeated analysis. If multiple measurements are made on one specimen, the spread of results will be small for a precise method and large for an imprecise one. Lack of precision means that results may be scattered, and unpredictably high or low.


A 'dashboard' analogy is often used to illustrate the different meanings of the terms accuracy and precision, and this is illustrated in Figure 1.1.

The standard deviation (SD) is the usual measure of scatter around a mean value. If the spread of results is wide, the SD is large, whereas if the spread is narrow, the SD is small. For data that have a Gaussian distribution, as is nearly always the case for analytical errors, the shape of the curve (Figure 1.2) is completely defined by the mean and the SD, and these characteristics are such that:





**Figure 1.1** The 'dartboard' analogy can be used to illustrate accuracy and precision.



**Figure 1.2** Diagram of a Gaussian (normal or symmetrical) distribution curve. The span (A) of the curve, the distance between the mean  $\pm 2$  SD, includes about 95% of the 'population'. The narrower span (B), the distance between the mean  $\pm 1$  SD, includes about 67% of the 'population'.

- About 67% of results lie in the range mean  $\pm 1$  SD.
- About 95% of results lie in the range mean  $\pm 2$  SD.
- Over 99% of results lie in the range mean  $\pm 3$  SD.

**Blunders** are grossly inaccurate results that bear no constant or predictable relationship to the true value. They arise, for instance, from mislabelling of specimens at the time of collection, or transcription errors when preparing or issuing reports (see Table 1.1).

If different results for the same test are obtained on two or more occasions on the same patient, then an

important question that arises is whether that difference is due to analytical imprecision or to a true change in the patient's clinical condition. Statistically, if the results differ by more than 2.8 times the analytical SD then there is a chance of over 95% that a genuine change in concentration of the substance has occurred.

## Biological causes of variation

Test results also show biological variation in both health and disease. The concentrations of all analytes in blood vary with time due to diverse physiological factors *within* the individual. There are also differences *between* individuals.

The following may be important causes of within-individual variation:

- 1 **Diet:** Variations in diet can affect the results of many tests, including serum triglyceride, the response to glucose tolerance tests and urinary calcium excretion.
- 2 **Time of day:** Several plasma constituents show diurnal variation (variation with the time of day), or a sleep/wake cycle. Examples include iron, adrenocorticotrophic hormone (ACTH) and cortisol concentrations.
- 3 **Posture:** Proteins and all protein-bound constituents of plasma show modest differences in concentration between blood collected from upright individuals and blood from recumbent individuals. Examples include serum calcium, cholesterol, cortisol and total thyroxine concentrations.
- 4 **Muscular exercise:** Recent exercise, especially if vigorous or unaccustomed, may increase serum creatine kinase (CK) activity and blood lactate, and lower blood pyruvate.
- 5 **Menstrual cycle:** Several substances show variation with the phase of the cycle. Examples include serum iron, and the serum concentrations of the pituitary gonadotrophins, ovarian steroids and their metabolites, as well as the amounts of these hormones and their metabolites excreted in the urine.
- 6 **Drugs:** These can have marked effects on chemical results. Attention should be drawn particularly to the many effects of oestrogen-containing oral contraceptives on serum constituents (Chapter 10: Steroid contraceptives).

Even after allowing for known physiological factors that may affect plasma constituents and for analytical imprecision, there is still considerable residual individual



### CASE 1.1

A 52-year-old man taking a statin drug to reduce his cholesterol level attended for a routine follow-up appointment. He was well, and had recently started training for a half-marathon as part of his determination to get fitter and reduce his risk of future cardiovascular problems. Statins sometimes cause muscle side effects, so among his other blood tests a creatine kinase (CK) was checked and it was very high. Should his statin be stopped?

**Comments:** Muscular exertion, especially if unaccustomed or severe, can give rise to high CK results. He was asked to refrain from training for a few days, and on repeat a CK level was normal.

variation (Table 1.7). The magnitude of this variation depends on the analyte, but it may be large and must be taken into account when interpreting successive values from a patient.

Differences between individuals can affect the concentrations of analytes in the blood. The following are the main examples:

- 1 **Age:** Examples include serum phosphate and alkaline phosphatase (ALP) activity, and serum and urinary concentrations of the gonadotrophins and sex hormones.
- 2 **Sex:** Examples include serum creatinine, iron and urate concentrations, and serum and urinary concentrations of the sex hormones.
- 3 **Race:** Racial differences have been described for serum cholesterol and protein. It may be difficult to distinguish racial from environmental factors, such as diet.

## Reference ranges

When looking at results, we need to compare each result with a set of results from a particular defined (or reference) population. This reference range is

determined, in practice, by measuring a set of reference values from a sample of that population, usually of healthy individuals. The nature of the reference population should be given whenever reference ranges are quoted, although a healthy population is usually assumed. Even age-matched and sex-matched reference ranges are often difficult to obtain, since fairly large numbers of individuals are needed.

When results of analyses for a reference population are analysed, they are invariably found to cluster around a central value, with a distribution that may be symmetrical (often Gaussian, Figure 1.3a) or asymmetrical (often log-Gaussian, Figure 1.3b). However, reference ranges can be calculated from these data without making any assumptions about the distribution of the data, using nonparametric methods.

Because of geographical, racial and other biological sources of variation between individuals, as well as differences in analytical methods, each laboratory should ideally define and publish its own reference ranges. By convention, these encompass the central 95% of the results obtained for each analysis from the reference population.


Analytical factors can affect the reference ranges for individual laboratories. If an inaccurate method is used, the reference range will reflect the method bias. If an imprecise method is used, the reference range will be widened, that is, the observed span of results (reflected in the SD) will be greater. In statistical terms, the observed variance (i.e. the square of the SD) of the population results will equal the sum of the true or biological variance of the population plus the analytical variance of the method.

## How do results vary in disease?

Biochemical test results do not exist in isolation, and when laboratory tests are requested, the clinician will often have made a list of differential diagnoses based on the patient's history, symptoms and signs, and may have a provisional diagnosis that is the likeliest possibility from this list

**Table 1.7** Residual individual variation of some serum constituents (expressed as the approximated day-to-day, within-individual coefficient of variation). CV = coefficient of variation.

Serum constituent	CV (%)	Serum constituent	CV (%)
Sodium	1	ALT activity	25
Calcium	1–2	AST activity	25
Potassium	5	Iron	25
Urea	10		



**Figure 1.3** Histograms showing the relative frequency with which results with the values indicated were obtained when serum  $\text{Na}^+$  and  $\gamma$ -glutamyltransferase (GGT) activities were measured in a reference population of healthy adult women. (a) The sodium data are symmetrically distributed about the mean whereas (b) the GGT data show a log-Gaussian distribution.

For example, in a patient with severe abdominal pain, tenderness and rigidity, there may be several diagnoses to consider including acute pancreatitis, perforated peptic ulcer and acute cholecystitis. In all three conditions, the serum amylase activity may be raised above the upper reference value for healthy adults. We need to know how the serum amylase activity might vary in the clinically likely diagnoses. It would be useful to know, for instance, that very high serum amylase activities are more likely to be associated with one of these diagnostic possibilities (pancreatitis), than with the other two.

To summarise, to interpret results on patients adequately, we need to know:

- the reference range for healthy individuals of the appropriate age range and sex;
- the values to be expected for patients with the disease, or diseases, under consideration;
- the prevalence of the disease, or diseases, in the population to which the patient belongs.

## The assessment of diagnostic tests – sensitivity and specificity

In evaluating and interpreting a test, it is necessary to know how it behaves in health and disease. Central to


this understanding are the terms “sensitivity” and “specificity.” These define how well a test performs in the diagnosis of a disease, but in order to calculate them it is necessary to know whether the disease is present or not by some method (this could be some other definitive test, or may be a diagnosis made later once the clinical course has made this more obvious).

- Test sensitivity refers to how effective the test is in detecting individuals who have the disease in question. It is expressed as the percentage of true positives in all the individuals who have disease (all the individuals with disease will encompass the true positives (TP) and false negatives (FN)). So:

$$\text{Sensitivity} = \text{TP} / (\text{TP} + \text{FN}) \times 100\%$$

- Test specificity is a measure of how good the test is at providing a negative result in the absence of disease. It is expressed as the percentage of true negatives in all those without the disease (all the individuals without disease will encompass the true negatives (TN) and the false positives (FP)). So:

$$\text{Specificity} = \text{TN} / (\text{TN} + \text{FP}) \times 100\%$$




**Figure 1.4** Diagrammatic representation of the distributions of results obtained with a test (a) that completely separates healthy people from people with a disease without any overlap between the distribution curves (i.e. an ideal test with 100% sensitivity and 100% specificity), and a test (b) that is less sensitive and less specific, in which there is an area of overlap between the distribution curves for healthy people and people with disease.

The ideal test is 100% sensitive (positive in all patients with the disease) and 100% specific (negative in all patients without the disease). We can illustrate this by means of the following hypothetical example shown diagrammatically in Figure 1.4a. This ideal is rarely achieved; there is usually overlap between the healthy and diseased populations (Figure 1.4b). In practice, we have to decide where to draw dividing lines that most effectively separate 'healthy' from 'diseased' groups, or disease A from disease B.

In evaluating tests for decision making, it is important to decide on the relative importance of sensitivity versus specificity in the context for which a test is used, and to compare the performance of different tests. In defining the presence or absence of a disease, a cut-off may be assigned to a test. Consider the situation where a high value for a particular test equates with the presence of a particular disease. A value above the cut-off would then define the presence of the disease and a value below the cut-off, the absence of disease. A cut-off set at a higher level will increase the test specificity at the expense of test sensitivity (more false negatives), while a cut-off set at a lower value will increase test sensitivity at the expense of test specificity (more false positives).

One way of comparing different tests and of visualising the balance between sensitivity and specificity at different test cut-offs is to plot the test sensitivity against specificity in a so-called 'receiver operating characteristic' (ROC) curve.



**Figure 1.5** Schematic representation of a receiver operating characteristic (ROC) plot. A random test produces a straight line set at 45° to the axes. A discriminatory, good test produces a graph with a steep slope from the origin, displaying high sensitivity at high specificity. Less discriminatory tests produce curves at intermediate positions, as shown. (Adapted from: Roulston, J.E. and Leonard, R.F.C. (1993). *Serological Tumour Markers: An Introduction*. Reproduced with permission from Elsevier.)

These curves will highlight which test is best suited to which requirement and will also help to define which cut-off to select in order to balance specificity versus sensitivity. This is illustrated in Figure 1.5.

## The assessment of diagnostic tests – positive and negative predictive values

The effectiveness of a test can also be defined in terms of the predictive value of a positive result and the predictive value of a negative result. These reflect the reality of clinical practice more than sensitivity and specificity do, in that the presence of a positive or negative test provides information about how likely a disease is to be present or not.

The positive predictive value is the proportion of the positive results that are true positives:

$$TP / (TP + FP) \times 100\%$$

A test with a high positive predictive value will, by definition, have few false positives. This would be important in a situation where a high number of false positives would otherwise lead to extensive and costly further investigation.

The negative predictive value is the proportion of the negative results that are true negatives:

$$TN / (TN + FN) \times 100\%$$

A test with a high negative predictive value would, by definition, have few false negatives. This would be particularly important, for example, in a test that was used for a screening programme where it is essential not to miss a case of the disease in question.

What is not immediately intuitive is that the predictive values depend not just on the sensitivity and specificity of a test but on the prevalence of the condition being tested for in the population of patients being tested.

To illustrate this, first consider screening neonates for an inherited metabolic disorder with an incidence of 1:5000; this is similar to that of some of the more common, treatable, inherited metabolic diseases such as phenylketonuria (PKU) or congenital hypothyroidism. Assume that we have a test with excellent performance, defined by sensitivity and specificity each of 99.5%, and that 1 000 000 neonates are tested (Table 1.8). The known incidence of the condition means that there will be 200 affected babies: 199 of these 200 babies will have a positive result (this is the sensitivity of 99.5%); 999 800 of the babies screened will not have the condition, and of these 994 801 will have a negative result (this is the specificity of 99.5%). However, there will be 4999 babies without the condition who receive a positive result, despite the apparently excellent specificity of the test, because the test

**Table 1.8 A hypothetical set of results of a screening test for a relatively common inherited metabolic disorder in neonates.**

Diagnostic category	Positive results	Negative results	Total
Disease present	199	1	200
Disease absent	4999	994 801	999 800
Total	5198	994 802	1 000 000
Predictive value	3.8%	100%	

Assumptions: sensitivity of the test 99.5%; specificity 99.5%; prevalence of the disorder 1:5000; 1 000 000 neonates screened. Note that the prevalence of PKU and of hypothyroidism in the UK is about 1:5000 live births, and that about 800 000 neonates in the UK are screened annually.

**Table 1.9 A hypothetical set of results of a test for a myocardial infarction.**

Diagnostic category	Positive results	Negative results	Total
Disease present	225	25	250
Disease absent	75	675	750
Total	300	700	1000
Predictive value	75%	96%	

Assumptions: sensitivity of the test 90%, specificity 90%, prevalence of the disorder 1:4; 1000 patients tested.

has been performed on so many babies. This means that the predictive value of a positive result is as low as 3.8%, while the predictive value of a negative result is virtually 100%. For every neonate affected by the disorder who has a positive test result, there will be about 25 (4999/199) neonates who also have a positive test but who do not have the disease.

Next consider a new test for myocardial infarction (MI) that has been shown to have slightly lower (but still acceptable) sensitivity and specificity of 95%. This is used to test patients presenting at an emergency department with severe central chest pain and it has previously been established in this department that 25% of such patients have suffered an MI (Table 1.9). The known incidence of MI in this population means that there will be 250 affected patients: 225 of these 250 patients will have a positive result (this is the sensitivity of 95%); 750 of the patients tested will not have had an MI and will be suffering from some other cause of chest pain, and of these 675 will have a negative result (this is the specificity of 95%). There will be 75 patients without an MI who receive a positive

result, meaning that the predictive value of a positive result is 75% (225 out of 300 patients), while the predictive value of a negative result is 96% (675 out of 700 patients). A negative result has virtually excluded the possibility of an MI, while a positive result has made an MI more than likely.

The second of these examples has shown a much higher positive predictive value (75%) than the first (3.8%) despite the lower sensitivity of the test used. This is because the condition being tested for was much more likely to be present. This provides an important lesson about how laboratory investigations can be used to make diagnoses, showing that tests perform better when a diagnosis is at least a likely possibility, and less well when a test is performed speculatively looking for an unlikely diagnosis.

Two important points regarding screening tests (those used to look for a condition in a basically healthy population) follow on from this. First, tests with very high sensitivity and with very low false-positive rates are required. Secondly, a heavy investigative load will result from the screening programme, since all the false positives will have to be followed up to determine whether or not they indicate the presence of disease. The traditional 95% reference range is not relevant to screening for rare conditions, since the rate of false positives would be far too high. The cut-off value has to be altered to decrease the false-positive rate, at the probable expense of missing some patients who have the condition for which screening is being carried out.

**Comment:** This is best examined by constructing a table as follows:

	Positive results	Negative results	Totals
Heart failure present	190 TP	10 FN	200
Heart failure absent	240 FP	560 TN	800
Total	430	570	1000

Because the test has a relatively high sensitivity, the table shows that it identifies the majority of patients with heart failure which is what is required in a test to rule out heart failure. Because the test lacks specificity, it can also be seen from the table that it identifies a considerable number of patients with positive results who do not have heart failure. In fact, the test is positive on more occasions in patients who do not have heart failure than in those with heart failure. Because other tests are available to the clinician, the false-positive patients can be separated from the true-positive patients on the basis of these further investigations. The 560 patients where the result is a true negative would then not need to go through more expensive further investigations. In this example, the test has been valuable in ruling out patients who would not require further investigation but ruling in those who would benefit. Clearly, it is not a perfect test but it would potentially prevent costly further investigations in a significant number of patients and, provided that the test itself is not too expensive, ultimately be worthy of consideration in terms of health economics.




### CASE 1.2

A new test is marketed which claims to diagnose heart failure. The test has a specificity of 70% and a sensitivity of 95% at the manufacturer's recommended cut-off for diagnosis. The Admissions Unit decides to use the test as part of an admission profile on breathless patients over the age of 65 years admitted for further assessment in order to exclude heart failure. Assuming a prevalence of 20% for heart failure in this population, calculate how many false negatives would be recorded after the first 1000 patients meeting the testing criteria had passed through the unit. Given that other tests can be used to establish a diagnosis of heart failure, do you think that the cut-off selected is sensible? (Prevalence figures are for illustrative purposes only.)

## Clinical audit in laboratory medicine

Clinical audit is a process that monitors the procedures involved in patient care to improve the delivery of the service. The principles are applicable to all clinical and investigational specialties (e.g. radiology), as well as laboratory-based specialties such as clinical biochemistry. For example, the monitoring of laboratory performance may identify a delay in analysing samples from the emergency department. This would precipitate a review of the way tests are requested, how samples are delivered to the department, the possible need for these samples to be prioritised in some way, and the way results are communicated back to the clinicians. Any necessary changes would



**Figure 1.6** The audit cycle.

be instituted, and the process re-monitored to ensure that the original problem had been overcome.

## The audit cycle

There is an essential sequence to auditing activities (Figure 1.6):

- 1 Identify an area of concern or interest, particularly if it is felt that there is room for improvement in the service, or if the same quality of service can be provided more economically.
- 2 Review and analyse the present procedures.
- 3 Identify specific aspects that might be capable of improvement.
- 4 Identify alternative procedures or standards that might lead to improvement.

- 5 Take the practical steps necessary to implement any changes proposed.
- 6 Compare the performance after the changes with that before them.

It must be emphasised that the final stage of analysis of the effects of any change is an integral part of the audit process; it is essential to know whether the measures taken have improved the service or made it more cost-effective. Sometimes, changes have no effect, or even have adverse effects.



## FURTHER READING

Asher, R. (1954) Straight and crooked thinking in medicine. *British Medical Journal* **2**, 460–2.



# Disturbances of water, sodium and potassium balance

---

## Learning objectives

To understand:

- ✓ the distribution of water,  $\text{Na}^+$  and  $\text{K}^+$  in the different fluid compartments of the body, and their control by hormonal and other factors;
- ✓ the clinical effects and management of different types of loss, retention or redistribution of fluid;
- ✓ the causes of hypernatraemia, hyponatraemia, hyperkalaemia and hypokalaemia, and what further investigations might be useful.

---

## Introduction

Fluid loss, retention or redistribution are common clinical problems in many areas of clinical practice. The management of these conditions is often urgent, and requires a rapid assessment of the history and examination, and of biochemical and other investigations. Both internal and external balance of these analytes must be considered. The internal balance is the distribution between different body compartments, while the external balance matches input with output.

plasma and gastrointestinal (GI) secretions, provide the potential for large losses, with consequent serious and rapid alterations in internal balance. For example, about 25 000 mmol of  $\text{Na}^+$  are filtered at the glomerulus over 24 h, normally with subsequent reabsorption of more than 99%. Likewise, 1000 mmol of  $\text{Na}^+$  enter the GI tract in various secretions each day, but less than 0.5% (5 mmol) is normally lost in the faeces.

## Internal distribution of water and sodium

In a 70 kg adult, total body water is about 42 L comprising about 28 L of intracellular fluid (ICF) and 14 L of extracellular fluid (ECF) water. ECF water is distributed as 3 L of plasma water and 11 L of interstitial water. Total body  $\text{Na}^+$  is about 4200 mmol and is mainly extracellular – about 50% is in the ECF, 40% in bone and 10% in the ICF.

---

## Water and sodium balance

The continuous movements of  $\text{Na}^+$  and water between plasma and glomerular filtrate, or between



Two important factors influence the distribution of fluid between the ICF and the intravascular and extravascular compartments of the ECF:

- **Osmolality:** This affects the movement of water across cell membranes.
- **Colloid osmotic pressure:** Together with hydrodynamic factors, this affects the movement of water and low molecular mass solutes (predominantly NaCl) between the intravascular and extravascular compartments.

## Osmolality and tonicity

The *osmolality* is the number of solute particles per unit weight of water, irrespective of the size or nature of the particles. Therefore, a given weight of low molecular weight solutes contributes much more to the osmolality than the same weight of high molecular weight solutes. The units are mmol/kg of water. This determines the osmotic pressure exerted by a solution across a membrane. Most laboratories can measure plasma osmolality, but it is also possible to calculate the approximate osmolality of plasma using a number of formulae of varying complexity. The following formula has the benefit of being easy to calculate and performs approximately as well as more complex versions (all concentrations must be in mmol/L):

Calculated osmolality

$$= 2[\text{Na}^+] + 2[\text{K}^+] + [\text{glucose}] + [\text{urea}]$$

This formula includes all the low molecular weight solutes that contribute to plasma osmolality. Values for  $\text{Na}^+$  and  $\text{K}^+$  are doubled in order to make allowance for their associated anions, such as chloride. The formula is approximate and is not a complete substitute for direct measurement. Calculated osmolality is usually close to measured osmolality, but they may differ considerably for two different types of reason:

- There may be a large amount of unmeasured low molecular mass solute (e.g. ethanol) present in plasma. This will contribute to the measured osmolality, but will obviously not be taken into account in the osmolality calculated from this formula. This will cause an 'osmole gap' with measured osmolality being greater than calculated osmolality.
- Alternatively, there may be a gross increase in plasma protein or lipid concentration, both of which decrease the plasma water per unit volume. This affects some methods of measurement of  $\text{Na}^+$ , giving an artefactually low result ('pseudohyponatraemia', see Chapter 2: Other causes of

hyponatraemia). This will result in an erroneously low calculated osmolality.

The osmolality of urine is usually measured directly, but is also linearly related to its specific gravity (which can be measured using urine dipsticks), unless there are significant amounts of glucose, protein or X-ray contrast media present.

**Tonicity** is a term often confused with osmolality. It relates to the osmotic pressure due to those solutes (e.g.  $\text{Na}^+$ ) that exert their effects across cell membranes, thereby causing movement of water into or out of the cells. Substances that can readily diffuse into cells down their concentration gradients (e.g. urea, alcohol) contribute to plasma osmolality but not to plasma tonicity, since after equilibration their concentration will be equal on both sides of the cell membrane. Tonicity is not readily measurable.

The tonicity of ICF and ECF equilibrate with one another by movement of water across cell membranes. An increase in ECF tonicity causes a reduction in ICF volume as water moves from the ICF to the ECF to equalise the tonicity of the two compartments, whereas a decrease in ECF tonicity causes an increase in ICF volume as water moves from the ECF to the ICF.



### CASE 2.1

A 45-year-old man was brought into the A&E department late at night in a comatose state. It was impossible to obtain a history from him, and clinical examination was difficult, but it was noted that he smelt strongly of alcohol. The following analyses were requested urgently.

Why is his measured osmolality so high?


Serum	Result	Reference ranges (adult male)
Urea	4.7	2.5–6.6 mmol/L
$\text{Na}^+$	137	135–145 mmol/L
$\text{K}^+$	4.3	3.6–5.0 mmol/L
Total $\text{CO}_2$	20	22–30 mmol/L
Glucose	4.2	mmol/L
Osmolality	465	280–296 mmol/kg

**Comments:** The osmolality can be calculated as 291.5, using the formula in Chapter 2: Osmolality and tonicity. The difference between this figure and the value for the directly measured osmolality (465 mmol/L) could be explained by the presence of another low molecular mass solute in plasma.

From the patient's history, it seemed that ethanol might be contributing significantly to the plasma osmolality, and plasma ethanol was measured the following day, on the residue of the specimen collected at the time of emergency admission. The result was 170 mmol/L, very close to the difference between the measured and calculated osmolalities.

## Colloid osmotic pressure (oncotic pressure)

The osmotic pressure exerted by plasma proteins across cell membranes is negligible compared with the osmotic pressure of a solution containing NaCl and other small molecules, since they are present in much lower molar concentrations. In contrast, small molecules diffuse freely across the capillary wall, and so are not osmotically active at this site, but plasma proteins do not readily do so. This means that plasma protein concentration and hydrodynamic factors together determine the distribution of water and solutes across the capillary wall, and hence between the intravascular and interstitial compartments (Figure 2.1).



**Figure 2.1** Movements of water and low molecular mass solutes across the capillary wall when the plasma protein concentration is (a) normal and (b) low. The effects shown are: hydrostatic pressure, which drives water and low molecular mass solutes *outwards* and decreases along the length of the capillary; and plasma oncotic pressure, which attracts water and low molecular mass solutes *inwards* and is constant along the length of the capillary. The net movement of water and low molecular mass solutes across the capillary wall is governed by the net effect of hydrostatic and plasma oncotic pressures.

## Regulation of external water balance

Typical daily intakes and outputs of water are given in Table 2.1. Water intake is largely a consequence of social habit and is very variable, but is also controlled by the sensation of thirst. Its output is controlled by the action of vasopressin, also known as antidiuretic hormone (ADH). In states of pure water deficiency, plasma tonicity increases, causing a sensation of thirst and stimulating vasopressin secretion, both mediated by hypothalamic osmoreceptors. Vasopressin then promotes water reabsorption in the distal nephron, with consequent production of small volumes of concentrated urine. Conversely, a large intake of water causes a fall in tonicity, suppresses thirst and reduces vasopressin secretion, leading to a diuresis, producing large volumes of dilute urine.


**Table 2.1** Average daily water intake and output of a normal adult in the UK.

Intake of water	mL	Output of water	mL
Water drunk	1500	Urine volume	1500
Water in food	750	Water content of faeces	50
Water from metabolism of food	250	Losses in expired air and insensible perspiration	950
Total	2500	Total	2500

Secretion of vasopressin is normally sensitively controlled by small changes in ECF tonicity, but it is also under tonic inhibitory control from baroreceptors in the left atrium and in the great vessels on the left side of the heart. When haemodynamic factors (e.g. excessive blood loss, heart failure) reduce the stretch on these receptors, often without an accompanying change in ECF tonicity, a reduction in tonic inhibitory control stimulates vasopressin secretion. The resulting water retention causes hyponatraemia, and is relatively ineffective in expanding the intravascular compartment, since water diffuses freely throughout all compartments (Figure 2.2).

## Regulation of external sodium balance

Dietary intakes of  $\text{Na}^+$  (and  $\text{Cl}^-$ ) are very variable worldwide. A typical 'Western' diet provides 100–200 mmol




**Figure 2.2** Different effects on the body's fluid compartments of fluid gains of 3 L of (a) water and (b) isotonic saline. The volumes shown relate to a 70 kg adult.

of both  $\text{Na}^+$  and  $\text{Cl}^-$  daily, but a normal total body  $\text{Na}^+$  can be maintained even if intake is less than 5 mmol or greater than 750 mmol daily. Urinary losses of  $\text{Na}^+$  normally closely match intake. There is normally little loss of these ions through the skin or in the faeces, but in disease the GI tract can become a major source of  $\text{Na}^+$  loss.

The amount of  $\text{Na}^+$  excreted in the urine controls the ECF volume since, when osmoregulation is normal, the amount of extracellular water is controlled to maintain a constant concentration of extracellular  $\text{Na}^+$ . A number of mechanisms are important regulators of  $\text{Na}^+$  excretion:

- **The renin-angiotensin-aldosterone system:** Renin is secreted in response to a fall in renal afferent arteriolar pressure or to a reduction in supply of  $\text{Na}^+$  to the distal tubule. It converts angiotensinogen in plasma to angiotensin I (AI), which in turn is converted to angiotensin II (AII) by angiotensin-converting enzyme (ACE). Both AII and its metabolic product angiotensin III (AIII) are physiologically active, and stimulate the release of aldosterone from the adrenal cortex. Aldosterone acts on the distal tubule to promote  $\text{Na}^+$  reabsorption in exchange for urinary loss of  $\text{H}^+$  or  $\text{K}^+$ . Since  $\text{Na}^+$  cannot enter cells freely, its retention (with iso-osmotically associated water) contributes solely to ECF volume expansion, unlike pure water retention (Figures 2.2 and 2.3). Although the renin-angiotensin-aldosterone system causes relatively slow responses to  $\text{Na}^+$  deprivation or  $\text{Na}^+$  loading, evidence suggests that this is the main regulatory mechanism for  $\text{Na}^+$  excretion.
- **The glomerular filtration rate (GFR):** The rate of  $\text{Na}^+$  excretion is often related to the GFR. When the GFR falls acutely, less  $\text{Na}^+$  is filtered and excreted, and



**Figure 2.3** Different effects on the body's fluid compartments of fluid losses of 3L of (a) water and (b) isotonic saline. The volumes shown relate to a 70kg adult.

vice versa. However, this only becomes a limiting factor in  $\text{Na}^+$  excretion at very low levels of GFR.

- **Atrial natriuretic peptide (ANP):** This peptide secreted by cardiocytes of the right atrium of the heart promotes  $\text{Na}^+$  excretion by the kidney, apparently by causing a marked increase in GFR. The importance of the ANP regulatory mechanism is not yet clear, but it probably only plays a minor role. Other structurally similar peptides have been identified, including brain or B-type natriuretic peptide (BNP), secreted by the cardiac ventricles and with similar properties to ANP. BNP is increasingly being used in the assessment of patients suspected of having cardiac failure (see Chapter 12: The diagnosis of heart failure).

## Disorders of water and sodium homeostasis

It is important to remember that the concentration of any substance is a consequence of the amount both of the solute (here  $\text{Na}^+$ ) and of the solvent (here water).

The concentration of the solute may change because of changes in either the amount of solute, the amount of solvent, or both. Although the physiological control mechanisms for water and for  $\text{Na}^+$  are distinct, they need to be considered together when seeking an understanding of a patient's  $\text{Na}^+$  and water balance, and of the plasma  $\text{Na}^+$  concentration.

Whereas losses or gains of pure water are distributed across all fluid compartments, losses or gains of  $\text{Na}^+$  and water, as isotonic fluid, are borne by the much smaller ECF compartment (Figures 2.2 and 2.3). Thus, it is usually more urgent to replace losses of isotonic fluid than losses of water. For the same reason, circulatory overload is more likely with excessive administration of isotonic  $\text{Na}^+$ -containing solutions than with isotonic dextrose (effectively water administration, since the dextrose is metabolised, leaving water).

Plasma  $\text{Na}^+$  concentration cannot be used as a simple measure of body  $\text{Na}^+$  status since it is very often abnormal as a result of losses or gains of water rather than of  $\text{Na}^+$ . The plasma  $\text{Na}^+$  concentration must be interpreted in relation to the patient's history and the

**Table 2.2 Causes of depletion of and excess water.**

Categories	Examples
Depletion of water	
• Inadequate intake	Infants, patients in coma or who are very sick, or have symptoms such as nausea or dysphagia
• Abnormal losses via	
Lungs	Inadequate humidification in mechanical ventilation
Skin	Fevers and in hot climates
Renal tract	Diabetes insipidus, lithium therapy
Excess water	
• Excessive intake	
Oral	Psychogenic polydipsia
Parenteral	Hypotonic infusions after operations
• Renal retention	Excess vasopressin (SIADH, Table 2.6), hypoadrenalism, hypothyroidism

findings on clinical examination, and if necessary backed up by other investigations.

The main causes of depletion and excess of water are summarised in Table 2.2, and of  $\text{Na}^+$  in Table 2.3. Although some of these conditions may be associated with abnormal plasma  $\text{Na}^+$  levels, it must be emphasised that this is not necessarily always the case. For example, patients with acute losses of isotonic fluid (e.g. plasma, ECF, blood) may be severely and dangerously hypovolaemic and  $\text{Na}^+$  depleted, and very possibly in shock, but their plasma  $\text{Na}^+$  concentration may nevertheless be normal or even raised.

## Hyponatraemia


Hyponatraemia is the most common clinical biochemical abnormality, occurring in up to 30% of hospitalised patients, with a wide spectrum of clinical symptoms from mild to life-threatening. It is usually primarily a disorder of water balance, with a relative excess of total body water. Most patients with hyponatraemia also have a low plasma osmolality. Unless an unusual cause of hyponatraemia is suspected (see Chapter 2: Other causes of hyponatraemia), measurement of plasma osmolality contributes little or no extra information.

**Table 2.3 Causes of depletion of and excess sodium.**

Categories	Examples
Depletion of sodium	
• Inadequate oral intake	Rare, by itself
• Abnormal losses via	
Skin	Excessive sweating, dermatitis, burns
GI tract	Vomiting, aspiration, diarrhoea, fistula, paralytic ileus, blood loss
Renal tract	Diuretic therapy, osmotic diuresis, renal tubular disease, mineralocorticoid deficiency
Excess of sodium	
• Excessive intake	
Oral	Sea water (drowning), salt tablets, hypertonic NaCl administration (this is rare)
Parenteral	Post-operatively, infusion of hypertonic NaCl
• Renal retention	Acute and chronic renal failure, primary and secondary hyperaldosteronism, Cushing's syndrome

Patients with hyponatraemia can be divided into three categories, on the basis of the ECF volume being low, normal or increased. These categories in turn reflect a total body  $\text{Na}^+$  that is low, normal or increased, respectively. The value of this classification is two-fold. First, the clinical history and examination often indicate the ECF volume and therefore the total body  $\text{Na}^+$  status. Secondly, treatment often depends on the total body  $\text{Na}^+$  status rather than the  $\text{Na}^+$  concentration. One possible way of narrowing the differential diagnosis of a patient with hyponatraemia, based on this subdivision, is shown in Figure 2.4.

It is also helpful to assess the rate at which the hyponatraemia is likely to have developed, and the severity of any symptoms, since these are important factors in determining treatment. Acute hyponatraemia develops over a period of less than 48h, with chronic hyponatraemia developing over a longer period. Neurological symptoms of hyponatraemia are due to the development of cerebral oedema (Table 2.4).



**Figure 2.4** Schematic diagram to assist in the diagnosis of some of the more common causes of hyponatraemia. In practice more than one cause may be present, and the findings may be influenced by the recent clinical history and oral or IV fluid intake. (GI, gastrointestinal; SIADH, syndrome of inappropriate secretion of antidiuretic hormone; 'third space' losses, leakage of fluid from intravascular to extravascular space, seen in bowel obstruction, pancreatitis, sepsis, etc.)

**Table 2.4 Neurological symptoms of hyponatraemia.**

Severity	Symptoms
Moderately severe	Nausea without vomiting, headache, confusion
Severe	Vomiting, cardio-respiratory distress, drowsiness, seizures, coma

Cerebral oedema is less likely to have developed in acute hyponatraemia, so treatment is aimed at preventing this from developing. In chronic hyponatraemia, rapid correction of a low sodium can result in osmotic demyelination syndrome, with consequences including dysarthria, dysphagia, spastic quadriparesis, seizures and death. Correction of hyponatraemia should therefore be performed at a controlled rate, in a setting where biochemical monitoring can be performed. It

**Table 2.5 Causes of hyponatraemia.**

ECF volume	Categories	Examples
Decreased total body Na <sup>+</sup> (loss of Na <sup>+</sup> >H <sub>2</sub> O)		
• Extrarenal losses of Na <sup>+</sup> (urine Na <sup>+</sup> <20 mmol/L)	GI tract Skin 'Internal'	Vomiting, diarrhoea Burns, severe dermatitis Paralytic ileus, peritoneal fluid
• Renal losses of Na <sup>+</sup> (urine Na <sup>+</sup> >20 mmol/L)	Diuretics Kidneys  Adrenals	Diuretic phase of renal tubular necrosis Mineralocorticoid deficiency
Normal or near-normal total body Na <sup>+</sup>	Acute conditions   Chronic conditions Anti-diuretic drugs Kidneys Adrenals Vasopressin excess Osmoregulator	Parenteral administration of water, after surgery or trauma, or during or after delivery   Opiates, chlorpropamide Chronic renal failure Glucocorticoid deficiency SIADH (Table 2.6) Low setting in carcinomatosis
Increased total body Na <sup>+</sup>	Acute conditions Chronic conditions	Acute renal failure Oedematous states (see Chapter 2: Hyponatraemia with increased ECF volume)

should be directed at the underlying cause where possible. Possible treatments include infusion of hypertonic saline, fluid restriction and stopping contributory drugs. The use of drugs such as demeclocycline or vasopressin antagonists is not usually recommended.

### Hyponatraemia with low ECF volume

The patient has lost Na<sup>+</sup> and water in one or more body fluids (e.g. GI tract secretions, urine, inflammatory exudate) or may have been treated with a diuretic (Table 2.5). The low ECF volume leads to tachycardia, orthostatic hypotension, reduced skin turgor and oliguria. The hypovolaemia causes secondary aldosteronism with a low urinary Na<sup>+</sup> concentration (usually <20 mmol/L), unless diuretic treatment is the cause, when urinary Na<sup>+</sup> remains high. The hypovolaemia also provides a 'volume stimulus' to vasopressin secretion, resulting in oliguria and a concentrated urine. The consequent water retention can further contribute to the hyponatraemia. Treatment is usually straightforward here and requires administration of isotonic saline.

### Hyponatraemia with normal ECF volume

The hyponatraemia results from excessive water retention, due to inability to excrete a water load. This may develop acutely, or it may be chronic (Table 2.5).

- *Acute water retention:* Plasma vasopressin is acutely increased after trauma or major surgery, as part of the metabolic response to trauma, and during delivery and post-partum. Administration of excessive amounts of water (e.g. as 5% dextrose) in these circumstances may exacerbate the hyponatraemia and cause acute water intoxication.
- *Chronic water retention:* Perhaps the most widely known chronic 'cause' of this form of hyponatraemia is dilutional hyponatraemia, often known as the syndrome of inappropriate secretion of ADH (SIADH) (Table 2.6). Whether this concept is of value in understanding its aetiology, or valid in terms of altered physiology, is uncertain. As the name implies, ADH (or rather vasopressin) is being secreted in the absence of an 'appropriate' physiological stimulus, of either fluid depletion or hypernatraemia. As water is



**Table 2.6 SIADH.**

Characteristics of the syndrome	Causes (and examples)
Low plasma Na <sup>+</sup> and osmolality	Malignant disease of the bronchus, prostate, pancreas, etc.
Inappropriately high urine osmolality	Chest diseases, e.g. pneumonia, bronchitis, tuberculosis
Excessive renal excretion of Na <sup>+</sup>	Central nervous system (CNS) diseases, including brain trauma, tumours, meningitis
No evidence of volume depletion	
No evidence of oedema	Drug treatment, e.g. carbamazepine, chlorpropamide, opiates
Normal renal and adrenal function	Miscellaneous conditions, including porphyria, psychosis, post-operative states

retained, the potential for expansion of the ECF volume is limited by a reduction in renin and an increase in sodium excretion. A new steady state is achieved, with essentially normal, or only mildly increased, ECF volume. If the causative disorder (Table 2.6) is transient, plasma Na<sup>+</sup> returns to normal when the primary disorder (e.g. pneumonia) is treated. However, in patients with cancer, the hyponatraemia is presumably due to production of vasopressin or a related substance by the tumour, and is usually persistent. If symptoms are mild, they may be treated by fluid restriction. Increased solute intake with urea or a combination of oral sodium chloride and a low dose loop diuretic can also be used in more profound hyponatraemia. Other causes of chronic retention of water include:

**CASE 2.2**

A 76-year-old man was making reasonable post-operative progress following major abdominal surgery for a carcinoma of the colon. Two days after the operation he appeared well, and there were no signs of dehydration or oedema. The following results were obtained:

What is the most likely cause of this man's low plasma Na<sup>+</sup> concentration?

Serum	Result	Reference ranges (adult male)
Urea	4.3	2.5–6.6 mmol/L
Na <sup>+</sup>	128	135–145 mmol/L
K <sup>+</sup>	4.3	3.6–5.0 mmol/L
Total CO <sub>2</sub>	25	22–30 mmol/L

**Comments:** Hyponatraemia is often seen in post-operative patients receiving IV fluids. At this time the ability to excrete water is reduced as part of the metabolic response to trauma. If excessive amounts of hypotonic fluids (usually 5% dextrose) are given, hyponatraemia will result. It may also be at least partly due to the 'sick cell syndrome'. There are usually no clinical features of water intoxication, and all that is required is review of the patient's fluid balance, and adjustment of the prescription for IV fluids. This man had received a total of 4.5L of fluid since his operation, and his fluid balance chart showed that he had a positive balance of 2L.

**CASE 2.3**

A 63-year-old coal miner had had a persistent chest infection, with cough and sputum, for the previous 2 months. He was a smoker. Clinical examination revealed finger clubbing, crackles and wheezes throughout the chest, and a small pleural effusion. There were no signs of dehydration or oedema.

Examination of blood and of a simultaneous urine specimen yielded the following results:

Serum	Result	Reference ranges (adult male)
Urea	2.3	2.5–6.6 mmol/L
Na <sup>+</sup>	118	135–145 mmol/L
K <sup>+</sup>	4.3	3.6–5.0 mmol/L
Total CO <sub>2</sub>	26	22–30 mmol/L
Osmolality	260	280–296 mmol/L
Urine	Result	
Na <sup>+</sup>	74	mmol/L
Osmolality	625	mmol/kg

What is the most likely cause of this man's low plasma Na<sup>+</sup> and osmolality?



**Comments:** This patient is not diluting his urine in response to low plasma osmolality and hyponatraemia: this suggests inappropriate ADH secretion. There tends to be a continuing natriuresis despite the hyponatraemia in these patients, as the retention of water leads to mild expansion of the ECF, and hence reduced aldosterone secretion. The presence of a dilutional hyponatraemia is also supported by the low plasma urea concentration. Before diagnosing the syndrome of inappropriate ADH secretion (SIADH – see Chapter 2: Hyponatraemia with normal ECF volume and Table 2.6), it is important to exclude adrenal, pituitary and renal disease. In this patient, possible explanations include the recurrent chest infections, and/or an underlying bronchogenic carcinoma, with ectopic secretion of ADH.

- **Chronic renal disease:** Damaged kidneys may be unable to concentrate or to dilute urine normally, tending to produce a urine of osmolality about that of plasma. Thus, the ability to excrete a water load is severely impaired, and excess water intake (oral or IV) readily produces a dilutional hyponatraemia. These patients may also be overloaded with  $\text{Na}^+$ .
- **Glucocorticoid deficiency:** Whether due to anterior pituitary disease or abrupt withdrawal of long-term glucocorticoid therapy, this may lead to inability to excrete a water load, and to hyponatraemia.
- **Resetting of the osmostat:** Some patients with malnutrition, carcinomatosis or tuberculosis seem to have their osmostat reset at a low level, with plasma  $\text{Na}^+$  of 125–130 mmol/L. The cause is uncertain.

## Hyponatraemia with increased ECF volume

Significant increases in total body  $\text{Na}^+$  give rise to clinically detectable oedema (Table 2.5). Generalised oedema is usually associated with secondary aldosteronism, caused by a reduction in renal blood flow, which stimulates renin production. Patients fall into at least three categories:

- **Renal failure:** Excess water intake in a poorly controlled patient with acute or chronic renal disease can lead to hyponatraemia with oedema.
- **Congestive cardiac failure:** In cardiac failure there is reduced renal perfusion and an 'apparent' volume deficit, and also increased venous pressure, with altered fluid distribution between the intravascular and interstitial compartments (Figure 2.1). These

lead to secondary aldosteronism and increased vasopressin secretion, causing  $\text{Na}^+$  overload and hyponatraemia.

- **Hypoproteinaemic states:** Low plasma protein, especially low albumin, leads to excessive losses of water and low molecular mass solutes from the intravascular to the interstitial compartments (Figure 2.1). Hence interstitial oedema is accompanied by reduced intravascular volume, with consequent secondary aldosteronism and stimulation of vasopressin release.

## 'Sick cell syndrome'

Some ill patients may have a hyponatraemia that is very resistant to treatment, and has no immediately obvious cause. The effective arterial plasma volume may be contracted, with a consequent secondary hyperaldosteronism and  $\text{Na}^+$  retention. The total ECF volume may in contrast be increased, possibly because of stress-induced vasopressin secretion, or other causes of SIADH. These, however, may not be the whole explanation for the observed pathophysiology, since plasma aldosterone and vasopressin concentrations are not always raised. The hyponatraemia may be due, at least in part, to the 'sick cell syndrome', in which there is an inability to maintain a  $\text{Na}^+$  gradient across the cell membrane, because of an increase in permeability, with or without impaired  $\text{Na}^+$  pump activity.

## Other causes of hyponatraemia

In all the examples of hyponatraemia discussed above, the low plasma  $\text{Na}^+$  occurs in association with reduced plasma osmolality. Where this is not the case, the following possibilities should be considered:

- **Artefact:** 'Hyponatraemia' is often caused by collection of a blood specimen from a vein close to a site at which fluid (typically 5% dextrose) is being administered intravenously.
- **Pseudohyponatraemia:** This is an artefactual result due to a reduction in plasma water caused by marked hyperlipidaemia or hyperproteinaemia (e.g. multiple myeloma). Normally, lipids and proteins make up a relatively small proportion, by volume, of plasma.  $\text{Na}^+$  and other electrolytes are dissolved in the plasma water, and do not enter the lipid or protein fraction of the plasma. This means that methods which measure  $\text{Na}^+$  concentration in the plasma water give similar results to those that measure  $\text{Na}^+$  concentration in total plasma. Most commonly used methods for measuring  $\text{Na}^+$  measure the amount of  $\text{Na}^+$  in a given volume of plasma. These methods include